

Date of publication: 09 December 2011

Implementation: To be determined by each Service

Change Notification UK National Blood Services No. 27 - 2011

Hepatitis C

Applies to the Tissue Donor Selection Guidelines Live Donors, Tissue Donor Selection Guidelines Deceased Donors, Bone Marrow and PBSC Donor Selection Guidelines and the Cord Blood Donor Selection Guidelines

1. Affected Individual

Obligatory

Must not donate.

Discretionary

If the individual has been told that he/she is HCV antibody negative, then samples should be taken to determine eligibility.

See if Relevant

Tissues Safety Entry

Additional Information

Hepatitis C is a serious viral infection that can lead to chronic liver disease, liver cancer (hepatoma) and chronic fatigue syndrome. It has also been linked with malignant lymphomas and autoimmune disease. The infection is very easily spread by transfusion.

Individuals who are chronically infected are sometimes referred to as 'carriers'. They often have no, or minimal, symptoms associated with their infection.

Many cases are linked to previous drug use and, before the introduction of HCV screening of blood donations, to transfusion.

Individuals who have had Hepatitis C infection in the past, and have been told that they have been successfully treated, will usually remain HCV antibody positive for many years. As a negative HCV antibody screening test is required before their donation can be issued, their tissues/cells cannot be used.

Reason for Change

'Additional Information' has been added.

\Continued

Sheila MacLennan

Dr Sheila MacLennan

Professional Director - Joint UKBTS/HPA Professional Advisory Committee

☎ Direct Dial: (0113) 820 8638

✉ sheila.maclennan@nhsbt.nhs.uk