

Whole Blood and Component Donor Selection Guidelines (WB-DSG)

Edition 203 - 01 June 2007

Release 75 - 17 February 2025

Issue 01

Introduction

These guidelines form a constituent part of Chapter 3 Care and selection of whole blood and component donors (including donors of pre-deposit autologous blood) of the Guidelines for the Blood Transfusion Services in the United Kingdom.

These criteria are reviewed regularly to ensure that the blood collected is of the highest quality and of sufficient quantity to meet the needs of the UKBTS.

JPAC is responsible for this document and JPAC receives professional advice from the Standing Advisory Committees that form part of its structure.

Users of these guidelines must ensure that they have the latest version and that recent changes have been implemented (usually within three months) by their national service.

Comments about the content of these guidelines, including notification of errors, omissions and suggestions for improvements, should be sent to the Chair of the Standing Advisory Committee for the Care and Selection of Donors (SACCSD):

Dr Angus Wells angus.wells@nhsbt.nhs.uk

Contents

Introduction	1
Document and Change Control	3
General Principles	4
Medication	6
Inspection of the Donor	7
Use of Alphabetical Listing	8
A-Z Index	9
A-Z	52
Updates	206
Appendix 1 - Estimated Blood Volume for Female donors (after Nadler) by height and weight	215
Appendix 2 - Table of Immunisations	216
Appendix 3 - Maximum permitted Extra Corporeal Volume for component donors	218
Appendix 4 - Management of post donation illness	219
Drugs and Platelet Donation	222

Document and Change Control

These guidelines are under the continuing review of the Standing Advisory Committees for the Care and Selection of Donors (SACCSD) and for Transfusion Transmitted Infection (SACTTI). This is to ensure that they are accurate and up to date. All changes have the approval of the UKBTS Joint Professional Advisory Committee (JPAC).

Change Notification.

A Change Notification Letter notifies changes to the **Medical Director** and the **Quality Manager** of each of the four national services. The **Professional Director of JPAC** is responsible for this notification. All changes will have the approval of the JPAC.

Implementation of changes is the responsibility of the individual Services.

Document version terminology.

A version shall be any of the following:

Extensive revisions of this document are known as '**Editions**'.

Changes following the issue of 'Change Notification Letters' are known as '**Releases**'.

Changes to the website, which do not involve a change to the medical or scientific content, are given an '**Issue**' number.

Edition Date, Release Date and Issue Date is the date on which an Edition, Release or Issue is first published on the UKBTS website.

Changes to off-line versions.

The **Quality Manager** of each Blood Service will effect changes. They will be informed when a new version is released. The **Quality Manager** is responsible for ensuring that there is an effective Version Control and Change Procedure in operation within their service to ensure that only up to date versions are in use and that all authorized copies, electronic and paper, are traceable.

Individual users of these guidelines are responsible for ensuring that they are using an up-to-date version.

Changes to the website versions.

The website will always display the up to date version. Any errors should be notified to JPACOffice@nhsbt.nhs.uk

General Principles

These guidelines apply to donors giving whole blood or blood components (red cells, platelets, plasma and granulocytes) for therapeutic use.

Donors are selected firstly to ensure that they do not come to harm from giving their donation and secondly to ensure that their donation is unlikely to harm any recipient. The ultimate responsibility for the selection of donors rests with the respective **National Medical Director**.

The immediate responsibility is with the **Qualified Healthcare Professional** in clinical charge of an individual donor session. When it is not clear from these guidelines if an individual donor is suitable, no donation should be taken until it has been discussed and agreed with a **Designated Clinical Support Officer**.

Only persons in good health should be accepted as donors. The prospective donor must be evaluated for their fitness to donate on the day by a suitably qualified person who has undergone appropriate training to use this document to select or defer donors. They must verify their assessment by signing the donation record.

Special note must be taken of the content of the **Blood Safety Entry** in the **A-Z Topics**.

It is the responsibility of session staff to ensure that donors clearly understand the nature of the donation process and the associated risks involved, as explained in the available literature. The donors must also understand the health check and other medical information presented to them. Donors are asked about confidential aspects of their medical history, hence great care must be taken over privacy and confidentiality. This means that third party interpreters can only be used as described in the A–Z Topic entry on **Communication Difficulties**.

Where there is separate guidance for **Whole Blood** and for **Component** donors, this is made clear. When there is a recognised risk to either the donor or the recipient, the guidelines **must** be followed.

The following terms may be used:

Also Known As

Lists alternative names for the topic entry.

Including

Lists any other terms which may be covered by the Guideline.

Definition

Where additional clarity is required, a definition is provided.

Obligatory

This will indicate how the donor **must** be dealt with by the use of several terms:

Must not donate

The donor **must not donate** if any of the statements apply to them, unless a **discretion** clearly applies. Often the deferral will depend on time related factors. If this is the case, the donor must be advised clearly when they will again become eligible to donate. If the deferral is not time limited (ie. it is likely to be permanent) the donor **must** be clearly advised why they cannot donate.

Refer to a Designated Clinical Support Officer

Is used when there is a need to seek further advice. The Designated Clinical Support Officer is a suitably trained person authorised to undertake this task by the National Medical Director or their nominated deputy.

Discretionary

Gives reasons why a donor may be permitted to donate. The statements are conditional. All statements that must be fulfilled come before the final statement that they may be accepted. If the donor fulfils these requirements, as well as all others that apply, then they can be accepted.

See

Means that the specified A-Z Topic entry **must** be consulted.

See if Relevant

Is used when an A-Z Topic entry may or may not need to be consulted, depending upon the information provided by the donor.

Additional Information

This provides background information as to why any particular action is required.

Information

This provides specific information as to the status of the guidance (e.g. required by the Blood Safety and Quality Regulations).

Update Information

The information here shows in which edition and release of the guidelines that this advice first appeared in its current form.

Reason for Change

This provides the background to any changes made to the entry since the last Edition or Release.

Some or all of these terms may be used under each subject heading or sub-heading.

Autologous Transfusion.

These guidelines do not apply to donors wishing to give their blood for Autologous Transfusions. Specific guidelines should be referred to eg Transfusion Medicine 1993, 3, 307-316.

Non-Therapeutic Donations.

Donors whose serum, plasma or cells will be used for laboratory, rather than therapeutic, purposes are generally subject to the same medical selection criteria. However, some decisions regarding their suitability to donate may be varied by a Designated Clinical Support Officer.

Therapeutic Venesection.

Patients referred for therapeutic venesection must not be accepted at donor sessions. The exception is donors with haemochromatosis. They may be accepted after referral to, and consideration by, a Designated Clinical Support Officer.

This section was last updated in WB-DSG Edition 203, Release 01

Medication

The underlying illness suffered by a donor, rather than the properties of any drug they are taking, is the usual reason for them not being eligible to donate.

In general, traces of drugs in donations are harmless to their recipients. However, donors treated with certain drugs are deferred for periods associated with the pharmacokinetic properties of the drug. Examples are some drugs used to treat acne, psoriasis, and some prostate problems. All such drugs have their own entry in the **A-Z Index**.

Drugs that can affect platelet function are listed in the **Drug Index** together with the deferral period required before a donor's blood can be used for platelet production.

This section was last updated in WB-DSG Edition 203, Release 01

Inspection of the Donor

All donors.

The donor should appear to be in good health. Intoxication, either by alcohol or drugs, should be a reason for not accepting a donor (see Addiction & Drug Abuse in the **A-Z** Topics).

A qualified clinical professional must assess disabled donors (see Disabled Donors in the **A-Z** Topics).

The skin at the venepuncture site should be free from disease.

This section was last updated in WB-DSG Edition 203, Release 01

Use of Alphabetical Listing

Any medical condition, or possible contraindication to donation, elicited at any point during the donation process, must be managed as indicated in the **A - Z** Topic section of these guidelines. Any collected material, which as a result is unsuitable for clinical use, **must** be clearly labelled as **unfit for use**.

If there is more than one contraindication to donation, any indicating the need to permanently defer the donor must be applied. This will mean that the donor is withdrawn from the donor panel. If withdrawal is not required, then the longest applicable deferral period must be applied.

Donors who undergo component donation procedures may be subject to additional or separate criteria compared to whole blood donors. Reference should be made to Chapter 3 Care and selection of whole blood and component donors (including donors of pre-deposit autologous blood) of the Guidelines for the Blood Transfusion Services in the United Kingdom, 8th Edition, 2013.

Any new health risks identified by this process should be notified to the Standing Advisory Committee on Care and Selection of Donors, so that they can be considered for incorporation into future revisions of these guidelines.

Donations must not be accepted from donors who exhibit health risks that are not listed in this guidance, without referral to, and acceptance by, the Designated Clinical Support Officer.

This section was last updated in WB-DSG Edition 203, Release 20 Issue 01

A-Z Index

A

Abdominoplasty	Surgery	182
Ablation for SVT	Arrhythmias	62
Abnormal Smear	Cervical Dysplasia	75
Abortion	Pregnancy	163
Abrasions Following a Punch	Non-Consented Exposure to Human Body Fluids	149
Abscess	Infection - Acute	127
Absences	Epilepsy	94
Accident	Accident	52
Acellular non human matrix graft recipient	Tissue and Organ Recipients	189
Acellular Porcine Collagen Matrix	Xenotransplantation	203
Achilles' Tendonitis	Tendonitis	185
Acid Indigestion	Acid Indigestion	52
Acid Reflux	Acid Indigestion	52
Acitretin - for Acne	Acne	53
Acitretin - for Psoriasis	Psoriasis	166
Acitretin - for Skin Disease	Skin Disease	174
Acne	Acne	53
Acne Rosacea	Acne	53
Acoustic Neuroma – Causing Vertigo	Vertigo	199
Acoustic Neuroma – Surgery	Neurosurgery	148
Acromegaly	Pituitary Disorders	157
ACTH Secreting Tumour	Pituitary Disorders	157
Acupuncture	Complementary Therapy	80
Acute Bronchitis	Infection - Acute	127
Acute glomerulonephritis	Kidney and Bladder Disease	133
Acute Intermittent Porphyrria	Porphyria	162
Acute nephritis	Kidney and Bladder Disease	133
Acute Porphyria	Porphyria	162
Acute tubular interstitial nephritis	Kidney and Bladder Disease	133
Addiction and Drug Abuse	Addiction and Drug Abuse	54
Addison's Disease	Adrenal Failure	54
Adenocarcinoma	Malignancy	141
ADHD	Neurobehavioral Disorders	148
Adrenal Failure	Adrenal Failure	54
Adrenocorticotrophic Hormone	Steroid Therapy	180
Afibrinogenaemia	Bleeding Disorder	65
African Trypanosomiasis	Infection - Chronic	128

For	See	Page
Age	Age	55
Age-Related Macular Degeneration	Eye Disease	95
AIDS - Infection	HIV	119
AIDS - Risk	Blood Safety Entry	68
Air Crew and Air Traffic Controllers	Air Crew and Air Traffic Controllers	55
Air Traffic Controller	Air Crew and Air Traffic Controllers	55
Air Travel	Travel	193
Alcohol	Addiction and Drug Abuse	54
Alitretinoin – for Dermatitis	Dermatitis	85
Alitretinoin – for Skin Disease	Skin Disease	174
Allergic Rhinitis	Allergy	56
Allergy	Allergy	56
Alopecia	Alopecia	57
Alpha Blocker - for Blood Pressure	Blood Pressure - High	67
Alpha Blockers - for Prostate Problems	Prostate Problems	165
Alpha One Antitrypsin Deficiency - Causing Breathing Problems	Respiratory Disease	170
Alpha One Antitrypsin Deficiency - Not Causing Problems	Inherited Diseases	131
Alpha-1 Antitrypsin Deficiency – causing cirrhosis	Liver Disease	137
Alprostadiil	Erectile Dysfunction	94
Alternative Therapies	Complementary Therapy	80
Alzheimer's Disease	Central Nervous System Disease	73
Ambulance Emergency Response Vehicle Driver	Hazardous Activity	108
AMD	Eye Disease	95
Amfebutamone®	Smoking Cessation	177
Anaemia	Anaemia	58
Anaesthetic	Anaesthetic	59
Anal Warts	Warts	201
Analgesics	Pain Killers	154
Anaphylaxis	Allergy	56
Androcur®	Anti-Androgens	60
Androgen Antagonists	Anti-Androgens	60
Aneurysm	Cardiovascular Disease	72
Angina	Cardiovascular Disease	72
Animal Bite (Non-Human)	Animal Bite (Non-Human)	59
Ankle - Injury	Accident	52
Ankylosing Spondylitis	Autoimmune Disease	63
Antabuse®	Addiction and Drug Abuse	54
Antacids	Acid Indigestion	52

For	See	Page
Antenatal Anti-D	Immunoglobulin Therapy	126
Anthrax - Contact With	Infectious Diseases - Contact With	130
Anthrax - Immunisation	Immunisation - Non-Live	125
Anthrax - Infection	Infection - Acute	127
Anti Smoking Treatments	Smoking Cessation	177
Anti-Androgens	Anti-Androgens	60
Antibiotic Therapy	Antibiotic Therapy	61
Anticoagulant Therapy	Anticoagulant Therapy	61
Anticonvulsant Therapy - for Chronic or Neuropathic Pain	Pain Killers	154
Anticonvulsant Therapy - for Epilepsy	Epilepsy	94
Anti-D antenatal intra muscular injections	Immunoglobulin Therapy	126
Anti-D post natal intra muscular injections	Immunoglobulin Therapy	126
Antidepressant Therapy - for Chronic or Neuropathic Pain	Pain Killers	154
Antidepressant Therapy - for Mental Health Problems	Mental Health Problems	143
Antifungals	Infection - General	129
Antihistamine Use - for Allergy	Allergy	56
Antihistamine Use - for Sedation	Hypnotics	123
Antihypertensives	Blood Pressure - High	67
Anti-Inflammatory Drugs - Nonsteroidal	Nonsteroidal Anti-Inflammatory Drugs	151
Anti-Inflammatory Drugs - Steroids	Steroid Therapy	180
Anti-Obesity Drugs	Donor Weight	90
Antiphospholipid Syndrome - as part of an Autoimmune Disease	Autoimmune Disease	63
Antiphospholipid Syndrome - Causing Infertility	Fertility	97
Antiphospholipid Syndrome - Causing Thrombosis	Thrombosis and Thrombophilia	186
Antithrombin III Deficiency	Thrombosis and Thrombophilia	186
Anti-Thyroid Tablets	Thyroid Disease	189
Antivirals	Infection - General	129
Antivirals - for Herpes	Herpes Simplex	118
Anxiety Disorders	Mental Health Problems	143
Anxiolytics	Mental Health Problems	143
Apheresis Donation Frequency	Frequency of Donation	99
Appendectomy	Surgery	182
Apraxia	Neurobehavioral Disorders	148
Arachnoid Cyst	Accept	52
Arnold – Chiari Malformation	Neurosurgery	148
Arrhythmias	Arrhythmias	62
Arterial Thrombosis	Thrombosis and Thrombophilia	186
Arthritis - due to Osteoarthritis (wear and tear)	Osteoarthritis	152

For	See	Page
Arthritis - due to Psoriasis	Psoriasis	166
Arthritis - type not known but taking Nonsteroidal Anti-Inflammatory Drugs	Nonsteroidal Anti-Inflammatory Drugs	151
Arthropod (Tick) Borne Encephalitis - Contact With	Non-Contagious Diseases - Contact With	150
Arthropod (Tick) Borne Encephalitis - Immunisation	Immunisation - Non-Live	125
Arthropod (Tick) Borne Encephalitis - Infection	Infection - Acute	127
Arthroscopy	Endoscopy	93
Asperger's Syndrome	Neurobehavioral Disorders	148
Aspirin - for Heart Problems	Cardiovascular Disease	72
Aspirin - for Pain or Inflammation	Nonsteroidal Anti-Inflammatory Drugs	151
Asthma	Asthma	63
Athlete's Foot	Skin Disease	174
Atrial Fibrillation	Arrhythmias	62
Atrial Flutter	Arrhythmias	62
Atrial Tachycardia	Arrhythmias	62
Atrioventricular Nodal Reentrant Tachycardia	Arrhythmias	62
Attention Deficit Hyperactivity Disorder	Neurobehavioral Disorders	148
Auditory Processing Disorder	Neurobehavioral Disorders	148
Autism Spectrum Disorder	Neurobehavioral Disorders	148
Autistic	Neurobehavioral Disorders	148
Autoimmune Disease	Autoimmune Disease	63
Autologous matrix implant recipient	Tissue and Organ Recipients	189
Autologous Platelet Rich Plasma	Transfusion	191
Autologous tissue recipient	Tissue and Organ Recipients	189
Autologous Transfusion	Transfusion	191
A-V Block	Arrhythmias	62
A-V Tachycardia	Arrhythmias	62
Avastin®	Eye Disease	95
AVNRT	Arrhythmias	62
Avodart® - for Hair Loss	Alopecia	57
Avodart® - for Prostate Problems	Anti-Androgens	60
Awaiting Test Results	Investigations	132
Axillary Vein Thrombosis	Thrombosis and Thrombophilia	186
B		
B12 Deficiency	Anaemia	58
Babesiosis	Infection - Chronic	128
Baby	Pregnancy	163
Baby by Caesarean	Pregnancy	163
Bacillus Calmette-Guerin Immunisation	Immunisation - Live	125
Back Pain	Back Problems	64

For	See	Page
Back Problems	Back Problems	64
Baldness	Alopecia	57
Barrett's Oesophagus	Malignancy	141
Basal Cell Carcinoma	Malignancy	141
BBB	Cardiovascular Disease	72
BCG Immunisation	Immunisation - Live	125
Beconase® Nasal Spray	Allergy	56
Behcet's Syndrome	Autoimmune Disease	63
Bell's Palsy	Central Nervous System Disease	73
Benign Intracranial Hypertension	Central Nervous System Disease	73
Benign Prostatic Hypertrophy	Prostate Problems	165
Berry Aneurysm	Cerebrovascular Disease and Intracranial Haemorrhage	74
Beta Blocker - for Blood Pressure	Blood Pressure - High	67
Beta Blockers - for Heart Problems	Cardiovascular Disease	72
Beta Blockers - for Thyroid Disease	Thyroid Disease	189
Bevacizumab	Eye Disease	95
Bicalutamide	Anti-Androgens	60
Bilharzia	Infection - Acute	127
Biological Modalities and Therapies	Monoclonal antibody therapy and other Biological Modalities	144
Bipolar Disorder	Mental Health Problems	143
Birth by Caesarean	Pregnancy	163
Bisphosphonates - for Malignancy	Malignancy	141
Bisphosphonates - for Osteoporosis	Osteopenia	153
Bisphosphonates - for Paget's Disease	Paget's Disease of Bone	154
Bite - Animal	Animal Bite (Non-Human)	59
Bite - Human	Non-Consented Exposure to Human Body Fluids	149
Bite - Snake	Animal Bite (Non-Human)	59
Bladder - Catheter	Urinary Catheterisation	198
Bladder Infection	Infection - General	129
Bladder problems	Kidney and Bladder Disease	133
Bladder Surgery	Surgery	182
Bleeding Disorder	Bleeding Disorder	65
Blind Donor	Disabled Donor	88
Blood Clot	Thrombosis and Thrombophilia	186
Blood Derivatives	Transfusion	191
Blood Donation Frequency	Frequency of Donation	99
Blood Exposure - by Transfusion	Transfusion	191
Blood Exposure - Splash or Inoculation	Non-Consented Exposure to Human Body Fluids	149

For	See	Page
Blood Pressure - High	Blood Pressure - High	67
Blood Pressure - Low	Blood Pressure - Low	68
Blood Pressure Tablets	Blood Pressure - High	67
Blood Products	Transfusion	191
Blood Safety Entry	Blood Safety Entry	68
Blood Splash	Non-Consented Exposure to Human Body Fluids	149
Blood Thinners	Anticoagulant Therapy	61
Blood Transfusion	Transfusion	191
Body Building Drugs	Addiction and Drug Abuse	54
Body Piercing	Body Piercing	71
Boil	Infection - Acute	127
Bone Cancer	Malignancy	141
Bone Graft	Tissue and Organ Recipients	189
Bone Implanted Hearing Aid	Surgery	182
Borrelioses	Infection - Acute	127
Botox Injection	Complementary Therapy	80
Botulism - Immunisation	Immunisation - Non-Live	125
Bowel Cancer	Malignancy	141
Bowen's Disease	Malignancy	141
Bradycardia	Arrhythmias	62
Brain Cyst	Accept	52
Brain Surgery	Neurosurgery	148
Brain Tumour	Central Nervous System Disease	73
Breast Augmentation	Surgery	182
Breast Biopsy	Surgery	182
Breast Cancer	Malignancy	141
Breast Enlargement	Surgery	182
Breast Implant	Surgery	182
Breast Lump - Awaiting Investigation	Investigations	132
Breast Lump - Following Surgery	Surgery	182
Breast Reduction	Surgery	182
Breastfeeding	Breastfeeding	71
Broken Bone	Accident	52
Bromocriptine for Central Nervous System Disease	Central Nervous System Disease	73
Bromocriptine for Pituitary Disorders	Pituitary Disorders	157
Bronchiectasis	Respiratory Disease	170
Bronchitis - Acute	Infection - Acute	127
Bronchitis - Chronic	Respiratory Disease	170

For	See	Page
Bronchoscopy	Endoscopy	93
Brucella, contact with	Non-Contagious Diseases - Contact With	150
Brucellosis	Infection - Chronic	128
Bruising, Excessive	Bleeding Disorder	65
Budd Chiari Syndrome	Surgery	182
Bundle Branch Block	Cardiovascular Disease	72
Bunion Operation	Surgery	182
Bupropion	Smoking Cessation	177
Burns	Wounds, Mouth and Skin Ulcers	203
Burr Hole Surgery	Neurosurgery	148
Bursitis	Tendonitis	185
Bus Driver	Hazardous Activity	108
Byetta®	Diabetes Mellitus	86
Bypass Operation - Blocked Artery	Cardiovascular Disease	72
Bypass Operation - Other	Surgery	182
C		
Caesarean Birth	Pregnancy	163
Caisson Disease	Decompression Illness	83
Cancer	Malignancy	141
Candida	Thrush	188
Cannabis	Addiction and Drug Abuse	54
Cardiac Surgery	Cardiac Surgery	71
Cardiomyopathy	Cardiovascular Disease	72
Cardiovascular Disease	Cardiovascular Disease	72
Carotid Artery Disease	Cerebrovascular Disease and Intracranial Haemorrhage	74
Carpal Tunnel Syndrome Operation	Tendonitis	185
Cartilage Operation	Surgery	182
Casodex®	Anti-Androgens	60
Cataracts	Eye Disease	95
Catarrh	Catarrh	73
Catarrh - Acute	Infection - Acute	127
Catheter - Urinary	Urinary Catheterisation	198
Cauda Equina Syndrome	Neurosurgery	148
Caverject®	Erectile Dysfunction	94
Cellulitis	Infection - General	129
Central Nervous System Disease	Central Nervous System Disease	73
Cerebral Embolism	Cerebrovascular Disease and Intracranial Haemorrhage	74
Cerebral Palsy	Disabled Donor	88
Cerebral Shunt	Shunts, Stents and Devices	173

For	See	Page
Cerebrovascular Accident	Cerebrovascular Disease and Intracranial Haemorrhage	74
Cerebrovascular Disease and Intracranial Haemorrhage	Cerebrovascular Disease and Intracranial Haemorrhage	74
Cervarix®	Immunisation - Non-Live	125
Cervical Cone Biopsy	Cervical Dysplasia	75
Cervical Dysplasia	Cervical Dysplasia	75
Cervical Intraepithelial Neoplasia	Cervical Dysplasia	75
CFS	Chronic Fatigue Syndrome	77
Chagas' Disease	South American Trypanosomiasis	177
Chagas' Risk	South American Trypanosomiasis	177
Champix®	Smoking Cessation	177
Charcot Marie Tooth	Disabled Donor	88
Chemotherapy - Antibiotics	Antibiotic Therapy	61
Chemotherapy - for Autoimmune Disease	Autoimmune Disease	63
Chemotherapy - for Malignancy	Malignancy	141
Chest Infection	Infection - General	129
Chest Pain	Chest Pain	76
Chiari Malformation	Neurosurgery	148
Chicken Pox - Contact With	Infectious Diseases - Contact With	130
Chicken Pox - Immunisation	Immunisation - Live	125
Chicken Pox - Infection	Infection - Acute	127
Chik V	Tropical Viruses	194
Chikungunya Virus	Tropical Viruses	194
Chiropody	Chiropody	76
Chlamydia - Lymphogranuloma Venereum	Infection - Chronic	128
Chlamydia - Other	Infection - Acute	127
Cholecystectomy	Gall Bladder Disease	100
Cholecystitis	Gall Bladder Disease	100
Cholera - Contact With	Non-Contagious Diseases - Contact With	150
Cholera - Immunisation	Immunisation - Non-Live	125
Cholera - Infection	Infection - Acute	127
Cholesteatoma	Surgery	182
Cholesterol - High	Hypercholesterolaemia	122
Chondrocyte Transplant	Tissue and Organ Recipients	189
Chondromalacia	Chondromalacia	76
Christmas Disease	Bleeding Disorder	65
Chronic Alcohol Problems	Addiction and Drug Abuse	54
Chronic Bronchitis	Respiratory Disease	170
Chronic Fatigue Syndrome	Chronic Fatigue Syndrome	77

For	See	Page
Chronic Myeloid Leukaemia	Malignancy	141
Chronic Nephritis	Kidney and Bladder Disease	133
Chronic Venous Insufficiency	Varicose Veins and Chronic Venous Insufficiency	199
CIN	Cervical Dysplasia	75
Cirrhosis	Liver Disease	137
CIS	Malignancy	141
CJD	Prion Associated Diseases	164
Clamelle®	Antibiotic Therapy	61
Climbing	Hazardous Activity	108
Climbing Ladders	Hazardous Activity	108
Climbing Scaffolding	Hazardous Activity	108
Clinical Trials	Clinical Trials	77
Clomid®	Fertility	97
Clomiphene	Fertility	97
Clonidine - for Blood Pressure	Blood Pressure - High	67
Clonidine - for Menopausal Flushing	Menopause	143
Clonidine - for Migraine	Migraine	143
Clopidogrel	Clopidogrel	78
Clotting Factor Treatment	Bleeding Disorder	65
Clotting Factors	Bleeding Disorder	65
CNS Disease	Central Nervous System Disease	73
Coagulation Factor Concentrates	Bleeding Disorder	65
Coastguard Emergency Response Vehicle Driver	Hazardous Activity	108
Cocaine	Addiction and Drug Abuse	54
Cochlear Implant	Surgery	182
Co-Cyprindiol	Acne	53
Cod Liver Oil	Vitamins and Other Nutritional Supplements	200
Coeliac Disease	Coeliac Disease	78
Cold - Contact With	Infectious Diseases - Contact With	130
Cold - Infection	Infection - Acute	127
Cold Sore	Herpes Simplex	118
Colitis - Infection	Infection - General	129
Colitis - Inflammatory Bowel Disease	Inflammatory Bowel Disease	131
Collagen Injections	Complementary Therapy	80
Colloid Cyst	Accept	52
Colonic Hydrotherapy	Complementary Therapy	80
Colonic Irrigation	Complementary Therapy	80
Colonoscopy	Endoscopy	93

For	See	Page
Colostomy	Stoma	181
Communication Difficulties	Communication Difficulties	78
Complementary Therapy	Complementary Therapy	80
Component Donation Frequency	Frequency of Donation	99
Conceiving - Difficulty in	Fertility	97
Concussion	Accident	52
Cone Biopsy	Cervical Dysplasia	75
Congenital Erythropoietic Porphyria	Porphyria	162
Conjunctivitis	Eye Disease	95
Conn's syndrome	Conn's Syndrome	81
Conn's Syndrome	Conn's Syndrome	81
Contact With Infectious Disease	Infectious Diseases - Contact With	130
Contagious Pustular Dermatitis - Contact With	Infectious Diseases - Contact With	130
Contagious Pustular Dermatitis - Infection	Infection - Acute	127
Contraceptive Use	Contraceptive Use	81
Corneal Graft	Eye Disease	95
Corneal Reshaping - by Laser	Laser Treatment	136
Coronary Artery Bypass Surgery	Cardiovascular Disease	72
Coronary Thrombosis	Cardiovascular Disease	72
Coronavirus Contact	Coronavirus Infection (COVID-19)	81
Coronavirus Infection (COVID-19)	Coronavirus Infection (COVID-19)	81
Cortisone Treatment	Steroid Therapy	180
Cosmetic Fillers	Complementary Therapy	80
Cosmetic Laser Treatment	Laser Treatment	136
Cosmetic Surgery	Surgery	182
CoV	Coronavirus Infection (COVID-19)	81
COVID-19	Coronavirus Infection (COVID-19)	81
CPAP	Sleep Apnoea	175
Crane Operator	Hazardous Activity	108
Creutzfeldt-Jakob Disease	Prion Associated Diseases	164
Crimean-Congo Fever	Viral Haemorrhagic Fever	200
Crimean-Congo Fever – Contact With	Viral Haemorrhagic Fever	200
Crohn's Disease	Inflammatory Bowel Disease	131
Cross-sex Hormone Therapy	Hormone Replacement and Sex Hormone Therapy	120
Cupping	Complementary Therapy	80
Cushing's Disease of Pituitary Cause	Pituitary Disorders	157
CVA	Cerebrovascular Disease and Intracranial Haemorrhage	74
Cyprostat®	Anti-Androgens	60

For	See	Page
Cyproterone Acetate	Anti-Androgens	60
Cystic Fibrosis	Respiratory Disease	170
Cystitis	Infection - General	129
Cystostomy	Stoma	181
Cysts - Removal of	Surgery	182
Cytomegalovirus	Infection - General	129
D		
D&C	Surgery	182
D&V	Gastroenteritis	101
Dangers at Work	Hazardous Activity	108
Deaf Donor	Disabled Donor	88
Decompression Illness	Decompression Illness	83
Decompression Sickness	Decompression Illness	83
Deep Sea Diving	Hazardous Activity	108
Deep Vein Thrombosis	Thrombosis and Thrombophilia	186
Delayed Faint	Faints	96
Dementia	Central Nervous System Disease	73
Dengue (Virus) Fever - Contact With	Non-Contagious Diseases - Contact With	150
Dengue (Virus) Fever - Infection	Tropical Viruses	194
Dental Abscess	Infection - Acute	127
Dental Treatment	Dental Treatment	84
Depression	Mental Health Problems	143
Derma-Rolling	Body Piercing	71
Dermatitis	Dermatitis	85
Dermoid Cyst	Accept	52
Desensitisation Injections for Hay Fever	Allergy	56
Detox - Complementary Therapy	Complementary Therapy	80
Detox - for Alcohol or Drug Abuse	Addiction and Drug Abuse	54
Diabetes - Sugar	Diabetes Mellitus	86
Diabetes Insipidus	Diabetes Insipidus	86
Diabetes Insipidus - Cranial	Pituitary Disorders	157
Diabetes Mellitus	Diabetes Mellitus	86
Diabetic Ulcer	Diabetes Mellitus	86
Dianette	Acne	53
Diarrhoea	Gastroenteritis	101
Diarrhoea and Vomiting – Contact With	Infectious Diseases - Contact With	130
Diet Pills	Donor Weight	90
Digoxin - for Heart Failure	Cardiovascular Disease	72
Dilatation and Curettage	Surgery	182

For	See	Page
Diphtheria - Contact With	Infectious Diseases - Contact With	130
Diphtheria - Immunisation	Immunisation - Non-Live	125
Diphtheria - Infection	Infection - Acute	127
Diphtheria, Tetanus and Poliomyelitis - Immunisation	Immunisation - Non-Live	125
Diphtheria, Tetanus, Pertussis and Poliomyelitis - Immunisation	Immunisation - Non-Live	125
Diphtheria, Tetanus, Pertussis, Poliomyelitis and Haemophilus Type b - Immunisation	Immunisation - Non-Live	125
Disabled Donor	Disabled Donor	88
Disc Surgery	Back Problems	64
Discoid Lupus Erythematosus	Skin Disease	174
Disulfiram	Addiction and Drug Abuse	54
Diuretics	Diuretics	89
Diverticular Disease	Diverticular Disease	90
Diverticulitis	Diverticular Disease	90
Diverticulosis	Diverticular Disease	90
Diving	Hazardous Activity	108
Dizzy Spells - Feeling Faint	Faints	96
Dizzy Spells - Vertigo	Vertigo	199
Dog Bite	Animal Bite (Non-Human)	59
Donated Human Egg Recipient	Fertility	97
Donated Human Embryo Recipient	Fertility	97
Donated Human Sperm Recipient	Fertility	97
Donation Frequency	Frequency of Donation	99
Donor Weight	Donor Weight	90
Dopamine Receptor Agonist Drugs	Central Nervous System Disease	73
Double Red Cell Donation Frequency	Frequency of Donation	99
Drogenil®	Anti-Androgens	60
Dropped Beats	Arrhythmias	62
Drug Abuse	Addiction and Drug Abuse	54
Drug Allergy	Allergy	56
Drug Index	Drug Index	91
Drug Treatment	Drug Treatment	91
Drunk	Addiction and Drug Abuse	54
Dry Needling	Complementary Therapy	80
DTP Immunisation	Immunisation - Non-Live	125
Ductal Carcinoma in Situ	Malignancy	141
Duodenal Ulcer	Peptic Ulcer	156
Dura Mater Graft	Prion Associated Diseases	164
Dutasteride - for Hair Loss	Alopecia	57

For	See	Page
Dutasteride - for Prostate Problems	Anti-Androgens	60
DVT	Thrombosis and Thrombophilia	186
Dysentery	Infection - Acute	127
Dysentery - Amoebic - Contact With	Infectious Diseases - Contact With	130
Dysentery - Amoebic - Infection	Infection - Acute	127
Dyspraxia	Neurobehavioral Disorders	148
E		
Ear Infection	Infection - General	129
Ear Piercing	Body Piercing	71
Ebola Fever - Contact With	Viral Haemorrhagic Fever	200
Ebola Fever - Infection	Viral Haemorrhagic Fever	200
EBV - Contact With	Infectious Diseases - Contact With	130
EBV - Infection	Infection - Acute	127
Ecstasy	Addiction and Drug Abuse	54
Ectopic Pregnancy	Pregnancy	163
Ectopics - Heart	Arrhythmias	62
Eczema	Dermatitis	85
EDS	Ehlers Danlos Syndrome	92
Ehlers Danlos Syndrome	Ehlers Danlos Syndrome	92
Electrolysis	Hair Removal	108
Elidel® - for Dermatitis	Dermatitis	85
Elidel® - for Skin Disease	Skin Disease	174
Elliptocytosis	Haemolytic Anaemia	107
Emergency Response Vehicle Drivers	Hazardous Activity	108
Emphysema	Respiratory Disease	170
Empty Sella Syndrome	Pituitary Disorders	157
Encephalitis	Infection - General	129
Encephalitis Japanese - Immunisation	Immunisation - Non-Live	125
Endocarditis	Endocarditis	93
Endometrial Ablation	Surgery	182
Endometriosis	Endometriosis	93
Endoscopy	Endoscopy	93
Enterocolitis - Crohn's	Inflammatory Bowel Disease	131
Enterocolitis - Infective	Gastroenteritis	101
Epidermoid Cyst	Accept	52
Epilation	Hair Removal	108
Epilepsy	Epilepsy	94
Episcleritis	Eye Disease	95
Epistaxis	Nose Bleeds	152

For	See	Page
Erectile Dysfunction	Erectile Dysfunction	94
Erosions - Duodenal	Peptic Ulcer	156
Erosions - Gastric	Peptic Ulcer	156
Erythema Infectiosum	Parvovirus B19	155
Erythropoietic Protoporphyrria	Porphyria	162
Essential Thrombocythaemia	Haematological Disease	103
Etretinate	Etretinate	95
Exenatide	Diabetes Mellitus	86
Exercise	Exercise	95
Exposure to Hazardous Materials	Hazardous Activity	108
Eye Disease	Eye Disease	95
Eye Drops	Eye Disease	95
Eye Surgery	Eye Disease	95
F		
Face Lift	Surgery	182
Factor I (one) deficiency	Bleeding Disorder	65
Factor II (two) deficiency	Bleeding Disorder	65
Factor IX Deficiency	Bleeding Disorder	65
Factor V (five) deficiency	Bleeding Disorder	65
Factor V Leiden	Thrombosis and Thrombophilia	186
Factor VII Deficiency	Bleeding Disorder	65
Factor VIII Deficiency	Bleeding Disorder	65
Factor X Deficiency	Bleeding Disorder	65
Factor XI (eleven) deficiency	Bleeding Disorder	65
Factor XIII (thirteen) deficiency	Bleeding Disorder	65
Faecal Microbiota Transplantation	Complementary Therapy	80
Faints	Faints	96
Familial CJD	Prion Associated Diseases	164
Familial Pseudohyperkalaemia	Familial Pseudohyperkalaemia	97
Fatal Familial Insomnia	Prion Associated Diseases	164
Fatty Liver	Liver Disease	137
Fertility	Fertility	97
Fertility Treatment	Fertility	97
Fever	Pyrexia	167
Fibrinogen deficiency	Bleeding Disorder	65
Fibroid Operation	Surgery	182
Fibromyalgia	Fibromyalgia	99
Fibromyositis	Fibromyalgia	99
Fibrositis	Fibromyalgia	99

For	See	Page
Fifth Disease	Parvovirus B19	155
Filariasis	Infection - Chronic	128
Finasteride	Anti-Androgens	60
Fire Service Emergency Response Personnel	Hazardous Activity	108
Fish Pedicure	Accept	52
Fits	Epilepsy	94
Flagellation	Body Piercing	71
Flash glucose monitor	Shunts, Stents and Devices	173
Flu - Contact With	Infectious Diseases - Contact With	130
Flu - Immunisation	Immunisation - Non-Live	125
Flu - Infection	Infection - Acute	127
Flu Jab	Immunisation - Non-Live	125
Flutamide	Anti-Androgens	60
Flying - Air Crew	Air Crew and Air Traffic Controllers	55
Flying - Passenger	Travel	193
Folate Deficiency	Anaemia	58
Follicle Stimulating Hormone	Fertility	97
Food Allergy	Allergy	56
Food Poisoning	Gastroenteritis	101
Foreign Travel	Travel	193
Fracture	Accident	52
Frequency of Donation	Frequency of Donation	99
Frozen Shoulder	Tendonitis	185
Fungal Infection	Infection - General	129
G		
G6PD Deficiency	Haemolytic Anaemia	107
Gall Bladder Disease	Gall Bladder Disease	100
Gall Bladder Infection	Gall Bladder Disease	100
Gall Bladder Operation	Gall Bladder Disease	100
Gall Stones	Gall Bladder Disease	100
Ganglion Removal	Surgery	182
Gangrene	Cardiovascular Disease	72
Gardasil® - Immunisation	Immunisation - Non-Live	125
Gastrectomy	Surgery	182
Gastric Banding	Surgery	182
Gastric Flu	Gastroenteritis	101
Gastritis	Acid Indigestion	52
Gastroenteritis	Gastroenteritis	101
Gastrointestinal Disease	Gastrointestinal Disease	101

For	See	Page
Gastro-Oesophageal Reflux Disorder (GORD)	Acid Indigestion	52
Gastroscopy	Endoscopy	93
Gastrostomy	Stoma	181
Gender Reassignment	Transgender and Non-Binary Individuals	192
Gender-affirming Hormone Therapy	Hormone Replacement and Sex Hormone Therapy	120
General Anaesthetic	Anaesthetic	59
Genital Herpes Infection	Herpes Simplex	118
Genital Warts	Warts	201
Genitourinary Medicine	Sexually Transmitted Disease	172
German Measles - Contact With	Infectious Diseases - Contact With	130
German Measles - Immunisation	Immunisation - Live	125
German Measles - Infection	Infection - Acute	127
Gerstmann-Sträussler-Scheinker Disease	Prion Associated Diseases	164
Giardiasis	Giardiasis	101
Gilbert's Disease	Liver Disease	137
Gilbert's Syndrome	Liver Disease	137
Gingivitis	Dental Treatment	84
Glandular Fever - Contact With	Infectious Diseases - Contact With	130
Glandular Fever - Infection	Infection - Acute	127
Glaucoma	Eye Disease	95
Glomerulonephritis	Kidney and Bladder Disease	133
Glucosamine	Vitamins and Other Nutritional Supplements	200
Glucose monitor	Shunts, Stents and Devices	173
Glycogen Storage Disease	Glycogen Storage Disease	102
Goitre	Thyroid Disease	189
Golfer's Elbow	Tendonitis	185
Gonadotrophin Treatment - Infertility	Fertility	97
Gonorrhoea	Sexually Transmitted Disease	172
GORD	Acid Indigestion	52
Gout	Gout	102
Grand Mal	Epilepsy	94
Granuloma Inguinale	Infection - Chronic	128
Grave's Disease	Thyroid Disease	189
Growth - Malignant	Malignancy	141
Growth Hormone	Growth Hormone	102
Guillain-Barre Syndrome	Guillain-Barre Syndrome	103
GUM Clinic	Sexually Transmitted Disease	172
Gum disease	Dental Treatment	84

For	See	Page
H		
H2 Receptor Antagonists	Acid Indigestion	52
Haematological Disease	Haematological Disease	103
Haematuria	Haematuria	104
Haemochromatosis	Haemochromatosis	104
Haemoglobin Disorders	Haemoglobin Disorders	105
Haemoglobin Estimation	Haemoglobin Estimation	105
Haemoglobin, high	Polycythaemia and Raised Haemoglobin	161
Haemolytic Anaemia	Haemolytic Anaemia	107
Haemophilia	Bleeding Disorder	65
Haemophilia - Household Member / Carer	Bleeding Disorder	65
Haemophilia - Sexual Partner	Bleeding Disorder	65
Haemophilia Carrier	Bleeding Disorder	65
Haemophilus Influenzae - Contact With	Infectious Diseases - Contact With	130
Haemophilus Influenzae - Immunisation	Immunisation - Non-Live	125
Haemophilus Influenzae - Infection	Infection - Acute	127
Haemorrhoids	Haemorrhoids	107
Hair Loss Treatments	Alopecia	57
Hair Removal	Hair Removal	108
Hair Transplant	Alopecia	57
Hand, Foot and Mouth Disease	Infection - Acute	127
Hand, Foot and Mouth Disease - Contact With	Infectious Diseases - Contact With	130
Hang Gliding	Hazardous Activity	108
Hashimoto's Disease	Thyroid Disease	189
Hay Fever	Allergy	56
Hay Fever Desensitisation	Allergy	56
Hazardous Activity	Hazardous Activity	108
Hazardous Hobbies	Hazardous Activity	108
Hazardous Occupation	Hazardous Activity	108
Head Injury	Accident	52
Head Lice	Infection - Acute	127
Head Lice - Contact With	Non-Contagious Diseases - Contact With	150
Headache	Headache	109
Heaf Test	Tuberculin PPD Test	196
Health Care Worker	Health Care Worker	109
Heart Attack	Cardiovascular Disease	72
Heart Block	Arrhythmias	62
Heart Condition	Cardiovascular Disease	72
Heart Failure	Cardiovascular Disease	72

For	See	Page
Heart Murmur	Cardiovascular Disease	72
Heart Operation	Cardiac Surgery	71
Heart Pills	Cardiovascular Disease	72
Heart Valve Abnormality	Cardiovascular Disease	72
Heartburn - Including Treatment	Acid Indigestion	52
Heavy Goods Vehicle Driver	Hazardous Activity	108
Heavy Machine Operator	Hazardous Activity	108
Heavy Periods	Periods	156
Help Lines	National Help Lines	147
Henna Painting	Henna Painting	110
Henoch Schonlein Purpura	Autoimmune Disease	63
Heparin	Anticoagulant Therapy	61
Hepatic Porphyria	Porphyria	162
Hepatitis	Hepatitis	110
Hepatitis A	Hepatitis A	111
Hepatitis A - Immunisation	Hepatitis A	111
Hepatitis A - Infection	Hepatitis A	111
Hepatitis A - Person Sharing Home	Hepatitis A	111
Hepatitis A - Sexual Partner of Affected Individual	Hepatitis A	111
Hepatitis B	Hepatitis B	112
Hepatitis B - Immunisation	Hepatitis B	112
Hepatitis B - Infection/Carrier	Hepatitis B	112
Hepatitis B - Person Sharing Home	Hepatitis B	112
Hepatitis B - Sexual Partner of Affected Individual	Hepatitis B	112
Hepatitis C	Hepatitis C	115
Hepatitis C - Person Sharing Home	Hepatitis C	115
Hepatitis C - Person with current Hepatitis C infection	Hepatitis C	115
Hepatitis C - Sexual Partner of Affected Individual	Hepatitis C	115
Hepatitis C – Person with treated Hepatitis C infection	Hepatitis C	115
Hepatitis E	Hepatitis E	117
Hepatitis of Unknown Cause	Hepatitis of Unknown Cause	117
Hepatitis of Unknown Cause - Affected Individuals	Hepatitis of Unknown Cause	117
Hepatitis of Unknown Cause - Person Sharing Home	Hepatitis of Unknown Cause	117
Hepatitis of Unknown Cause - Sexual Partner of Affected Individual	Hepatitis of Unknown Cause	117
Herbal Remedy	Complementary Therapy	80
Hereditary Coproporphyrria	Porphyria	162
Hereditary Elliptocytosis	Haemolytic Anaemia	107
Hereditary Spherocytosis	Haemolytic Anaemia	107

For	See	Page
Hernia Repair	Surgery	182
Herpes - Genital/Oral	Herpes Simplex	118
Herpes Simplex	Herpes Simplex	118
Herpes Zoster - Contact With	Infectious Diseases - Contact With	130
Herpes Zoster - Immunisation	Immunisation - Live	125
Herpes Zoster - Infection	Infection - Acute	127
Heterografts	Xenotransplantation	203
Heterosexual Rape	Non-Consented Exposure to Human Body Fluids	149
Heterosexual sexual assault	Non-Consented Exposure to Human Body Fluids	149
HGV Driver	Hazardous Activity	108
Hiatus Hernia	Acid Indigestion	52
Hiatus Hernia Repair	Surgery	182
HIB - Immunisation	Immunisation - Non-Live	125
High Affinity Haemoglobin	Haemoglobin Disorders	105
High Blood Pressure	Blood Pressure - High	67
High Cholesterol	Hypercholesterolaemia	122
High haemoglobin	Polycythaemia and Raised Haemoglobin	161
High platelet count	Platelet Count	158
Hina	Henna Painting	110
Hip Replacement	Surgery	182
HIV	HIV	119
HIV - Infection	HIV	119
HIV - Person Sharing Home	HIV	119
HIV - Sexual Partner of Affected Individual	HIV	119
Hives	Allergy	56
Hodgkin's Lymphoma	Malignancy	141
Homeopathy	Complementary Therapy	80
Hormone Replacement and Sex Hormone Therapy	Hormone Replacement and Sex Hormone Therapy	120
Hospital Staff	Health Care Worker	109
HPV - Immunisation	Immunisation - Non-Live	125
HRT	Hormone Replacement and Sex Hormone Therapy	120
HSV-1	Herpes Simplex	118
HSV-2	Herpes Simplex	118
HTLV	HTLV	120
HTLV - Infection	HTLV	120
HTLV - Person Sharing Home	HTLV	120
HTLV - Sexual Partner of Affected Individual	HTLV	120
Hughes' Syndrome - as part of an Autoimmune Disease	Autoimmune Disease	63

For	See	Page
Hughes' Syndrome - Causing Infertility	Fertility	97
Hughes' Syndrome - Causing Thrombosis	Thrombosis and Thrombophilia	186
Human Bite	Non-Consented Exposure to Human Body Fluids	149
Human Papilloma Virus - Immunisation	Immunisation - Non-Live	125
Human Parvovirus	Parvovirus B19	155
Human Tissue Recipient	Tissue and Organ Recipients	189
Huntington's Chorea	Huntington's Disease	122
Huntington's Disease	Huntington's Disease	122
Hydatid Disease	Infection - Chronic	128
Hydatidiform Mole	Pregnancy	163
Hydrocephalus	Hydrocephalus	122
Hydroxychloroquine (Alopecia)	Alopecia	57
Hydroxychloroquine (Arthritis)	Autoimmune Disease	63
Hydroxychloroquine for Rheumatoid Arthritis	Autoimmune Disease	63
Hyperaldosteronism	Conn's Syndrome	81
Hypercholesterolaemia	Hypercholesterolaemia	122
Hyperkalaemia	Familial Pseudohyperkalaemia	97
Hypertension	Blood Pressure - High	67
Hyperthyroidism	Thyroid Disease	189
Hypnotics	Hypnotics	123
Hypofibrinogenaemia	Bleeding Disorder	65
Hypotension	Blood Pressure - Low	68
Hypothalamic Tumour	Pituitary Disorders	157
Hypothyroidism	Thyroid Disease	189
Hysterectomy	Surgery	182
I		
IBS	Irritable Bowel Syndrome	132
Idiopathic Intracranial Hypertension	Central Nervous System Disease	73
Idiopathic Thrombocytopenic Purpura	Immune Thrombocytopenia	123
IgA deficiency	Immunodeficiency	126
IIH	Central Nervous System Disease	73
Ileostomy	Stoma	181
Immune Thrombocytopenia	Immune Thrombocytopenia	123
Immunisation	Immunisation	124
Immunisation - Live	Immunisation - Live	125
Immunisation - Non-Live	Immunisation - Non-Live	125
Immunodeficiency	Immunodeficiency	126
Immunoglobulin A deficiency	Immunodeficiency	126
Immunoglobulin Therapy	Immunoglobulin Therapy	126

For	See	Page
Immunosuppression	Immunodeficiency	126
Impetigo - Contact With	Infectious Diseases - Contact With	130
Impetigo - Infection	Infection - Acute	127
Implanted Devices	Shunts, Stents and Devices	173
Implanted electrodes	Shunts, Stents and Devices	173
Impotence	Erectile Dysfunction	94
Imvanex	Mpox (Monkeypox)	145
Indigestion - Including Treatment	Acid Indigestion	52
Infected Eye	Infection - Acute	127
Infected Wound	Wounds, Mouth and Skin Ulcers	203
Infection - Acute	Infection - Acute	127
Infection - Chronic	Infection - Chronic	128
Infection - General	Infection - General	129
Infection - Tropical	Infection - General	129
Infectious Diseases - Contact With	Infectious Diseases - Contact With	130
Infectious Mononucleosis	Infection - Acute	127
Infectious Mononucleosis - Contact With	Infectious Diseases - Contact With	130
Infective Colitis	Infection - General	129
Infertility	Fertility	97
Inflammation - due to Infection	Infection - General	129
Inflammation - using Nonsteroidal Anti-Inflammatory Drugs	Nonsteroidal Anti-Inflammatory Drugs	151
Inflammation - using Steroid Treatment	Steroid Therapy	180
Inflammatory Bowel Disease	Inflammatory Bowel Disease	131
Inflammatory Eye Disease	Eye Disease	95
Influenza - Contact With	Infectious Diseases - Contact With	130
Influenza - Immunisation	Immunisation - Non-Live	125
Influenza - Infection	Infection - Acute	127
Inguinal Hernia Repair	Surgery	182
Inguinale Granuloma	Infection - Chronic	128
Inhaler - for Asthma	Asthma	63
Inhaler - for Bronchitis	Respiratory Disease	170
Inherited Diseases	Inherited Diseases	131
Injectable Tanning Agents	Blood Safety Entry	68
Injected Drugs	Addiction and Drug Abuse	54
Injection in Joint - for Infection	Infection - General	129
Injection in Joint - Steroid	Steroid Therapy	180
Injection of Prostaglandins for Impotence	Erectile Dysfunction	94
Injections - Iron	Anaemia	58

For	See	Page
Injuries	Accident	52
INOCA	Cardiovascular Disease	72
Inoculation Injury	Non-Consented Exposure to Human Body Fluids	149
Inoculations	Immunisation	124
Intermittent Claudication	Cardiovascular Disease	72
Intermittently scanned glucose monitor	Shunts, Stents and Devices	173
Intersex	Transgender and Non-Binary Individuals	192
Interstitial Cystitis	Kidney and Bladder Disease	133
Interval Between Donations	Frequency of Donation	99
Intoxication	Addiction and Drug Abuse	54
Intra-Articular Injection - for Infection	Infection - General	129
Intra-Articular Injection - Steroid	Steroid Therapy	180
Intraepithelial squamous cell carcinoma	Malignancy	141
Intravenous Drug Use	Addiction and Drug Abuse	54
Investigations	Investigations	132
Investigations Pending	Investigations	132
IPV - Immunisation	Immunisation - Non-Live	125
Iritis	Eye Disease	95
Iron Deficiency	Anaemia	58
Iron Treatment	Anaemia	58
Irregular Heartbeat	Arrhythmias	62
Irritable Bowel Syndrome	Irritable Bowel Syndrome	132
Ischaemic Heart Disease	Cardiovascular Disease	72
Isotretinoin	Acne	53
ITP	Immune Thrombocytopenia	123
IV Drug Abuse	Addiction and Drug Abuse	54
IVDM	Addiction and Drug Abuse	54
IVDU	Addiction and Drug Abuse	54
IVF	Fertility	97
J		
Jabs	Immunisation	124
Japanese Encephalitis - Immunisation	Immunisation - Non-Live	125
Jaundice	Jaundice	132
Joint Injection - for Infection	Infection - General	129
Joint Injection - for Osteoarthritis	Osteoarthritis	152
Joint Injection - Steroid	Steroid Therapy	180
K		
Kala-Azar	Infection - Chronic	128
Kidney and Bladder Disease	Kidney and Bladder Disease	133
Kidney Donor	Surgery	182

For	See	Page
Kidney Failure	Kidney and Bladder Disease	133
Kidney Infection	Infection - General	129
Kidney Recipient	Tissue and Organ Recipients	189
Kidney Stones	Kidney and Bladder Disease	133
Klinefelter's Syndrome	Klinefelter's Syndrome	136
Knocked Out	Accident	52
L		
Labyrinthitis	Infection - Acute	127
Ladders - Working at Heights	Hazardous Activity	108
Laminectomy	Back Problems	64
Language Processing Disorder	Neurobehavioral Disorders	148
Laparoscopy	Endoscopy	93
Large Goods Vehicle Driver	Hazardous Activity	108
Laryngitis	Infection - General	129
Laser Refractive Surgery	Laser Treatment	136
Laser Skin Treatment	Laser Treatment	136
Laser Treatment	Laser Treatment	136
Lassa Fever - Contact With	Viral Haemorrhagic Fever	200
Lassa Fever - Infection	Viral Haemorrhagic Fever	200
Latent Tuberculosis	Tuberculosis	197
Latex Allergy	Allergy	56
LBBB	Cardiovascular Disease	72
Left Bundle Branch Block	Cardiovascular Disease	72
Legionnaires' Disease	Infection - Acute	127
Leishmaniasis	Infection - Chronic	128
Lentigo Maligna	Malignancy	141
Lentigo Maligna Melanoma	Malignancy	141
Leptospirosis	Infection - Acute	127
Leukaemia	Malignancy	141
Leukocyte Donation Frequency	Frequency of Donation	99
LGV (infection)	Sexually Transmitted Disease	172
LGV Driver	Hazardous Activity	108
Lichen Planus	Skin Disease	174
Lichen Sclerosus	Skin Disease	174
Ligament Reconstruction - with Graft	Tissue and Organ Recipients	189
Ligament Reconstruction - without Graft	Surgery	182
Lipoma Removed	Surgery	182
Liraglutide – for Diabetes Mellitus	Diabetes Mellitus	86
Liraglutide – for weight loss	Donor Weight	90

For	See	Page
Listeriosis	Infection - Acute	127
Lithium Treatment	Mental Health Problems	143
Liver Disease	Liver Disease	137
Liver Failure	Liver Disease	137
Local Anaesthetic	Anaesthetic	59
Local Injection to Numb	Anaesthetic	59
Loop Diathermy - Cervix	Cervical Dysplasia	75
Loop Diathermy - Other	Surgery	182
Low platelet count	Platelet Count	158
Lucentis®	Eye Disease	95
Lumbar Puncture	Investigations	132
Lumbar Peritoneal Shunt	Shunts, Stents and Devices	173
Lumpectomy	Surgery	182
Lung Cancer	Malignancy	141
Lupus Erythematosus	Autoimmune Disease	63
Luteinising Hormone	Fertility	97
Lyme Disease	Infection - Acute	127
Lymphocyte Donation Frequency	Frequency of Donation	99
Lymphoedema	Wounds, Mouth and Skin Ulcers	203
Lymphogranuloma Venereum	Sexually Transmitted Disease	172
Lymphoma	Malignancy	141
M		
Machine Operator - Heavy Machinery	Hazardous Activity	108
Macugen®	Eye Disease	95
Magic Eye	Endoscopy	93
Malaria	Malaria	140
Malaria - Contact With	Non-Contagious Diseases - Contact With	150
Malaria - plasmapheresis donors	Malaria - plasmapheresis donors	140
Malaria Risk	Malaria	140
Male Homosexual Rape	Non-Consented Exposure to Human Body Fluids	149
Male Homosexual sexual assault	Non-Consented Exposure to Human Body Fluids	149
Malignancy	Malignancy	141
Malignancy	Cervical Dysplasia	
Malignant Hypertension	Blood Pressure - High	67
Malignant Melanoma	Malignancy	141
Manic Depression	Mental Health Problems	143
Mantoux Test	Tuberculin PPD Test	196
Marburg Fever - Contact With	Viral Haemorrhagic Fever	200
Marburg Fever - Infection	Viral Haemorrhagic Fever	200

For	See	Page
Marfan's Syndrome	Marfan's Syndrome	142
Mastectomy - for Malignancy	Malignancy	141
Mastectomy - for Other Reasons (Including Increased risk of malignancy)	Surgery	182
Mastoid Operation	Surgery	182
Mastoiditis	Infection - General	129
ME	Chronic Fatigue Syndrome	77
Measles - Contact With	Infectious Diseases - Contact With	130
Measles - Immunisation	Immunisation - Live	125
Measles - Infection	Infection - Acute	127
Measles and Rubella - Immunisation	Immunisation - Live	125
Measles, Mumps and Rubella - Immunisation	Immunisation - Live	125
Medication	Drug Treatment	91
Mehndi	Henna Painting	110
Melanoma - Malignant	Malignancy	141
Memorial Tattoos	Body Piercing	71
Meniere's Disease	Meniere's Disease	142
Meningitis - Contact With	Infectious Diseases - Contact With	130
Meningitis - Immunisation	Immunisation - Non-Live	125
Meningitis - Infection	Infection - Acute	127
Meningococcal Meningitis - Immunisation	Immunisation - Non-Live	125
Menopause	Menopause	143
Menstruation	Periods	156
Mental Health Problems	Mental Health Problems	143
MERS	Coronavirus Infection (COVID-19)	81
Methicillin Resistant Staphylococcus Aureus - Carrier	Infection - General	129
Methicillin Resistant Staphylococcus Aureus - Contact With	Non-Contagious Diseases - Contact With	150
Methicillin Resistant Staphylococcus Aureus - Infection	Infection - General	129
Methotrexate (Arthritis)	Autoimmune Disease	63
Methotrexate (Ectopic Pregnancy)	Pregnancy	163
Metrodin HP	Fertility	97
MGEN (infection)	Sexually Transmitted Disease	172
MGUS	Haematological Disease	103
MI	Cardiovascular Disease	72
Middle Eastern Respiratory Syndrome	Coronavirus Infection (COVID-19)	81
Migraine	Migraine	143
Miner Working Underground	Hazardous Activity	108
Mini Stroke	Cerebrovascular Disease and Intracranial Haemorrhage	74
Miscarriage	Pregnancy	163

For	See	Page
Missed Period	Periods	156
Mitral Valve Prolapse	Cardiovascular Disease	72
MMR	Immunisation - Live	125
Modafinil	Narcolepsy	146
Molar Pregnancy	Pregnancy	163
Mole Pregnancy	Pregnancy	163
Mole Removal - Other	Surgery	182
Mole Removal - Related to Pregnancy	Pregnancy	163
Molluscum Contagiosum	Warts	201
Moniliasis	Thrush	188
Monkeypox	Mpox (Monkeypox)	145
Monoclonal Antibody Therapy	Monoclonal antibody therapy and other Biological Modalities	144
Monoclonal antibody therapy and other Biological Modalities	Monoclonal antibody therapy and other Biological Modalities	144
Monoclonal Gammopathy of Unknown Significance	Haematological Disease	103
Mood Swings	Mental Health Problems	143
Morning After Pill	Contraceptive Use	81
Motor Neurone Disease	Central Nervous System Disease	73
Motor Sport	Hazardous Activity	108
Mountaineering	Hazardous Activity	108
Mouth Ulcers	Wounds, Mouth and Skin Ulcers	203
Moxibustion	Complementary Therapy	80
Mpox	Mpox (Monkeypox)	145
Mpox (Monkeypox)	Mpox (Monkeypox)	145
MRI Scan	Investigations	132
MRSA - Carrier	Infection - General	129
MRSA - Contact With	Non-Contagious Diseases - Contact With	150
MRSA - Infection	Infection - General	129
MS	Central Nervous System Disease	73
Multiple Endocrine Neoplasia Syndrome	Malignancy	141
Multiple Sclerosis	Central Nervous System Disease	73
Mumps - Contact With	Infectious Diseases - Contact With	130
Mumps - Immunisation	Immunisation - Live	125
Mumps - Infection	Infection - Acute	127
Murmur	Cardiovascular Disease	72
Muscular Dystrophy	Disabled Donor	88
MUSE®	Erectile Dysfunction	94
Myalgic Encephalomyelitis	Chronic Fatigue Syndrome	77
Myasthenia Gravis - Acquired	Autoimmune Disease	63

For	See	Page
Myasthenia Gravis - Congenital	Disabled Donor	88
Mycoplasma genitalium	Sexually Transmitted Disease	172
Myelodysplastic Syndrome	Haematological Disease	103
Myeloproliferative Syndrome	Haematological Disease	103
Myocardial Infarct	Cardiovascular Disease	72
Myocardial ischaemia with non-obstructive coronary arteries	Cardiovascular Disease	72
Myocarditis	Cardiovascular Disease	72
Myomectomy	Surgery	182
Myxoedema	Thyroid Disease	189
N		
NAFLD	Liver Disease	137
Nail Fungal Infection	Infection - Chronic	128
Nail Infection	Infection - General	129
Narcolepsy	Narcolepsy	146
Nasal Polyps	Allergy	56
NASH	Liver Disease	137
National Help Lines	National Help Lines	147
Navel Piercing	Body Piercing	71
Needle-Stick Injury	Non-Consented Exposure to Human Body Fluids	149
Neonatal Jaundice	Jaundice	132
Neotigason® - for Acne	Acne	53
Neotigason® - for Psoriasis	Psoriasis	166
Neotigason® - for Skin Disease	Skin Disease	174
Nephrectomy	Surgery	182
Nephritis	Kidney and Bladder Disease	133
Nephrostomy	Stoma	181
Nerve stimulator devices	Shunts, Stents and Devices	173
Nervous Breakdown	Mental Health Problems	143
Neurobehavioral Disorders	Neurobehavioral Disorders	148
Neurofibromatosis	Neurofibromatosis	148
Neurological Problems	Central Nervous System Disease	73
Neuromodulator devices	Shunts, Stents and Devices	173
Neurostimulation	Neurosurgery	148
Neurosurgery	Neurosurgery	148
Nicorette®	Smoking Cessation	177
Nicotine Replacement Therapy	Smoking Cessation	177
Nicotinell®	Smoking Cessation	177
Night Sweats	Night Sweats	149
Nipple Piercing	Body Piercing	71

For	See	Page
NiQuitin®	Smoking Cessation	177
Non-Alcoholic Fatty Liver Disease	Liver Disease	137
Non-alcoholic steatohepatitis	Liver Disease	137
Non-Alcoholic Steatohepatitis	Liver Disease	137
Non-Binary Individuals	Transgender and Non-Binary Individuals	192
Non-Consented Exposure to Human Body Fluids	Non-Consented Exposure to Human Body Fluids	149
Non-Contagious Diseases - Contact With	Non-Contagious Diseases - Contact With	150
Non-Hodgkin's Lymphoma	Malignancy	141
Non-Human Organ Perfusion	Xenotransplantation	203
Non-Specific Urethritis - Acute	Infection - Acute	127
Non-Specific Urethritis - Chronic	Infection - Chronic	128
Nonsteroidal Anti-Inflammatory Drugs	Nonsteroidal Anti-Inflammatory Drugs	151
Norovirus – Contact With	Infectious Diseases - Contact With	130
Nose Bleeds	Nose Bleeds	152
Nose Piercing	Body Piercing	71
Novel Coronavirus	Coronavirus Infection (COVID-19)	81
NSAID	Nonsteroidal Anti-Inflammatory Drugs	151
NSU - Acute	Infection - Acute	127
NSU - Chronic	Infection - Chronic	128
Null-Cell Adenoma	Pituitary Disorders	157
O		
Obesity	Donor Weight	90
Obsessive Compulsive Disorder	Neurobehavioral Disorders	148
OCD	Neurobehavioral Disorders	148
Ocular Surgery	Eye Disease	95
Ocular Tissue Recipient	Tissue and Organ Recipients	189
Operation	Surgery	182
Optic Neuritis	Central Nervous System Disease	73
Oral Herpes	Herpes Simplex	118
Oral Polio - Immunisation	Immunisation - Live	125
Oral Typhoid - Immunisation	Immunisation - Live	125
Orf	Infection - Acute	127
Organ Donor	Surgery	182
Organ Recipient	Tissue and Organ Recipients	189
Oseltamivir	Oseltamivir	152
Osteitis Deformans	Paget's Disease of Bone	154
Osteoarthritis	Osteoarthritis	152
Osteomalacia	Osteopenia	153
Osteomyelitis	Osteomyelitis	153

For	See	Page
Osteopenia	Osteopenia	153
Osteoporosis	Osteopenia	153
Ovarian Cancer	Malignancy	141
Ovarian Cyst - Malignant	Malignancy	141
Ovarian Cyst Removed - Non-Malignant	Surgery	182
Ovary Removed - Malignant	Malignancy	141
Ovary Removed - Non-Malignant	Surgery	182
Over Weight	Donor Weight	90
Overactive Thyroid	Thyroid Disease	189
P		
Pacemaker	Shunts, Stents and Devices	173
Paget's Disease of Bone	Paget's Disease of Bone	154
Paget's Disease of the Breast	Malignancy	141
Pain Killers	Pain Killers	154
Painful Bladder Syndrome	Kidney and Bladder Disease	133
Palpitations	Palpitations	154
Parachuting	Hazardous Activity	108
Paranoia	Mental Health Problems	143
Paratyphoid	Infection - Chronic	128
Parkinson's Disease	Central Nervous System Disease	73
Partially Sighted	Disabled Donor	88
Parvovirus	Parvovirus B19	155
Parvovirus B19	Parvovirus B19	155
Patent Foramen Ovale	Cardiovascular Disease	72
PBC	Liver Disease	137
PCOS	Accept	52
PE	Thrombosis and Thrombophilia	186
Pegaptanib sodium	Eye Disease	95
Pelvic Floor Repair	Surgery	182
Pemphigoid	Skin Disease	174
Pemphigus	Skin Disease	174
PEP	Pre- and Post-Exposure Prophylaxis for HIV prevention	162
Peptic Ulcer	Peptic Ulcer	156
Peri-Articular Injection - for Infection	Infection - General	129
Peri-Articular Injection - Steroid	Steroid Therapy	180
Pericarditis - Viral	Infection - Acute	127
Period Problems	Periods	156
Periodontal disease	Dental Treatment	84
Periodontitis	Dental Treatment	84

For	See	Page
Periods	Periods	156
Peripheral Vascular Disease	Cardiovascular Disease	72
Peritonitis - not Requiring Surgery	Infection - General	129
Peritonitis - Requiring Surgery	Surgery	182
Peritonsillar Abscess	Infection - Acute	127
Permanent Makeup	Body Piercing	71
Pernicious Anaemia	Anaemia	58
Peroneal Muscular Atrophy	Disabled Donor	88
Perthes' Disease	Perthes' Disease	157
Pertussis - Contact With	Infectious Diseases - Contact With	130
Pertussis - Immunisation	Immunisation - Non-Live	125
Pertussis - Infection	Infection - Acute	127
Petit Mal	Epilepsy	94
Pharyngitis	Infection - General	129
Phlebitis	Superficial Thrombophlebitis	181
Phlebothrombosis	Superficial Thrombophlebitis	181
Phlebotomus Fever	Infection - Acute	127
Piercing	Body Piercing	71
Piles	Haemorrhoids	107
Pilot - Aircraft	Air Crew and Air Traffic Controllers	55
Pimecrolimus - for Dermatitis	Dermatitis	85
Pimecrolimus - for Skin disease	Skin Disease	174
Pituitary Disorders	Pituitary Disorders	157
Pituitary Extract - Human	Prion Associated Diseases	164
Platelet Count	Platelet Count	158
Platelet Disorders	Platelet Disorders	159
Platelet Donation Frequency	Frequency of Donation	99
Platelet Rich Plasma Facial	Body Piercing	71
Pleurisy - Related to Infection	Infection - General	129
Pleurisy - Related to Malignancy	Malignancy	141
Pneumococcal Immunisation	Immunisation - Non-Live	125
Pneumonia	Infection - General	129
Pneumothorax	Pneumothorax	161
Podiatry	Chiropody	76
Police Emergency Response Vehicle Driver	Hazardous Activity	108
Polio - Contact With	Non-Contagious Diseases - Contact With	150
Polio - Injected Immunisation	Immunisation - Non-Live	125
Polio - Oral Immunisation	Immunisation - Live	125

For	See	Page
Poliomyelitis - Disability from	Disabled Donor	88
Poliomyelitis - Infection	Infection - Acute	127
Polycystic Kidney Disease	Kidney and Bladder Disease	133
Polycystic Ovary Syndrome	Accept	52
Polycythaemia and Raised Haemoglobin	Polycythaemia and Raised Haemoglobin	161
Polymyalgia Rheumatica	Autoimmune Disease	63
Polyposis Coli	Malignancy	141
Polyps - Malignant	Malignancy	141
Polyps - Nasal	Allergy	56
Polyps - Removal	Surgery	182
Porphyria	Porphyria	162
Porphyria Cutanea Tarda	Porphyria	162
Post natal anti-D	Immunoglobulin Therapy	126
Post Viral Fatigue Syndrome	Chronic Fatigue Syndrome	77
Posterior Pituitary Disorders	Pituitary Disorders	157
Post-exposure Prophylaxis	Pre- and Post-Exposure Prophylaxis for HIV prevention	162
Pramipexole	Central Nervous System Disease	73
Pre- and Post-Exposure Prophylaxis for HIV prevention	Pre- and Post-Exposure Prophylaxis for HIV prevention	162
Pre-cancerous Cells of the Cervix	Cervical Dysplasia	75
Pre-exposure Prophylaxis	Pre- and Post-Exposure Prophylaxis for HIV prevention	162
Pregnancy	Pregnancy	163
Premalignant Conditions	Malignancy	141
PREP	Pre- and Post-Exposure Prophylaxis for HIV prevention	162
Pressure Sore	Wounds, Mouth and Skin Ulcers	203
Primary Biliary Cholangitis	Liver Disease	137
Primary Sclerosing Cholangitis	Liver Disease	137
Prion Associated Diseases	Prion Associated Diseases	164
Proctitis	Proctitis	165
Proctoscopy	Endoscopy	93
Prolactinoma	Pituitary Disorders	157
Propecia®	Anti-Androgens	60
Propecia® - for Hair Loss	Alopecia	57
Prophylactic Antibiotics	Antibiotic Therapy	61
Proscar® - for Hair Loss	Alopecia	57
Proscar® - for Prostate Problems	Anti-Androgens	60
Prostaglandin Injection - for Erectile Dysfunction	Erectile Dysfunction	94
Prostate Cancer	Malignancy	141
Prostate Problems	Prostate Problems	165

For	See	Page
Prostate Specific Antigen	Prostate Problems	165
Prostatectomy	Prostate Problems	165
Prostatic Intraepithelial Neoplasia	Malignancy	141
Prostitute	Sex Worker	171
Protein C Deficiency	Thrombosis and Thrombophilia	186
Protein S Deficiency	Thrombosis and Thrombophilia	186
Prothrombin deficiency	Bleeding Disorder	65
Proton Pump Inhibitors	Acid Indigestion	52
Protopic - for Dermatitis	Dermatitis	85
Protopic - for Skin Disease	Skin Disease	174
Provigil	Narcolepsy	146
PRP Facial	Body Piercing	71
PSA	Prostate Problems	165
Pseudohyperkalaemia	Familial Pseudohyperkalaemia	97
Psoriasis	Psoriasis	166
Psoriatic Arthritis	Psoriasis	166
Psychiatric Problems	Mental Health Problems	143
Public Health Risk of CJD or vCJD	Prion Associated Diseases	164
Public Service Vehicle Driver	Hazardous Activity	108
Pulmonary Embolism	Thrombosis and Thrombophilia	186
Pulmonary Hypertension	Cardiovascular Disease	72
Punch injury	Non-Consented Exposure to Human Body Fluids	149
PUVA	Psoriasis	166
Pyelitis	Infection - General	129
Pyelonephritis	Kidney and Bladder Disease	133
Pyelonephritis	Infection - General	129
Pyrexia	Pyrexia	167
Pyruvate Kinase Deficiency	Haemolytic Anaemia	107
Q		
Q Fever	Infection - Chronic	128
Q Fever - Contact With	Non-Contagious Diseases - Contact With	150
Quinsy	Infection - Acute	127
R		
Rabies	Rabies	168
Rabies Immunisation	Rabies	168
Radiation Therapy	Radiation Therapy	169
Radioactive Iodine Injection	Radionuclides	169
Radionuclides	Radionuclides	169
Raloxifene - for Breast Malignancy Prophylaxis	Malignancy	141
Ranibizumab	Eye Disease	95

For	See	Page
Rape - Male Homosexual	Non-Consented Exposure to Human Body Fluids	149
Rape – Heterosexual	Non-Consented Exposure to Human Body Fluids	149
Raynaud's Syndrome	Raynaud's Syndrome	170
RBBB	Cardiovascular Disease	72
RCVS	Migraine	143
Real time continuous glucose monitor	Shunts, Stents and Devices	173
Recipient of Donated Human Egg	Fertility	97
Recipient of Donated Human Embryo	Fertility	97
Recipient of Donated Human Sperm	Fertility	97
Recipients of Normal Human Immunoglobulin	Immunoglobulin Therapy	126
Regional Anaesthetic	Anaesthetic	59
Relapsing Fever	Infection - Acute	127
Relenza®	Zanamivir	204
Renal Colic	Kidney and Bladder Disease	133
Renal Disease	Kidney and Bladder Disease	133
Renal Failure	Kidney and Bladder Disease	133
Renal Stent	Shunts, Stents and Devices	173
Renal Stent - Removal	Surgery	182
Renal Stones	Kidney and Bladder Disease	133
Repair Vocal Chords	Surgery	182
Respiratory Disease	Respiratory Disease	170
Restless Leg Syndrome	Accept	52
Resurfacing of Hip	Surgery	182
Retinal Detachment - Treated by Cryotherapy	Surgery	182
Retinal Detachment - Treated by Laser	Laser Treatment	136
Retinal Vein Thrombosis	Thrombosis and Thrombophilia	186
Retinitis Pigmentosa	Disabled Donor	88
Revatio® - for Pulmonary Hypertension	Cardiovascular Disease	72
Reversible Cerebral Vasoconstriction Syndrome	Migraine	143
Rheumatic Fever	Rheumatic Fever	170
Rheumatic Fever - Contact With Acute Infection	Infectious Diseases - Contact With	130
Rheumatic Fever - Heart Problems	Cardiovascular Disease	72
Rheumatoid Arthritis	Autoimmune Disease	63
Rhinoplasty	Surgery	182
Rickettsial Infections	Infection - Chronic	128
Right Bundle Branch Block	Cardiovascular Disease	72
Ringworm	Skin Disease	174
Roaccutane®	Acne	53

For	See	Page
Rock Climbing	Hazardous Activity	108
Rocky Mountain Spotted Fever	Infection - Chronic	128
Rodent Ulcer	Malignancy	141
Root Canal Treatment	Dental Treatment	84
Ropinirole	Central Nervous System Disease	73
Rosacea	Acne	53
Rotigotine	Central Nervous System Disease	73
Rubella - Contact With	Infectious Diseases - Contact With	130
Rubella - Immunisation	Immunisation - Live	125
Rubella - Infection	Infection - Acute	127
S		
Salmonella - other than Typhoid & Paratyphoid	Gastroenteritis	101
Salmonella - Typhoid & Paratyphoid	Infection - Chronic	128
Salpingitis	Infection - General	129
Sandfly Fever	Infection - Acute	127
Sarcoidosis	Sarcoidosis	171
SARS	Coronavirus Infection (COVID-19)	81
SARS-CoV-2	Coronavirus Infection (COVID-19)	81
Saxenda®	Donor Weight	90
SBE	Endocarditis	93
Scabies	Skin Disease	174
Scaffolder	Hazardous Activity	108
Scale and polish	Dental Treatment	84
Scarlet Fever - Contact With	Infectious Diseases - Contact With	130
Scarlet Fever - Infection	Infection - Acute	127
Schistosomiasis	Infection - Acute	127
Schizophrenia	Mental Health Problems	143
Sclera Graft Recipient	Tissue and Organ Recipients	189
Scleritis	Eye Disease	95
Scleroderma	Autoimmune Disease	63
Scrub Typhus	Infection - Chronic	128
Scuba Diving	Hazardous Activity	108
Sedation for Minor Procedure	Anaesthetic	59
Sedatives - for Mental Health Problems	Mental Health Problems	143
Sedatives - for Sleeping	Hypnotics	123
SEID	Chronic Fatigue Syndrome	77
Self Catheterisation	Urinary Catheterisation	198
Self-flagellation	Body Piercing	71
Self-Injected Drugs - Drug User	Addiction and Drug Abuse	54

For	See	Page
Self-Injected Drugs - for Erectile Dysfunction	Erectile Dysfunction	94
Semi-Permanent Makeup	Body Piercing	71
Seronegative Arthritis	Autoimmune Disease	63
Severe Acute Respiratory Syndrome	Coronavirus Infection (COVID-19)	81
Sex Assigned at Birth	Transgender and Non-Binary Individuals	192
Sex Change	Transgender and Non-Binary Individuals	192
Sex with a Man who has had Sex with a Man	Blood Safety Entry	68
Sex with a Prostitute	Blood Safety Entry	68
Sex with Anyone Meeting Sub-Saharan African Criteria	Blood Safety Entry	68
Sex with Anyone Who Has Used IV, IM or SC Drugs	Blood Safety Entry	68
Sex Worker	Sex Worker	171
Sexual Assault - male homosexual	Non-Consented Exposure to Human Body Fluids	149
Sexual assault – heterosexual	Non-Consented Exposure to Human Body Fluids	149
Sexual Partner of a Person with Non-Consented Exposure to Human Body Fluids	Non-Consented Exposure to Human Body Fluids	149
Sexually Transmitted Disease	Sexually Transmitted Disease	172
Sharps Injury	Non-Consented Exposure to Human Body Fluids	149
Sheehan's Syndrome	Pituitary Disorders	157
Shingles - Contact With	Infectious Diseases - Contact With	130
Shingles - Immunisation	Immunisation - Non-Live	125
Shingles - Immunisation	Immunisation - Live	125
Shingles - Infection	Infection - Acute	127
Shunts, Stents and Devices	Shunts, Stents and Devices	173
SIADH	Pituitary Disorders	157
Sickle-Cell	Haemoglobin Disorders	105
Sickle-Cell Trait	Sickle-Cell Trait	174
Sigmoidoscopy	Endoscopy	93
Sildenafil - for Erectile Dysfunction	Erectile Dysfunction	94
Sildenafil - for Pulmonary Hypertension	Cardiovascular Disease	72
Sinusitis	Infection - General	129
Sjogren's Syndrome	Autoimmune Disease	63
Skin Cancer	Malignancy	141
Skin Creams	Topical Medication	190
Skin Disease	Skin Disease	174
Skin Lesion Removed	Surgery	182
Skin Piercing	Body Piercing	71
Skin Tag Removal	Surgery	182
Skin Ulcer	Wounds, Mouth and Skin Ulcers	203
Sky Diving	Hazardous Activity	108

For	See	Page
Slapped Cheek Syndrome	Parvovirus B19	155
SLE	Autoimmune Disease	63
Sleep Apnoea	Sleep Apnoea	175
Sleeping Sickness	Infection - Chronic	128
Sleeping Tablets	Hypnotics	123
Slimming Tablets	Donor Weight	90
Slipped Disc Operation	Back Problems	64
Slow Heart Beat	Arrhythmias	62
Smallpox Immunisation	Smallpox Immunisation	176
Smallpox Immunisation - Contact	Smallpox Immunisation	176
Smear Test - Abnormal	Cervical Dysplasia	75
Smoking	Smoking	177
Smoking Cessation	Smoking Cessation	177
Snake Bite	Animal Bite (Non-Human)	59
Snip	Surgery	182
Sore Throat	Infection - General	129
South American Trypanosomiasis	South American Trypanosomiasis	177
South American Trypanosomiasis - Contact With	Non-Contagious Diseases - Contact With	150
South American Trypanosomiasis - Infection Risk	South American Trypanosomiasis	177
South American Trypanosomiasis - plasmapheresis donors	South American Trypanosomiasis - plasmapheresis donors	178
Spherocytosis	Haemolytic Anaemia	107
Spina Bifida	Spina Bifida	179
Spinal - Neurosurgery	Neurosurgery	148
Spinal - Orthopaedic Surgery	Surgery	182
Splenectomy	Splenectomy	179
Sporadic CJD	Prion Associated Diseases	164
Sport	Exercise	95
Spots - Acne	Acne	53
Spots - not Acne	Infection - General	129
Sprained Ankle	Accident	52
Squamous cell carcinoma in situ (CIS) of the skin	Malignancy	141
STD	Sexually Transmitted Disease	172
Stem Cell Donation Frequency	Frequency of Donation	99
Sterilisation	Surgery	182
Steroid Cream	Steroid Therapy	180
Steroid Inhaler	Steroid Therapy	180
Steroid Tablets	Steroid Therapy	180
Steroid Therapy	Steroid Therapy	180

For	See	Page
Steroids for Bodybuilding	Addiction and Drug Abuse	54
Steven's Johnson Syndrome	Allergy	56
Stoma	Stoma	181
Stomach Cancer	Malignancy	141
Stomach Ulcer - Malignant	Malignancy	141
Stomach Ulcer - Non-Malignant	Peptic Ulcer	156
Stroke	Cerebrovascular Disease and Intracranial Haemorrhage	74
Stye	Infection - Acute	127
Subacute Bacterial Endocarditis	Endocarditis	93
Subarachnoid Haemorrhage	Cerebrovascular Disease and Intracranial Haemorrhage	74
Sugar Diabetes	Diabetes Mellitus	86
Sulfasalazine (Arthritis)	Autoimmune Disease	63
Superficial Thrombophlebitis	Superficial Thrombophlebitis	181
Superficial Vein Thrombosis	Superficial Thrombophlebitis	181
Supraventricular Tachycardia	Arrhythmias	62
Surgery	Surgery	182
SVT	Arrhythmias	62
Syncope	Faints	96
Syndrome of Inappropriate Anti-Diuretic Hormone	Pituitary Disorders	157
Syphilis	Syphilis	184
Syphilis Sexual Contact	Syphilis	184
Syringomyelia	Neurosurgery	148
Systemic Exertion Intolerance Disease	Chronic Fatigue Syndrome	77
Systemic Lupus Erythematosus	Autoimmune Disease	63
T		
T. Cruzi	South American Trypanosomiasis	177
Tachycardia	Arrhythmias	62
Tacrolimus - for Skin Disease	Skin Disease	174
Tacrolimus - for Dermatitis	Dermatitis	85
Tamiflu®	Oseltamivir	152
Tamoxifen - for Cancer	Malignancy	141
Tamoxifen - for Infertility	Fertility	97
Tamoxifen – for Breast Malignancy Prophylaxis	Malignancy	141
Tattoo	Body Piercing	71
TB	Tuberculosis	197
TB Immunisation	Immunisation - Live	125
Temperature - High	Pyrexia	167
Temporal Arteritis	Autoimmune Disease	63
Tendinitis	Tendonitis	185

For	See	Page
Tendonitis	Tendonitis	185
Tennis Elbow	Tendonitis	185
Tension Free Vaginal Tape Operation	Surgery	182
Termination of Pregnancy	Pregnancy	163
Testicular Cancer	Malignancy	141
Testosterone replacement therapy	Hormone Replacement and Sex Hormone Therapy	120
Tests	Investigations	132
Tetanus	Tetanus	185
Tetanus - Contact With	Non-Contagious Diseases - Contact With	150
Tetanus - Immunisation	Tetanus	185
Tetracycline - for Acne	Acne	53
Tetracycline - for Other Infections	Infection - General	129
Thalassaemia Trait	Haemoglobin Disorders	105
The Bends	Decompression Illness	83
The Pill	Contraceptive Use	81
Thiazide	Diuretics	89
Threadworms	Threadworms	186
Three Day Fever	Infection - Acute	127
Thrombocythaemia	Haematological Disease	103
Thrombocytopenia - Immune	Immune Thrombocytopenia	123
Thrombocytopenia - Non-Immune	Platelet Count	158
Thrombocytosis	Platelet Count	158
Thrombophilia	Thrombosis and Thrombophilia	186
Thrombophlebitis	Superficial Thrombophlebitis	181
Thrombosis and Thrombophilia	Thrombosis and Thrombophilia	186
Thrombotic Thrombocytopenic Purpura	Thrombosis and Thrombophilia	186
Thrush	Thrush	188
Thyroid Disease	Thyroid Disease	189
Thyroidectomy - for Hyperthyroidism	Thyroid Disease	189
Thyroidectomy - for Malignancy	Malignancy	141
Thyrotoxicosis	Thyroid Disease	189
Thyroxine	Thyroid Disease	189
TIAs	Cerebrovascular Disease and Intracranial Haemorrhage	74
Tick-Borne Encephalitis - Contact With	Non-Contagious Diseases - Contact With	150
Tick-Borne Encephalitis - Immunisation	Immunisation - Non-Live	125
Tick-Borne Encephalitis - Infection	Infection - Acute	127
Tigason®	Etretinate	95
Tinea - Fungal Infection	Skin Disease	174

For	See	Page
Tissue and Organ Recipients	Tissue and Organ Recipients	189
Toctino® - Dermatitis	Dermatitis	85
Toctino® - Skin Disease	Skin Disease	174
Toenail Infection	Infection - General	129
Toenail Removed	Surgery	182
Tongue Piercing	Body Piercing	71
Tonsillectomy	Surgery	182
Tonsillitis	Infection - General	129
Too Soon After Donation	Frequency of Donation	99
Tooth Extraction	Dental Treatment	84
Topical Medication	Topical Medication	190
Topiramate	Valproate and Topiramate	198
Tourette's Syndrome	Neurobehavioral Disorders	148
Toxoplasmosis	Toxoplasmosis	191
Tracheitis	Infection - General	129
Tranquilisers	Mental Health Problems	143
Transfusion	Transfusion	191
Transgender and Non-Binary Individuals	Transgender and Non-Binary Individuals	192
Transient Global Amnesia	Central Nervous System Disease	73
Transient Ischaemic Attacks	Cerebrovascular Disease and Intracranial Haemorrhage	74
Transition	Transgender and Non-Binary Individuals	192
Transverse Myelitis	Central Nervous System Disease	73
Trapped Nerve Released	Surgery	182
Trauma	Accident	52
Travel	Travel	193
Treatment with Blood Components	Transfusion	191
Trigeminal Neuralgia - Drug Treatment	Pain Killers	154
Trigeminal Neuralgia -Requiring Surgery	Neurosurgery	148
Tropical Areas	Travel	193
Tropical Disease	Infection - General	129
Tropical Viruses	Tropical Viruses	194
Tropical Viruses - plasmapheresis donors	Tropical Viruses - plasmapheresis donors	195
Truvada®	Pre- and Post-Exposure Prophylaxis for HIV prevention	162
Trying for a Baby	Fertility	97
Trypanosoma Cruzi - Infection	Infection - Chronic	128
Trypanosoma Cruzi - Risk Through Exposure	South American Trypanosomiasis	177
TSH Secreting Tumour	Pituitary Disorders	157
TTP	Thrombosis and Thrombophilia	186

For	See	Page
Tuberculin PPD Test	Tuberculin PPD Test	196
Tuberculosis	Tuberculosis	197
Tuberculosis - Contact With	Tuberculosis	197
Tuberculosis Immunisation	Immunisation - Live	125
Tummy Tuck	Surgery	182
Tumour - Malignant	Malignancy	141
Tumour - Non-Malignant	Surgery	182
Tumour Chemotherapy	Malignancy	141
Turner's Syndrome	Turner's Syndrome	197
TVT Operation	Surgery	182
Type I (1) and II (2) Diabetes	Diabetes Mellitus	86
Types 1, 2 or 3 Von Willebrand disease	Bleeding Disorder	65
Typhoid	Infection - Chronic	128
Typhoid - Injected Immunisation	Immunisation - Non-Live	125
Typhoid - Oral Immunisation	Immunisation - Live	125
Typhoid and Hepatitis A - Immunisation	Immunisation - Non-Live	125
Typhus	Infection - Chronic	128
U		
Ulcer - Gullet, Stomach or Duodenum	Peptic Ulcer	156
Ulcer - Skin/Mouth	Wounds, Mouth and Skin Ulcers	203
Ulcerative Colitis	Inflammatory Bowel Disease	131
Unconscious	Accident	52
Under Weight	Donor Weight	90
Underactive Thyroid	Thyroid Disease	189
Undergoing Tests	Investigations	132
Undulant Fever - Contact With	Non-Contagious Diseases - Contact With	150
Undulant Fever - Infection	Infection - Chronic	128
Unhealed Wound	Wounds, Mouth and Skin Ulcers	203
Upset Tummy	Gastroenteritis	101
Ureteric Stent	Shunts, Stents and Devices	173
Ureteric Stent - Removal	Surgery	182
Urethral Stricture - Dilatation	Surgery	182
Urethral Stricture - Needing to Self Catheterise	Urinary Catheterisation	198
Urethritis Acute	Infection - Acute	127
Urinary Catheterisation	Urinary Catheterisation	198
Urinary Catheterisation	Urinary Catheterisation	198
Urinary Incontinence - Catheter	Urinary Catheterisation	198
Urinary Retention - Catheter	Urinary Catheterisation	198
Urinary Tract Infection	Infection - General	129

For	See	Page
Urostomy	Stoma	181
Urticaria	Allergy	56
UTI	Infection - General	129
V		
Vaccination	Immunisation	124
Vaccination - Live	Immunisation - Live	125
Vaccination - Non-Live	Immunisation - Non-Live	125
Vaginal Repair	Surgery	182
Valproate and Topiramate	Valproate and Topiramate	198
Valproic	Valproate and Topiramate	198
Valve Surgery	Cardiac Surgery	71
Vampire Facial®	Body Piercing	71
Varenicline	Smoking Cessation	177
Variant Creutzfeldt-Jakob Disease	Prion Associated Diseases	164
Varicella - Contact With	Infectious Diseases - Contact With	130
Varicella - Immunisation	Immunisation - Live	125
Varicella - Infection	Infection - Acute	127
Varicose Ulcer	Wounds, Mouth and Skin Ulcers	203
Varicose Vein Surgery	Surgery	182
Varicose Veins and Chronic Venous Insufficiency	Varicose Veins and Chronic Venous Insufficiency	199
Varicose Veins Injection	Varicose Veins and Chronic Venous Insufficiency	199
Varigate Porphyria	Porphyria	162
Vascular Dementia	Cerebrovascular Disease and Intracranial Haemorrhage	74
Vasculitis	Autoimmune Disease	63
Vasectomy	Surgery	182
Vasodilators - for Blood Pressure	Blood Pressure - High	67
Vasodilators - for Cardiovascular Disease	Cardiovascular Disease	72
Vasodilators - for Erectile Dysfunction	Erectile Dysfunction	94
Vasodilators - for Raynaud's Syndrome	Raynaud's Syndrome	170
v		
vCJD	Prion Associated Diseases	164
V		
Venereal Disease	Sexually Transmitted Disease	172
Venesection - for Haemochromatosis	Haemochromatosis	104
Venesection - for Polycythaemia	Polycythaemia and Raised Haemoglobin	161
Venous Thrombosis	Thrombosis and Thrombophilia	186
Ventricular Tachycardia	Arrhythmias	62
Ventriculoperitoneal Shunt	Shunts, Stents and Devices	173
Verruca	Warts	201
Vertigo	Vertigo	199
Vestibular Schwannoma – Causing Vertigo	Vertigo	199

For	See	Page
Vestibular Schwannoma – Surgery	Neurosurgery	148
Viagra® - for Erectile Dysfunction	Erectile Dysfunction	94
Viagra® - for Pulmonary Hypertension	Cardiovascular Disease	72
Vibration White Finger	Raynaud's Syndrome	170
Victoza®	Diabetes Mellitus	86
Viral Disease	Infection - General	129
Viral Haemorrhagic Fever	Viral Haemorrhagic Fever	200
Viral Hepatitis	Hepatitis	110
Viral Pericarditis	Infection - General	129
Viridal Duo®	Erectile Dysfunction	94
Vitamins and Other Nutritional Supplements	Vitamins and Other Nutritional Supplements	200
Vitiligo	Autoimmune Disease	63
Vomiting and Diarrhoea	Gastroenteritis	101
Von Recklinghausen's Disease	Neurofibromatosis	148
Von Willebrand disease	Bleeding Disorder	65
Vulval Carcinoma in Situ	Malignancy	141
W		
Waiting for Operation	Surgery	182
Waiting for Results of Tests	Investigations	132
Warfarin - for Abnormal Heart Rhythm	Arrhythmias	62
Warfarin - to stop Blood Clotting	Anticoagulant Therapy	61
Warts	Warts	201
Water on the Brain	Hydrocephalus	122
Water Tablets	Diuretics	89
Waxing	Hair Removal	108
Weight Loss Injections	Donor Weight	90
Weight Loss Pills	Donor Weight	90
Weight Problems	Donor Weight	90
Weil's Disease	Infection - Acute	127
West Nile Virus	West Nile Virus	202
West Nile Virus - Contact With	Non-Contagious Diseases - Contact With	150
West Nile Virus - plasmapheresis donors	West Nile Virus - plasmapheresis donors	202
Wet Cupping	Complementary Therapy	80
Wheelchair User	Disabled Donor	88
White Finger Syndrome	Raynaud's Syndrome	170
Whooping Cough - Contact With	Infectious Diseases - Contact With	130
Whooping Cough - Immunisation	Immunisation - Non-Live	125
Whooping Cough - Infection	Infection - Acute	127
Wilson's Disease	Liver Disease	137

For	See	Page
Wolff-Parkinson-White Syndrome	Arrhythmias	62
Working at Heights	Hazardous Activity	108
Wounds, Mouth and Skin Ulcers	Wounds, Mouth and Skin Ulcers	203
WPW	Arrhythmias	62
X		
Xenografts	Xenotransplantation	203
Xenotransplantation	Xenotransplantation	203
XMRV	XMRV	204
Y		
Yaws	Infection - Chronic	128
Yeast	Thrush	188
Yellow Fever - Contact With	Non-Contagious Diseases - Contact With	150
Yellow Fever - Infection	Tropical Viruses	194
Yellow Fever Immunisation	Immunisation - Live	125
YF	Tropical Viruses	194
Z		
Zanamivir	Zanamivir	204
Zika Virus	Tropical Viruses	194
Zoonosis	Non-Contagious Diseases - Contact With	150
Zoonotic infection	Non-Contagious Diseases - Contact With	150
Zyban®	Smoking Cessation	177

Accept

<i>Discretionary</i>	If use of medicines or other therapies are not a reason to defer, accept.
<i>See if Relevant</i>	<p><u>Complementary Therapy</u> <u>Drug Index - preparations which may affect platelet function</u> <u>Fertility</u> <u>Nonsteroidal Anti-Inflammatory Drugs</u> <u>Neurosurgery</u></p> <p>If taking Dopamine-receptor agonists (e.g. Rotigotine, Bromocriptine, Ropinirole, Pramipexole), see: <u>Central Nervous System Disease</u></p>
<i>Additional Information</i>	The condition that brought you to this entry is not a reason for deferral. It is however important to ensure that there are no other factors that may affect any donation, such as having symptoms related to hypotension as side effects from the use of medicines such as dopamine receptor agonist drugs (e.g. rotigotine, bromocriptine, ropinirole and pramipexole) that can cause hypotension and fainting, or complementary (alternative) therapy.
<i>Reason for change</i>	Fertility has been added to See if Relevant section.
<i>Update Information</i>	This entry was last updated in: WB-DSG Edition 203 Release 72

Accident

<i>Includes</i>	Fracture, head injury and trauma.
<i>Obligatory</i>	<p>Must not donate if:</p> <p>a) Any wound is not fully healed.</p> <p>b) Has any infection.</p> <p>c) Has a plaster-cast.</p>
<i>See if Relevant</i>	<p><u>Disabled Donor</u> <u>Epilepsy</u> <u>Infection - General</u> <u>Neurosurgery</u> <u>Surgery</u> <u>Tetanus - 2. Immunization</u> <u>Transfusion</u></p>
<i>Additional Information</i>	<p>An unhealed wound or sore is a risk for bacteria entering the blood. Bacteria can be a serious threat to anybody receiving blood or blood components. This is because bacteria can multiply to dangerous levels after collection.</p> <p>A plaster-cast can hide a wound or sore.</p>
<i>Reason for change</i>	<p>The previous entry unduly restricted individuals who had suffered an accident from donating by requiring them to be 'recovered'. An example would have been inappropriately preventing a person from donating because of a sprained ankle.</p> <p>Links have been added to 'Disabled Donor', 'Epilepsy' and 'Infection - General'.</p>
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 01.

Acid Indigestion

<i>Includes</i>	Acid reflux, gastritis, gastro-oesophageal reflux disorder (GORD), heartburn, hiatus hernia and indigestion.
<i>Obligatory</i>	Must not donate if: Waiting for investigations or results of investigations.
<i>Discretionary</i>	If symptoms are relieved by regular or sporadic use of medication and in good health, accept.
<i>See if Relevant</i>	<u>Endoscopy</u> <u>Malignancy</u> <u>Peptic Ulcer</u> <u>Surgery</u>
<i>Additional Information</i>	It is important only to accept people in good health. Where the cause of symptoms is not known and the individual is either waiting for investigations or the results of investigations, donation must be deferred. This is because the cause of the symptoms may be a reason for deferral.
<i>Reason for change</i>	The previous entry for 'Gastritis' has been revised and renamed 'Acid Indigestion'. Links and 'Additional Information' have been added.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 01.

Acne

<i>Includes</i>	Acne rosacea.
<i>Definitions</i>	With regard to drug treatment: Topical: Applied to the skin only. Systemic: Taken by mouth or other routes so that it has an affect on the whole body.
<i>Obligatory</i>	Must not donate if: a) Has ever taken etretinate (Tigason®). b) Less than 36 months from the last dose of acitretin (Neotigason®). c) Less than four weeks from the last dose of isotretinoin (Roaccutane®). d) There is secondary infection. e) Less than seven days from completing systemic antibiotic treatment for secondary infection.
<i>Discretionary</i>	If using topical treatments (including retinoids), or taking oral antibiotics, diuretics (such as spironolactone) or oral co-cyprindiol (Dianette® (cyproterone acetate and ethinylestradiol)), accept.
<i>Additional Information</i>	Etretinate (Tigason®), acitretin (Neotigason®) and isotretinoin (Roaccutane®) taken systemically can cause birth defects in babies exposed to them while inside the womb. It is important to allow time for the drug to be cleared from the blood of a donor. It takes longer to clear some drugs than others. There is no published data that topical retinoids cause birth defects. Secondary infection of acne is usually obvious with swelling and redness of affected spots. There is a risk of bacteria entering the blood. This could be a serious threat to anybody receiving blood or blood components. This is because bacteria can multiply to dangerous levels after collection.
<i>Reason for change</i>	The deferral period after acitretin therapy has increased from 24 to 36 months.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 55.

Addiction and Drug Abuse

Includes Alcohol, body building drugs and injected non-prescribed drugs.

Obligatory **Must not donate if:**

a) Has ever injected, or has been injected with, drugs; even a long time ago or only once. This includes bodybuilding drugs, injected tanning agents and injected chemsex drugs.

b) Adversely affected by any drug, including alcohol, which may affect the process of obtaining valid consent.

c) Less than seven days from taking disulfiram (Antabuse®).

Discretionary a) If any injected drugs were prescribed for the donor by a registered health care professional for a condition that would not lead to exclusion, accept.

b) If the donor is taking medication to support their abstinence from alcohol or other non-injected drugs and

- they are not adversely affected by drugs, including alcohol, and
- they understand and consent to the donation process and testing of their blood, accept.

See if Relevant Blood Safety Entry

For alcohol related problems:

Liver Disease

Additional Information Injecting drug users represent one of the groups of individuals within whom emerging infections have spread before they have been recognized. This was the case with HIV and HCV infection. Because of this, the BSQR requires that they are permanently excluded from becoming donors. It can be many years before any infection shows itself. Former drug users often do not realize that they can pass infection on to others many years after they last used drugs themselves.

Previous use of non-injected drugs does not necessarily require exclusion.

Anyone obviously affected by alcohol, or other drugs that can affect the mind, cannot give valid consent or fully understand why they are being asked certain questions. They can be a danger to themselves and to others. If the donor is deferred, this may be until the next session, or permanently, if the donor's use of alcohol and/or drugs is likely to continue.

Disulfiram (Antabuse®) may cause severe reactions in a recipient whose blood contains alcohol.

Other medications such as Acamprosate (Campral®) or Naltrexone may be prescribed to support abstinence from alcohol or drug use. If the donor is well and their alcohol or drug use has not caused any end-organ damage, then they can be accepted to donate.

Information Part of this entry is a requirement of the Blood Safety and Quality Regulations 2005.

Reason for change This entry has been revised to include guidance on the acceptance of donors who are prescribed medication to support their abstinence from alcohol. Other revisions to clarify the text.

Update Information This entry was last updated in:
DSG-WB Edition 203, Release 65

Adrenal Failure

Includes Addison's disease.

Obligatory **Must not donate.**

<i>Additional Information</i>	Adrenal failure is due to the adrenal glands producing insufficient steroid hormones to maintain health. There are many causes, including autoimmune disease and infection. Affected individuals take replacement steroid hormones. The dose of these must be increased during times of stress. It is considered that taking blood from people with adrenal failure may put them at unnecessary risk.
<i>Reason for change</i>	The title has been changed from 'Addison's Disease' to 'Adrenal Failure' and 'Additional Information' has been added.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 01.

Age

<i>Definitions</i>	<p>First Time Donor Is an individual who has not previously donated. It is also a person who has previously attended but, for whatever reasons, did not give a full donation (e.g. deferred because of selection guidelines, failed screen test, failed venepuncture, part bag etc).</p> <p>Regular Donor Is a donor who has been medically assessed at a donor session in the last 24 months. For component donors this must include mandatory infection screening and, for all donors over the age of 72 years, this must also include giving a full donation.</p> <p>Returning Donor Is a donor who has not attended a donation session or been medically assessed within the last 24 months, but who has previously given a full donation.</p> <p>Full Donation An amount above the minimum required volume has been collected.</p>
<i>Obligatory</i>	<p>Whole blood and component donors. Must not donate if:</p> <ul style="list-style-type: none"> a) They are under 17 years of age. b) They are a first time donor who has had their 66th birthday. c) They are a returning donor who has had their 72nd birthday.
<i>Additional Information</i>	<p>The lower age limit takes account of national laws on age of consent.</p> <p>Upper age limits for blood and component donation have traditionally been set to protect the donor's safety. There is however little evidence to support this. Donor haemovigilance data show a decreased incidence of adverse events in older donors compared to younger donors, although there is an increase in local complications of donating (i.e. bruising and rebleeds). Donor adverse event monitoring will continue to inform the need for any modification to this guidance.</p> <p>To donate after their 72nd birthday a donor must remain in good health and have given at least one full donation in the previous 24 months. To continue donating they must give no less than one full donation every 24 months.</p> <p>When appropriate, donors may be accepted on their birthday.</p>
<i>Information</i>	This entry is compliant with the Blood Safety and Quality Regulations 2005.
<i>Reason for change</i>	The age limit for returning donors has been increased to 72 years.
<i>Update Information</i>	This entry was last updated in: WB-DSG Edition 203 Release 72

Air Crew and Air Traffic Controllers

Includes

Air crew:
 Flight crew
 Cabin crew
 Military Aircrew
 Military Supernumerary crew

Air Traffic Controllers:
 Civilian controllers
 Military controllers

Obligatory **Air crew (except military) and Air Traffic Controllers (except military):
 Must not donate if:**
 On duty within the next 24 hours.

**Military air crew:
 Must not donate if:**
 On duty within the next 36 hours.

**Military controllers:
 Must not donate if:**
 On duty within the next 12 hours.

Discretionary **Non-military crew/controllers holding Class 3 medical certificates:**
 If not on duty within the next 12 hours, accept.

Additional Information The UK Civil Aviation Authority (CAA) guidelines state:

'In order to prevent the very slight risk of post-donation faintness or syncope, donating blood or plasma should be avoided during the 24 hours before duty for holders of Class 1 and 2 medical certificates, and during the 12 hours before duty for holders of Class 3 medical certificates.'

There is no CAA guidance for crew that are not required to hold these types of medical certification. Donors to whom this applies, including cabin crew and some private pilots, should be advised not to donate if they are on duty within the next 24 hours in line with commercial pilots and other flight crew.

The Ministry of Defence (MOD) guidelines state:
 'Aircrew and Supernumerary Crew should not fly until 36 hours have elapsed after donating blood.'
 and
 'Military and MOD Contracted Civilian Controllers should not control until 12 hours have elapsed after donating blood.'

Reason for change Updated Civil Aviation Authority advice for holders of Class 3 medical certificates has been added.

Update Information This entry was last updated in:
 DSG-WB Edition 203, Release 69

Allergy

Includes Allergic rhinitis, allergic conjunctivitis, anaphylaxis, hay fever and urticaria.

Obligatory **Ensure:**
 Procedures will not expose the donor to something they are allergic to, e.g. constituents of the arm cleaning preparation.

Must not donate if:
 a) Has severe symptoms due to an allergy or to the medication they are taking.
 b) The donor has taken oral or injected steroids within the last seven days.

<i>Discretionary</i>	<p>a) Donors taking medication, other than oral or injected steroids (including antihistamines, eye drops or intranasal steroids e.g. beclometasone (Beconase[®])), or with a history of allergy or anaphylaxis (including those who carry adrenaline/epinephrine for self administration, e.g. Anapen[®] or EpiPen[®]), provided they are well on the day and will not be exposed to anything they are allergic to, accept.</p> <p>b) Nickel allergy, accept.</p> <p>c) Donors undergoing desensitisation treatment for hay fever or for allergy to an insect sting, providing they are not experiencing any systemic or local reactions to the treatment, and are well on the day, accept.</p>
<i>See if Relevant</i>	<p><u>Asthma</u> <u>Dermatitis</u> <u>Steroid Therapy</u></p>
<i>Additional Information</i>	<p>Any person who is unwell should not be accepted as a donor. This is to ensure that a serious underlying condition, that could be a risk either to the donor or to a potential recipient, is not missed. For this reason, a potential donor with anything other than minor symptoms related to an allergy should not be accepted.</p> <p>Severe systemic reactions are not seen with nickel 'allergy'.</p>
<i>Reason for change</i>	<p>The list of potential allergens has been reworded. A reference to allergic conjunctivitis has been added. Clarification of the acceptance criteria for desensitisation treatment has been added. The link to 'Coeliac Disease' has been removed.</p>
<i>Update Information</i>	<p>This entry was last updated in: DSG-WB Edition 203, Release 61.</p>

Alopecia

<i>Includes</i>	<p>Baldness and hair loss treatments.</p>
<i>Obligatory</i>	<p>Must not donate if:</p> <p>a) Dutasteride (Avodart[®]) taken in the last six months.</p> <p>b) Finasteride (Propecia[®], Proscar[®]) taken in the last four weeks.</p> <p>c) Taking systemic anti-fungal treatment or oral steroids.</p> <p>d) Related to malignancy or to its treatment.</p>
<i>Discretionary</i>	<p>a) If the donor is on no treatment, accept.</p> <p>b) If the donor is only using topical treatment and is not attending specialist follow up, accept.</p> <p>c) If the donor is only taking oral Hydroxychloroquine, Spironolactone or Minoxidil, and is not under specialist follow up, accept.</p> <p>d) If the donor has recovered from hair transplant surgery and no further treatment or follow up is planned, accept.</p> <p>e) For all other cases, refer to a DCSO.</p>
<i>See if Relevant</i>	<p>For systemic anti-fungal treatment: <u>Infection - Chronic</u></p> <p><u>Malignancy</u></p> <p>For hair transplants: <u>Surgery</u></p> <p>For injected or oral steroid treatment: <u>Steroid Therapy</u></p>

<i>Additional Information</i>	<p>Hair loss can be related to several factors, including family history, hormone changes, scalp infections, medication and underlying medical disorders.</p> <p>Alopecia areata is an autoimmune condition which can vary in severity, from patchy to complete hair loss. Individuals may require local treatment, including injected steroids, or systemic treatment with steroids and/or immunosuppressants.</p> <p>Dutasteride and finasteride can cause abnormal development of the sexual organs of a male baby within the womb. As it is not possible to know if an individual donation may be transfused to a pregnant woman, whose baby may be at risk, donations cannot be taken from people who may have one of these drugs in their blood. They remain in the blood even after treatment has stopped.</p> <p>Given the range of causes and treatments, referral to a DCSO may be required to establish the donor's eligibility to donate blood.</p>
<i>Reason for change</i>	This entry has been revised by adding a discretion for oral Spironolactone and Minoxidil therapy. Further guidance has been added with regard to immunosuppressive treatments that may be required.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 61.

Anaemia

<i>Includes</i>	Iron deficiency, iron treatment, folate deficiency and pernicious anaemia.
<i>Obligatory</i>	<p>Must not donate if:</p> <p>a) Fails the haemoglobin screen test.</p> <p>b) Under investigation or on treatment for anaemia.</p>
<i>Discretionary</i>	<p>History of anaemia: This must be assessed regarding its cause, current status and what treatment has been received.</p> <p>1. Iron deficiency:</p> <p>a) If not under investigation and the underlying cause is not a reason to exclude, accept.</p> <p>b) If following treatment to cure anaemia, the donor is taking medication to prevent recurrence, accept.</p> <p>2. Other types:</p> <p>a) Medication to prevent recurrence, as opposed to treat anaemia (e.g. B12 for treated pernicious anaemia or folic acid for treated folate deficiency), accept.</p> <p>b) 'See if Relevant' conditions below.</p> <p>c) In other cases: Refer to a 'Designated Clinical Support Officer'.</p>
<i>See if Relevant</i>	<p><u>Haemoglobin Disorders</u> <u>Haemoglobin Estimation</u> <u>Haemolytic Anaemia</u> <u>Kidney and Bladder Disease</u> <u>Malignancy</u></p> <p>If treated with blood components or blood products or by plasma exchange or filtration: <u>Transfusion</u></p>
<i>Additional Information</i>	Donating blood will lower the haemoglobin concentration. People with a history of anaemia may not be able to make up this loss as easily as others.
<i>Information</i>	Part of this entry is a requirement of the Blood Safety and Quality Regulations 2005.

<i>Reason for change</i>	Additional links have been added together with specific mention of pernicious anaemia and folate deficiency. There have been other minor changes to improve clarity.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 22

Anaesthetic

<i>Includes</i>	General anaesthetic, local anaesthetic, regional anaesthetic and sedation for minor procedures.
<i>Obligatory</i>	<p>Must not donate if:</p> <p>a) The underlying condition for which the anaesthetic or sedation was given is not acceptable.</p> <p>b) Less than 24 hours since the anaesthetic or sedation was administered.</p>
<i>See if Relevant</i>	<p><u>Accident</u></p> <p><u>Dental Treatment</u></p> <p><u>Endoscopy</u></p> <p><u>Infection - General</u></p> <p><u>Surgery</u></p> <p><u>Transfusion</u></p>
<i>Additional Information</i>	A longer deferral period may be required due to the nature of the procedure or the underlying condition. Procedures requiring local anaesthetic will normally require a longer deferral period due to any associated infection risk. Treatment requiring any degree of sedation should be followed by a minimum deferral period of 24 hours. This is to ensure that consent and the response to questions can be considered valid.
<i>Reason for change</i>	A link has been added to 'Infection - General'. The 'Additional Information' has been modified.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 01.

Animal Bite (Non-Human)

<i>Includes</i>	Snake bite.
<i>Obligatory</i>	<p>Must not donate if:</p> <p>a) Ever bitten by a non-human primate (monkeys and apes).</p> <p>b) Less than 12 months since being bitten by a bat or exposed to bat saliva anywhere in the world.</p> <p>c) Less than 12 months since being bitten by any other mammal outside of the British Isles (UK and Ireland).</p> <p>d) Any wound is infected or not healed.</p> <p>e) The donor is not fully recovered.</p>
<i>See if Relevant</i>	<p><u>Immunisation</u></p> <p><u>Infection - General</u></p> <p><u>Rabies</u></p> <p><u>Surgery</u></p>

For a human bite:**Non-Consented Exposure to Human Body Fluids***Additional Information*

Animal bites may result in many different infections. Allowing all wounds to heal and for any obvious infection to have resolved should avoid problems. As well as local care of the bite, donors may receive other treatments such as antibiotics or immunisation; these may affect the donor's eligibility.

Bites from poisonous (venomous) snakes can cause extensive local tissue damage that can take a long time to heal and may require surgery. Depending on the species of biting snake, their venom can cause problems with blood clotting and, in severe cases, require transfusion of blood components (e.g. plasma or cryoprecipitate).

There is a concern that bites from non-human primates, because of close genetic links, may transmit diseases that could cause illness in people. It is known that some diseases have been transmitted by this route. For this reason any person who has ever been bitten by a non-human primate is not allowed to donate. Non-human primates include chimpanzees, gorillas, orangutans, monkeys (old and new world), tarsiers, lemurs and lorises.

Anyone who has been in unusual contact with a bat, such as handling a sick or injured bat, should be considered at risk of rabies. Bat bites are usually insignificant and easily overlooked. Merely being in a place where bats roost is not considered a risk.

Rabies, and similar diseases, have long incubation periods and do not show as a wound infection. There is no evidence that these infections have ever been transmitted through a blood transfusion. These diseases appear to be confined to the nervous system during their incubation periods. There is evidence that they have been transmitted through organ, tissue and ocular transplants.

Reason for change

Guidance on snake bites is now included in this entry. A reminder to assess treatments beyond local wound care has been added to the additional information section. The see if relevant section has been updated.

Update Information

This entry was last updated in:
WB-DSG Edition 203 Release 72

Anti-Androgens*Includes*

Bicalutamide (Casodex[®]), cyproterone acetate (Androcur[®], Cyprostat[®]), dutasteride (Avodart[®]), finasteride (Propecia[®], Proscar[®]) and flutamide (Drogenil[®]).

*Obligatory***Must not donate if:**

- a) Dutasteride (Avodart[®]) taken in the last six months.
- b) Finasteride (Propecia[®], Proscar[®]) taken in the last four weeks.
- c) Bicalutamide (Casodex[®]), cyproterone acetate (Androcur[®], Cyprostat[®]) or flutamide (Drogenil[®]) has been taken for a malignant condition.

Discretionary

1. Donors taking cyproterone acetate for non-malignant conditions, if not affected by the Blood Safety Entry, accept.
2. Donors using topical anti-androgen treatments for alopecia, including male pattern baldness, accept.

See if Relevant

Acne
Blood Safety Entry
Hair Removal
Malignancy
Prostate Problems

Additional Information

Dutasteride and finasteride can cause abnormal development of the sexual organs of a male baby within the womb. As it is not possible to know if an individual donation may be

transfused to a pregnant woman whose baby may be at risk, donations cannot be taken from people who may have one of these drugs in their blood. They remain in the blood even after treatment has stopped.

Cyproterone acetate (particularly in the form of Androcur®) may be used to treat male hypersexuality. In such cases a sensitive exploration of any relevant issues dealt with by the Blood Safety Entry should be undertaken.

Reason for change A discretion to accept donors using topical anti-androgens has been added.

Update Information This entry was last updated in:
WB-DSG Edition 203 Release 74

Antibiotic Therapy

See **If on antibiotics to treat an infection:**
Infection - General

See if Relevant **If on prophylactic antibiotics:**
Acne
Infection - General
Infectious Diseases – Contact With
Kidney and Bladder Disease
Splenectomy

Additional Information Treatment with antibiotics is not of itself a reason for deferral but the reason for the treatment may be. When treatment is being given to prevent infection, rather than to treat it, see if there is a relevant entry. If not, discuss with a '**Designated Clinical Support Officer**'.

Reason for change The See if Relevant section has been revised.

Update Information This entry was last updated in:
DSG-WB Edition 203, Release 61.

Anticoagulant Therapy

Definitions An **anticoagulant** is a drug taken to limit the ability of blood to form a clot. Examples include heparin, warfarin and direct-acting oral anticoagulants (DOACs) such as apixaban, rivaroxaban and dabigatran.

Obligatory **Must not donate if:**
a) Receiving anticoagulant treatment or has been treated with anticoagulants in the last seven days.

b) The underlying reasons for anticoagulant treatment requires deferral

Discretionary a) If prescribed for treatment of venous thromboembolism:

- See Thrombosis and Thrombophilia

b) Otherwise, if treatment was completed more than seven days ago and:

- The reason for treatment does not preclude donation, and
- The donor is not under investigation,

accept.

See if Relevant Cardiovascular Disease
Clopidogrel
Drug Index - preparations which may affect platelet function
Nonsteroidal Anti-Inflammatory Drugs (including aspirin)

Superficial Thrombophlebitis
Thrombosis and Thrombophilia

<i>Additional Information</i>	There are many reasons that someone might be treated with an anticoagulant drug. It is important that the underlying indication for treatment is included in the assessment of the donor's eligibility to donate.
	While on anticoagulant treatment, it is more likely that a donor will bleed or bruise after donation. The effect of treatment wears off over some days. After seven days the blood clotting mechanisms should be back to normal.
	Donors taking antiplatelet medication such as aspirin or clopidogrel should be assessed using the relevant entries for their medication and the underlying reason for treatment.
<i>Information</i>	Part of this entry is a requirement of the Blood Safety and Quality Regulations 2005.
<i>Reason for change</i>	The scope of the entry has been clarified, with addition of a definition for anticoagulant medication. Additional links have been added to the See if Relevant section.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 68

Arrhythmias

<i>Obligatory</i>	<p>1. Must not donate if:</p> <p>a) Symptomatic or requires treatment</p> <p>b) The donor is undergoing investigation</p> <p>c) The donor has a history of an arrhythmia (eg Atrial Fibrillation, Atrial Flutter, Supraventricular Tachycardia, Ventricular Tachycardia) even if their symptoms have now settled.</p> <p>2. In other cases: Refer to a 'Designated Clinical Support Officer'.</p>
<i>Discretionary</i>	<p>1. Donors with a previous history of an arrhythmia triggered by a non-cardiac medical condition which has now been treated (eg thyrotoxicosis), refer to a DCSO.</p> <p>2. Donors who have been treated by ablation therapy for Supraventricular Tachycardia caused by either AVNRT (Atrioventricular Nodal Reentrant Tachycardia) or Wolff-Parkinson White Syndrome, and</p> <ul style="list-style-type: none"> • it is at least six months since successful ablation therapy, and • the donor does not require regular or 'as required' medication for their SVT, and • there is no other associated heart disease, and • the donor has been discharged from follow up, <p>accept.</p> <p>3. Donors with a history of palpitations where the donor has been assessed clinically and a cardiac cause has been excluded, see the entry for 'Palpitations'.</p>
<i>See if Relevant</i>	<u>Cardiovascular Disease</u> <u>Palpitations</u>
<i>Additional Information</i>	Some heart irregularities may be made worse by giving blood. This includes a risk that donation could trigger a recurrence in someone with a history of a previous arrhythmia. In cases where the donor's eligibility is not clear, DCSO referral ensures further information can be sought regarding their condition.
<i>Reason for change</i>	Discretionary criteria revised.
<i>Update Information</i>	

This entry was last updated in:
WB-DSG Edition 203 Release 72

Asthma

<i>Obligatory</i>	<p>Must not donate if:</p> <p>a) Asthma is symptomatic.</p> <p>b) Taking, or has completed, a course of oral or injected steroids lasting more than 3 weeks within the last six months.</p> <p>c) The donor has needed long term (six months or more) treatment with oral or injected steroids within the last 12 months.</p> <p>d) The donor has taken a short course (less than three weeks) of oral or injected steroids in the last seven days.</p> <p>e) The donor has been treated with monoclonal antibodies, or other biological modalities, in the last six months.</p>
<i>Discretionary</i>	If b), c) or d) above do not apply and the potential donor is asymptomatic at the time of donation, even if taking regular preventive treatment, including inhaled steroids, accept.
<i>See if Relevant</i>	<p><u>Infection - General</u> <u>Monoclonal antibody therapy or other Biological Modalities</u> <u>Steroid Therapy</u></p>
<i>Additional Information</i>	<p>Taking blood from a person with symptomatic asthma will lower the amount of oxygen the blood can carry and could make their symptoms worse.</p> <p>Steroid therapy can hide the signs and symptoms of infection. Blood from an infected donor can be dangerous to the person receiving it.</p>
<i>Reason for change</i>	Guidance has been added for donors treated with monoclonal antibodies and other biological modalities.
<i>Update Information</i>	This entry was last updated in: WB-DSG Edition 203 Release 71

Autoimmune Disease

<i>Obligatory</i>	<p>Must not donate if:</p> <p>a) The donor has needed treatment with steroids or conventional disease modifying antirheumatic drugs to suppress the condition in the last 12 months.</p> <p>b) The donor has needed treatment with monoclonal antibody therapy or other biologic disease modifying antirheumatic drugs to suppress the condition in the last 6 months.</p> <p>c) The cardiovascular system is involved.</p> <p>d) The donor has ongoing lung disease or renal impairment due to their condition.</p>
<i>Discretionary</i>	<p>a) If the donor:</p> <ul style="list-style-type: none"> • has been established on a stable maintenance treatment for an Autoimmune Disease with only one of the following drugs: Methotrexate, Sulfasalazine, Hydroxychloroquine or Azathioprine, and • the dose of the drug has not increased in the previous 6 months, and • the donor is well,

accept.

b) If there is any uncertainty about the diagnosis or the nature of treatment, refer to a DCSO.

See Is there a specific A-Z index entry for the condition you are assessing?

See if Relevant Cardiovascular Disease
Disabled Donor
Drug Index - Drugs and Platelet Donation
Fertility
Inflammatory Bowel Disease
Liver Disease
Monoclonal antibody therapy and other Biological Modalities
Nonsteroidal Anti-inflammatory Drugs
Skin Disease
Steroid Therapy
Thrombosis and Thrombophilia

If treated with transfusion, immunoglobulin, plasma exchange or filtration:

Transfusion

Additional Information Conventional systemic Disease Modifying Antirheumatic drugs (csDMARDs) are viewed as disease-modifying drugs. They include Methotrexate, Sulfasalazine, Hydroxychloroquine and Azathioprine. Sulfasalazine and Hydroxychloroquine have limited effect on the immune system. If used for maintenance treatment, Methotrexate and Azathioprine are usually given at lower doses which do not cause a significant degree of immunosuppression.

If the donor is taking higher dose Methotrexate or Azathioprine, they should not be accepted. If there is uncertainty about the dose refer to the DCSO for assessment. Further information on these drugs and immunosuppression can be found in 'The Green Book: Immunisation against Infectious Disease' (available at www.gov.uk).

Nonsteroidal anti-inflammatory drugs do not suppress the donor's immune system.

Physical therapies such as physiotherapy and hydrotherapy are not considered treatments to suppress the condition.

Autoimmune disease can cause problems such as infertility and thrombosis (antiphospholipid or Hughes' syndrome).

Some autoimmune conditions can permanently damage the cardiovascular system. If this is known to have happened, the person should not donate as they are more likely to have a serious adverse event.

Reason for change The obligatory criteria have been expanded to specify the deferral period required depending on the type of treatment.

Update Information This entry was last updated in:
WB-DSG Edition 203 Release 74

Back Problems

Obligatory **See:**
Is there an entry for the underlying condition?

Must not donate if:

Not able to use the bleed facilities provided without risking their own safety or the safety of others (donors must not be bled in a wheelchair).

See if Relevant Autoimmune Disease
Disabled Donor
Drug Index - preparations which may affect platelet function
Neurosurgery
Nonsteroidal Anti-Inflammatory Drugs
Surgery
Pain Killers

Additional Information Back problems have many causes. It is important to be certain that, for any individual, the cause is not a reason for them to be deferred.

It is also important that neither the donor, nor anyone assisting them, should risk injury by inappropriately attempting to use the bleed facilities provided. Alternative facilities may be available in other venues that may allow a donor with limited mobility to donate safely.

<i>Reason for change</i>	A link to painkillers has been added.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 04 Issue 02.

Bleeding Disorder

<i>Includes</i>	Coagulation factor deficiencies: <ul style="list-style-type: none"> • Factor I (one) fibrinogen deficiency (afibrinogenaemia, hypofibrinogenaemia) • Factor II (two) prothrombin deficiency • Factor V (five) deficiency • Factor VII (seven) deficiency • Factor VIII (eight) deficiency (Haemophilia A) • Factor IX (nine) deficiency (Haemophilia B, Christmas Disease) • Factor X (ten) deficiency • Factor XI (eleven) deficiency • Factor XIII (thirteen) deficiency • Von Willebrand disease (types 1, 2 and 3)
<i>Excludes</i>	Platelet disorders – see <u>Platelet Disorders</u> <p>Individuals who received Coagulation Factor Concentrates (including Prothrombin Complex Concentrates):</p> <ul style="list-style-type: none"> • To treat and prevent the coagulopathy associated with trauma and/or massive transfusion • To reverse the effect of anticoagulants such as warfarin (see Transfusion)

1. Affected Individuals and Carriers

<i>Obligatory</i>	<p>Must not donate if:</p> <p>a) Diagnosed with Haemophilia A (Factor VIII deficiency), Haemophilia B (Factor IX deficiency), Type 2 Von Willebrand Disease, Type 3 Von Willebrand Disease</p> <p>b) The donor has received a transfusion since 1st January 1980.</p> <p>c) The donor has ever: <ul style="list-style-type: none"> • received coagulation factor concentrates, including blood derived and recombinant products, and/or • received or is currently on treatment to reduce or prevent excessive bleeding e.g. desmopressin, tranexamic acid, oral contraceptive pill and similar hormone therapies. </p> <p>d) The donor has required or been advised they will require prophylactic treatment for surgery, dental treatment, or for any other procedure.</p> <p>e) There is a history of excessive bleeding or bruising.</p> <p>f) The donor is requiring monitoring and/or follow-up.</p> <p>g) There is associated organ involvement e.g. liver damage.</p> <p>h) For acquired disorders, the underlying cause or treatment precludes donation e.g. malignancy, monoclonal antibody therapy.</p>
<i>Discretionary</i>	If the donor has Type 1 Von Willebrand Disease and:

1. has not received a transfusion since 1st January 1980, and
2. has never received any type of coagulation factor treatment, and
3. has never received any other treatment to reduce or prevent excessive bleeding, and
4. has not received or been advised that they will require prophylactic treatment, and
5. has never had any excessive bleeding or bruising, and
6. is not requiring monitoring or follow-up, and
7. the underlying cause and/or treatment does not preclude donation,

accept.

If the donor is a carrier of a coagulation factor deficiency, and:

1. has not received a transfusion since 1st January 1980, and
2. has never received any type of coagulation factor treatment, and
3. has never received any other treatment to reduce or prevent excessive bleeding, and
4. has not received or been advised that they will require prophylactic treatment, and
5. has never had any excessive bleeding or bruising, and f) is not requiring monitoring or follow-up,

accept.

See if Relevant

Autoimmune Disease
Ehlers Danlos Syndrome
Malignancy
Monoclonal antibody therapy and other Biological modalities
Platelet Disorders
Transfusion

Additional Information

Coagulation factor deficiencies can be inherited or can be acquired, associated with haematological, neoplastic, cardiovascular, liver or autoimmune disease.

Some deficiencies cause significant bleeding, either spontaneously or in response to even minimal trauma or minor procedures. Individuals will have been assessed and advised about their condition and bleeding risk. They may have received treatment or been informed regarding the need for treatment in the future. The donor may have also been provided with a Bleeding Disorders Information Card.

Some people with the carrier state (trait) may be at risk of bleeding (symptomatic carriers). The diagnosis of the milder forms or carrier status of coagulation factor deficiencies may arise from family screening, or through testing during investigation for menorrhagia (heavy periods), or bleeding during pregnancy or childbirth.

If someone has had problems with bleeding or bruising, they may be at increased risk of complications from donation.

The guidance contained in this entry is not intended for use for donors without a coagulation factor deficiency, for example for someone who may have taken tranexamic acid for heavy periods due to an underlying gynaecological cause.

The current International Society on Thrombosis and Haemostasis (ISTH) classification recognises three types of Von Willebrand Disease: Type 1 is a partial quantitative deficiency of Von Willebrand Factor and is typically a milder form; the levels of von Willebrand Factor may overlap with the levels found in unaffected individuals.

More severe effects are usually seen with Von Willebrand Disease types 2 and 3. Care should be taken to determine the type of Von Willebrand Disease, as only donors with type 1 are potentially eligible.

Information

Part of this entry is a requirement of the Blood Safety and Quality Regulations 2005.

2. Family Members, Carers and Sexual Partners of Individuals Treated with Blood Derived Coagulation Factor Concentrates

<i>Obligatory</i>	<p>Must not donate if:</p> <p>a) Treated with blood derived coagulation factor concentrates.</p> <p>b) A sexual partner, or former sexual partner, of a person treated with blood derived coagulation factor concentrates.</p> <p>c) Less than four months after the date of an inoculation injury with either blood derived coagulation factor concentrates, or from blood contamination from an affected individual.</p> <p>d) Diagnosed as affected (even mildly) by the disorder.</p>
<i>Discretionary</i>	If three months or more from the last sexual contact, accept.
<i>See if Relevant</i>	<u>Non-Consented Exposure to Human Body Fluids Transfusion</u>
<i>Additional Information</i>	<p>Blood derived coagulation concentrates are made from the blood of many hundreds of individual donors. They may put recipients at risk of infections that can be passed through blood. This risk may be shared by their sexual partners and anyone suffering an inoculation injury.</p> <p>Many bleeding disorders are inherited. Family members that are blood relations may be affected by the bleeding disorder. They could be at risk of excessive bleeding or bruising. Most close blood relations would have been screened by a haematologist from whom additional information may be available.</p> <p>Waiting three or four months from the last sexual contact or inoculation injury helps to ensure that the infections tested for by the Blood and Tissues Services will be picked up.</p> <p>This guidance presumes that a validated NAT test for hepatitis C is negative. If this test is stopped, the guidance will change.</p>
<i>Information</i>	Part of this entry is a requirement of the Blood Safety and Quality Regulations 2005.
<i>Reason for change</i>	<p>Expansion of the Includes section and addition of an Excludes section to clarify the scope of the guidance contained in this entry.</p> <p>Expansion of obligatory and discretionary criteria applicable to affected individuals and carriers.</p>
<i>Update Information</i>	This entry was last updated in: WB-DSG Edition 203 Release 72

Blood Pressure - High

<i>Obligatory</i>	<p>Must not donate if:</p> <p>a) The cause of hypertension is under investigation.</p> <p>b) Anti-hypertensive medication has been altered in the last four weeks.</p> <p>c) Is having problems with feeling faint, fainting or giddiness.</p> <p>d) Has suffered from heart failure.</p> <p>e) Has renal impairment requiring dialysis, the use of erythropoietin or similar drugs, or is either under active investigation or continued follow up for renal impairment.</p> <p>f) Has required surgery for a blocked or narrowed artery including any type of amputation.</p> <p>g) Has or has had gangrene.</p>
-------------------	---

<i>Discretionary</i>	<p>a) If the donor is being regularly assessed for high blood pressure but treatment has not been commenced, accept.</p> <p>b) If the donor is taking medication for raised blood pressure and neither the type nor the dose has been changed in the last four weeks and they are otherwise well, accept.</p> <p>c) If gangrene was not related to diabetes or peripheral vascular disease (e.g. it was due to hypothermia or meningococcal meningitis) and all wounds are fully healed, even if amputation was required, accept.</p>
<i>See if Relevant</i>	<p><u>Cardiovascular Disease</u> <u>Cerebrovascular Disease and Intracranial Haemorrhage</u> <u>Diabetes Mellitus</u> <u>Kidney and Bladder Disease</u></p>
<i>Additional Information</i>	<p>The rationale for not accepting donors on medication, other than beta blockers or diuretics, for the treatment of hypertension was reviewed by the Standing Advisory Committee for the Care and Selection of Donors in 2008. It was decided that available data did not support the deferral of all individuals with controlled hypertension taking other medications.</p> <p>In the UK about one in twenty individuals has hypertension. Most people with hypertension are in good health and are fit to donate blood.</p> <p>It is however important that complications due to raised blood pressure are carefully assessed and, where necessary, donors are excluded from donating (e.g. those with heart failure or damage to their kidneys, or those experiencing hypotensive side effects from their medication).</p>
<i>Reason for change</i>	Link updated from 'Central Nervous System Disease' to 'Cerebrovascular Disease and Intracranial Haemorrhage' in the 'See if Relevant' section.
<i>Update Information</i>	This entry was last updated in: WB-DSG Edition 203 Release 73

Blood Pressure - Low

<i>Also Known As</i>	Hypotension.
<i>Discretionary</i>	If the donor is in good health and does not have faints or dizzy spells, accept.
<i>See if Relevant</i>	<u>Faints</u>
<i>Additional Information</i>	<p>Low blood pressure is not normally a problem. It is common in women and seems to be linked with the female sex hormone oestrogen.</p> <p>Low blood pressure can be caused by serious heart disease. In such cases a donation would not be taken.</p> <p>Fainting can put a donor at risk of injury. Any donor who has problems with faints or dizzy spells should not donate.</p>
<i>Reason for change</i>	A link has been added to 'Faints'.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 01.

Blood Safety Entry

<i>Definitions</i>	<p>Individual risk is based on the donor's sexual behaviour, including new partners and number of partners.</p> <p>Partner risk is based on sexual contact with a partner who may, at a population level, be at higher risk of acquiring infection, as described in this entry.</p>
--------------------	---

Sexual contact is defined as oral, vaginal or anal sex.

Anal sex is defined as penile-anal intercourse only. It does not apply to oro-anal sex or the use of sex toys.

Chemsex is sex while using stimulant drugs taken for the specific purpose of enhancing sexual experience and reducing inhibitions. Chemsex does not refer to sex after using alcohol or recreational drugs for other purposes, nor the use of drugs such as Viagra or Cialis to treat erectile dysfunction.

Obligatory Information must be provided so that individuals at risk do not donate. The reasons for donor self-exclusion must be understood.

1. You must not donate if:

You think you need a test for HIV/AIDS, HTLV or hepatitis.

2. You must never donate if:

- a) You are HIV positive or receiving treatment for HIV.
- b) You are HTLV positive.
- c) You are a hepatitis B carrier.
- d) You are a hepatitis C carrier.
- e) You have ever been diagnosed with syphilis, even if treated
- f) You have ever injected or been injected with drugs; even a long time ago or only once. This includes bodybuilding drugs, injected tanning agents and injected chemsex drugs. You may be able to give if a doctor prescribed the drugs. Please ask.

3. You must not donate for at least three months if:

You are working as a sex worker. You may be accepted for donation if it is longer than three months since you last received money or drugs for anal, vaginal or oral sex.

4. Individual risk criteria

- a) You must not donate for at least three months if you have taken part in chemsex activity, including the use of stimulant drugs. This risk applies for all sexual contact
- b) You must not donate if you have been diagnosed with gonorrhoea, until at least three months after completion of treatment and discharge from further follow up.
- c) You must not donate if in the last three months,
 - you have had more than one sexual partner, AND
 - you have had anal sex with any of your partners.
- d) You must not donate if in the last three months, you have had anal sex with a new sexual partner. For the purpose of donor selection, a new partner is someone that you have not had sex with before or a previous partner with whom you have restarted a sexual relationship.

If you are in a sexual relationship with one partner only, you can donate once it is three months from the date of first sexual contact, even if you are having anal sex.

5. Partner risk criteria

You must not donate for at least three months after sexual contact with a partner who is, or you think may be:

- a) HIV or HTLV positive.
- b) A hepatitis B carrier.
- c) A hepatitis C carrier.
- d) A partner who has ever received money or drugs for sex.
- e) A partner who has ever injected, or been injected with, drugs: even a long time ago or only once. This includes bodybuilding drugs, injected tanning agents and injected chemsex drugs. You may be able to give if a doctor prescribed the drugs. Please ask.

6. You must not donate for at least three months if:

- a) You have taken Pre-Exposure Prophylaxis (PrEP) / Truvada[®] for prevention of HIV
- b) You have taken or been prescribed Post-Exposure Prophylaxis (PEP) for prevention of

HIV.

If the underlying reason for taking PrEP or PEP warrants a longer deferral period, this should be applied.

See if Relevant

Addiction and Drug Abuse

Hepatitis B

Hepatitis C

HIV

HTLV

Infection - General

Non-consented Exposure to Human Bodily Fluids

Pre- or Post-Exposure Prophylaxis for HIV

Sexually Transmitted Disease

Syphilis

Additional Information

The FAIR (For the Assessment of Individualised Risk) study considered changes to the donor selection policy to allow a more individualised risk-based approach to donor selection policy. In their 2020 report, the FAIR group specifically looked at the guidelines which applied to men who have sex with men (MSM) and recommended an approach based on assessment of a donor's recent sexual behaviour and experience. This approach has been agreed by SaBTO and has now been implemented by the UK Transfusion Services.

Changes to donor selection criteria mean that donors who were previously excluded because of their sexuality and/or gender can now be accepted, if they meet the individual risk criteria outlined in FAIR (see section 5, above). These rules must be applied equally to all donors.

FAIR identified several factors associated with a higher risk of blood borne infections. These include the recent diagnosis of a bacterial sexually transmitted disease and the following sexual behaviours:

- new or multiple sexual partners
- anal sex
- participation in chemsex activity

Drugs used for chemsex include methamphetamine, mephedrone and GHB/GBL, but other drugs may be used (e.g. ketamine, poppers, cocaine). Chemsex is a high risk activity because it usually involves multiple sexual partners, sometimes for extended periods of time. The drugs involved also reduce inhibition leading to riskier sexual activity.

Infection with some sexually transmitted diseases, particularly gonorrhoea or syphilis, is associated with a higher risk of acquiring blood borne infections. Donors who have had gonorrhoea can give three months after completing treatment. Donors who have had syphilis are permanently deferred as their blood will still react in screening tests, even if they have been successfully treated

Some partner risk criteria still apply to donors who have a partner in a population group at higher risk of infection. For affected donors, these risk criteria should be applied even if the donor is eligible under individual risk criteria. These risks are being reviewed and will be updated as further evidence is available.

The drugs used in both Pre- and Post-Exposure Prophylaxis for HIV (PrEP and PEP) may interfere with the routine HIV screening tests carried out on all blood donations. For this reason, donors who have taken PrEP or PEP should not donate for three months, even if they are otherwise eligible under individual risk criteria.

Information

This is a requirement of the Blood Safety and Quality Regulations 2005.

Reason for change

Removal of reference to the Blood Safety Leaflet Information.

Update Information

This entry was last updated in:
DSG-WB Edition 203, Release 69

Body Piercing

<i>Includes</i>	Derma-rolling, ear and body piercing, permanent and semi-permanent make-up, tattooing (including memorial tattoos) platelet rich plasma (PRP) facial, and ritual self-flagellation.
<i>Obligatory</i>	Must not donate if: Less than four months from last piercing.
<i>Discretionary</i>	Painting, stencilling or transfers applied to the skin without piercing, accept.
<i>Additional Information</i>	<p>Piercing has passed infection from person to person. Waiting four months helps to ensure that the infections tested for by the Blood and Tissues Services will be picked up.</p> <p>Platelet rich plasma (PRP) facials (also known as 'Vampire Facials') have been associated with HIV transmission.</p> <p>Ritual self-flagellation is carried out by some religious groups. The practice includes beating or flogging oneself with sharp objects. It may be associated with exposure to blood from other participants, either directly or through contamination of shared equipment.</p> <p>This guidance presumes that a validated NAT test for hepatitis C is negative. If this test is stopped, the guidance will change.</p>
<i>Information</i>	Part of this entry is a requirement of the Blood Safety and Quality Regulations 2005.
<i>Reason for change</i>	A deferral for ritual self-flagellation has been added.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 63

Breastfeeding

<i>See</i>	<u>Pregnancy</u>
<i>Additional Information</i>	Breastfeeding is not of itself a reason to defer but the time from giving birth may be.
<i>Reason for change</i>	'Additional Information' has been added.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 01.

Cardiac Surgery

<i>Obligatory</i>	Must not donate.
<i>Discretionary</i>	If surgery was for a congenital heart defect, cure has been achieved and donation is not excluded because of their transfusion history, accept.
<i>See if Relevant</i>	<u>Cardiovascular Disease</u> <u>Surgery</u> <u>Transfusion</u>
<i>Additional Information</i>	Individuals who have had cardiac surgery, other than for congenital abnormality, are unlikely to be fit enough to safely have a unit of blood removed. An individual who has had congenital abnormalities corrected can often lead a normal lifestyle and may be able to give blood safely.
<i>Reason for change</i>	To bring guidance into line with the Cardiovascular Disease Topic
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 36.

Cardiovascular Disease

- Obligatory* **Must not donate if has or has had:**
- a) An aneurysm.
 - b) Cardiomyopathy.
 - c) Ischaemic heart disease or angina regardless of cause including INOCA (myocardial ischaemia with non-obstructive coronary arteries).
 - d) Heart failure.
 - e) Myocarditis and it is less than 12 months from recovery.
 - f) Peripheral vascular disease (including intermittent claudication and gangrene).
 - g) Has required surgery for a blocked or narrowed artery including any type of amputation.
 - h) Valvular heart disease.
 - i) Heart Block or Bundle Branch Block.

- Discretionary*
- a) If an incidental heart murmur has been heard or a valve abnormality has been found at echo, which is asymptomatic and does not require follow up, accept
 - b) If asymptomatic and there is no treatment planned for Patent Foramen Ovale (PFO), accept
 - c) If a congenital heart defect has been treated medically or surgically, cure has been achieved (or the defect has spontaneously resolved) and donation is not excluded because of a transfusion history, accept.
 - d) If the donor has been diagnosed with Right Bundle Branch Block (RBBB), and the donor has been clinically assessed and found to have no evidence of cardiac or pulmonary disease, accept.
 - e) If the donor has been diagnosed with First Degree Heart Block, and the donor has been clinically assessed and found to have no evidence of cardiac disease, accept for whole blood donation.
 - f) If the donor has been found to have coronary atheroma as an incidental finding during routine investigations and
 - if the donor has not been advised to take antiplatelet agents (e.g. aspirin) and/or cholesterol lowering medication, and
 - there is no history of chest pain or other cardiac symptoms,
 accept.

See if Relevant

Arrhythmias
Blood Pressure - High
Cardiac Surgery
Central Nervous System Disease
Cerebrovascular Disease and Intracranial Haemorrhage
Endocarditis
Shunts, Stents and Devices
Superficial Thrombophlebitis
Thrombosis and Thrombophilia

Additional Information A history of 'Cardiovascular Disease' means that removing blood from their circulation may put the donor at risk of having a heart attack, stroke or other vascular incident.

Patent Foramen Ovale (PFO) is a normal variant found in up to 40% of the population at post mortem. If it is asymptomatic and no treatment or surgery is planned for this atrial septal defect, donors can be accepted.

Incidental heart murmurs and valve abnormalities are increasingly being found due to the sensitivity of new testing regimes. If the abnormality is of no clinical significance, i.e. the

donor is asymptomatic and does not require treatment or follow up, the donor may be accepted.

RBBB and first degree heart block can be diagnosed in individuals in the absence of heart disease. Provided the donor has been clinically assessed and there is no evidence of cardiovascular or pulmonary disease, the donor can be accepted. If there is any uncertainty about the diagnosis or the results of investigations, refer to a DCSO.

Reason for change Guidance on berry aneurysms has been moved to the Cerebrovascular Disease and Intracranial Haemorrhage entry.

Update Information This entry was last updated in:
WB-DSG Edition 203 Release 73

Catarrh

1. Acute

See Infection - Acute

Additional Information Catarrh may be due to infection or to allergy. If the problem is new, it should be treated as an infection.

Update Information This entry was last updated in:
DSG-WB Edition 203, Release 01.

Reason for change Additional Information' has been added.

2. Chronic

Obligatory **Must not donate if:**
Taking prescribed medication for catarrh other than antihistamines, a nasal decongestant or nasal steroids.

Discretionary If using antihistamines, a nasal decongestant or nasal steroids only, accept.

See if Relevant Allergy
Infection - General
Steroid Therapy

Additional Information Chronic catarrh may be due to infection or to allergy. A decision will need to be made as to the underlying cause.

Update Information This entry was last updated in:
DSG-WB Edition 203, Release 01.

Reason for change A reference to antihistamines have been added to 'Obligatory' and 'Discretionary'.

A link has been added to 'Allergy' and 'Steroid Therapy'. 'Additional Information' has been added.

Update Information This entry was last updated in:
DSG-WB Edition 203, Release 01.

Update Information This entry was last updated in:
DSG-WB Edition 203, Release 01.

Central Nervous System Disease

Excludes Cerebrovascular disease and all forms of intracranial haemorrhage.

Obligatory **Must not donate if has or has had:**
a) Dementia (e.g. Alzheimer's disease).

b) A history of CNS disease of unknown aetiology or suspected infective origin. These include, but are not limited to, neurodegenerative conditions, multiple sclerosis (MS), optic neuritis, clinically isolated syndrome, and transverse myelitis.

c) Malignant tumour.

d) Parkinson's Disease

e) If having symptoms related to hypotension while taking dopamine receptor agonist drugs such as rotigotine, ropinirole and pramipexole.

Discretionary

a) Individuals who have had Bell's palsy more than four weeks ago and have discontinued any treatment for the condition for at least seven days, once investigated and discharged from specialist follow-up even if they have residual paralysis, accept.

b) If the donor has been investigated and a definite diagnosis of transient global amnesia has been made, accept.

c) If diagnosed with Idiopathic (benign) intracranial hypertension (IIH) and

- the donor is asymptomatic, and
- if the donor is taking diuretics (e.g. Acetazolamide) for IIH and the dose has not changed in the last four weeks,

accept.

d) If taken for a condition other than Parkinson's Disease, as long as not having symptoms of hypotension related to dopamine receptor agonist drugs such as rotigotine, bromocriptine, ropinirole and pramipexole, accept.

See if Relevant

Cardiovascular Disease
Cerebrovascular Disease and Intracranial Haemorrhage
Epilepsy
Infection - General
Neurosurgery
Pituitary Disorders
Prion Associated Diseases
Steroid Therapy
Urinary Catheterisation

Additional Information

Donor safety:

Transient global amnesia is a temporary and isolated disorder of memory. Affected individuals are usually over 50 years of age and there is an association with migraine. There is no association with cerebrovascular disease.

Idiopathic or benign intracranial hypertension is a raised intracranial pressure where no mass or other disease is present.

Parkinson's disease is a movement disorder that can make venepuncture difficult and lead to damage to the donor's arm. It is also treated with dopamine receptor agonist drugs (e.g. rotigotine, bromocriptine, ropinirole and pramipexole) that can cause hypotension and fainting.

Recipient safety:

It is thought that degenerative brain disease in the form of vCJD has been transmitted by blood transfusion. Often the exact cause of a degenerative brain condition only becomes known after death. For this reason, when there is any doubt as to the underlying cause of a brain condition, it is considered safest not to accept a donation.

Information

This is a requirement of the Blood Safety and Quality Regulations 2005.

Reason for change

A discretion to accept donors on treatment for IIH has been added.

Update Information

This entry was last updated in:
 WB-DSG Edition 203 Release 73

Cerebrovascular Disease and Intracranial Haemorrhage

Includes Diseases of the vasculature of the brain. This includes:

- Stroke or Cerebrovascular accident (haemorrhagic or embolic)
- Transient Ischaemic Attack.
- Vascular Dementia
- Carotid Artery Disease
- Cerebral haemorrhages and haematomas that are intracranial, subdural, subarachnoid, or epidural

Obligatory **Must not donate.**

Discretionary If the donor has had one or more berry aneurysms treated by interventional radiology or surgery and

- the donor has recovered from any associated subarachnoid haemorrhage, and
- there is no residual neurological deficit, such as a stroke, and
- any surgery did not require the use of dural grafts and/or was performed in the UK after 1992,

refer to a DCSO.

If the donor has recovered from a single episode of intracranial haemorrhage due to trauma, and

- there is no underlying cerebral or cerebrovascular disease, and
- there is no underlying bleeding disorder, and
- there is no residual neurological deficit, such as a stroke,

accept.

See if Relevant Central Nervous System Disease
Epilepsy
Neurosurgery
Transfusion

Additional Information A history of thrombotic stroke or cerebral haemorrhage may increase the risk of donor adverse events. In order to reduce this risk, donors with a history of cerebrovascular disease must be deferred.

If the incident was due to trauma, and not intrinsic cerebrovascular pathology, and donor has fully recovered, it may be appropriate to accept the donor.

Reason for change This is a new entry.

Update Information This entry was last updated in:
WB-DSG Edition 203 Release 73

Cervical Dysplasia

Obligatory **Must not donate if:**
a) Undergoing investigation or treatment.

b) Diagnosed with invasive cervical carcinoma.

Discretionary a) If the donor had colposcopy treatment for abnormal cervical cells and has been discharged to routine screening, accept. It is not necessary to wait for a normal smear result before donating.

b) If only having regular review of smears, accept.

Additional Information Cervical screening includes testing for high risk Human Papilloma Virus (HR-HPV). Women who are positive for HR-HPV may be called for routine smear tests at more frequent intervals. They can donate provided they are not undergoing other tests or awaiting colposcopy investigation.

Women with abnormal cells on a smear test are triaged according to their risk of developing cervical carcinoma. Women at higher risk will be referred for investigation and treatment via colposcopy.

A colposcope is an instrument used to view the neck of the womb (cervix). It is not a flexible endoscope so its use is not a reason for deferral if the donor is otherwise eligible to donate.

Abnormalities identified at colposcopy include cervical intra epithelial neoplasia (CIN, Grades 1-3) and cervical glandular intra epithelial neoplasia (CGIN). CIN-3 is also known as cervical carcinoma in situ. By definition, patients with CIN or CGIN do not have invasive cervical carcinoma, so can be accepted once treated, fully healed and discharged. There is no need to wait for the results of their next routine smear, usually at 6 months post treatment, unless the donor has been advised that follow up will be necessary at the colposcopy clinic.

<i>Information</i>	This entry is compliant with the Blood Safety and Quality Regulations 2005.
<i>Reason for change</i>	Updated to clarify the scope of entry, when donor can return after treatment for cervical dysplasia and the significance of HR-HPV testing.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 55.

Chest Pain

<i>Obligatory</i>	Must not donate if: a) Due to heart disease. b) The cause is not known.
<i>Discretionary</i>	If donor has been investigated for chest pain and causes that would otherwise result in deferral have been excluded such as ischaemic heart disease, pulmonary embolism or infection, accept.
<i>See if Relevant</i>	<u>Autoimmune Disease</u> <u>Cardiovascular Disease</u>
<i>Additional Information</i>	It is important not to take a donation from an individual with ischaemic heart disease as any lowering of blood pressure could result in a heart attack. If the cause of any chest pain has not been investigated it could potentially be due to heart disease and a donation should not be taken.
<i>Reason for change</i>	To clarify the discretionary acceptance criteria.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 18.

Chiropody

<i>Also Known As</i>	Podiatry.
<i>Obligatory</i>	Must not donate if: There are open wounds or infection.
<i>See if Relevant</i>	<u>Infection - General</u> For fungal infection see: <u>Skin Disease</u>
<i>Additional Information</i>	An unhealed wound or sore is a risk for bacteria entering the blood. Bacteria in blood can be a serious threat to anybody receiving blood, products made from it, or tissues. This is because the bacteria can multiply to dangerous levels during storage.
<i>Reason for change</i>	'Additional Information' has been added together with a link to 'Infection - General' and 'Skin Disease'.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 01.

Chondromalacia

<i>Discretionary</i>	Accept.
<i>See if Relevant</i>	<u>Disabled Donor</u> <u>Drug Index - preparations which may affect platelet function</u> <u>Nonsteroidal Anti-Inflammatory Drugs</u> <u>Surgery</u>
<i>Additional Information</i>	This is caused by abnormal softening or degeneration of the cartilage of joints. It especially affects the knee in adolescents and is thought to be related to rapid growth. The condition itself is not a reason to defer but treatment or disability caused by the condition may be relevant to donation.
<i>Reason for change</i>	'Additional Information' and links have been added.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 01.

Chronic Fatigue Syndrome

<i>Also Known As</i>	CFS, myalgic encephalomyelitis (ME), post-viral fatigue syndrome and Systemic Exertion Intolerance Disease (SEID).
<i>Obligatory</i>	Must not donate
<i>Discretionary</i>	If donor gives a history of fatigue following a viral infection, e.g. Glandular fever, with no relapse of symptoms and it is at least six months since all symptoms resolved, accept.
<i>Additional Information</i>	CFS is generally diagnosed by excluding other conditions and may follow an infection that may or may not have been viral and which may be carried by the affected individual. It is most common between the ages of 25 and 45 years and women are affected more often than men. It is associated with easily induced and prolonged episodes of fatigue often accompanied by other symptoms. The condition is relapsing by nature and donation may make symptoms worse, or provoke a relapse in an affected individual. Post viral fatigue can occur after an acute viral infection. Symptoms of fatigue can last weeks or months and may follow a relapsing course. It is important that individuals have fully recovered before being accepted to donate.
<i>Reason for change</i>	A discretion to accept donors who have fully recovered from fatigue associated with an acute viral infection has been added.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 69

Clinical Trials

<i>Obligatory</i>	Must not donate if: Participating in a clinical trial. This includes the use of drugs of any kind (oral, injected, transcutaneous, etc.) and applies to healthy individuals participating as volunteers - for example in 'phase 1' clinical trials.
<i>Discretionary</i>	a) If a ' Designated Clinical Support Officer ' has examined and agreed the trial protocol, accept. b) If the trial does not involve the use of drugs (e.g. hypnotherapy, physiotherapy) and any underlying condition would not be a reason to defer, accept.
<i>See if Relevant</i>	

Complementary Therapy
Monoclonal antibody therapy and other Biological Modalities
Transfusion

<i>Additional Information</i>	It is important for the Blood Services to know that anything being given to a donor as part of a clinical trial will not affect either the safety of the donor or of any potential recipient.
	When a particular drug treatment is being assessed, trial participants may be randomly allocated to receive the treatment or a placebo drug. Participants should know which treatment is under investigation in their trial (or trial arm) but will not know whether they have had the treatment or not. They should be assessed for donation on the basis that they might have done.
	Some donors may not recall which treatment was under investigation in their trial (or trial arm). In this case, the donor should be asked to find out and contact us again when they have the information available.
<i>Reason for change</i>	Removal of specific details regarding COVID-19 trials. The See if Relevant section has been revised.
<i>Update Information</i>	This entry was last updated in: WB-DSG Edition 203 Release 71

Clopidogrel

Obligatory **Must not donate**

<i>Additional Information</i>	Clopidogrel is an antiplatelet drug which is used in the treatment and secondary prevention of cardiovascular disease and stroke. Occasionally Clopidogrel is used for primary prevention in patients who are intolerant of or hypersensitive to aspirin. The risk of bruising after blood donation while on Clopidogrel is not known, so any donor on this medication should be deferred, even if they are otherwise eligible.
<i>Reason for change</i>	This is a new entry.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 55.

Coeliac Disease

Discretionary Accept.

<i>Additional Information</i>	Coeliac disease is an abnormal immune response to gluten (contained in some cereals, in particular wheat) that damages the small bowel. This can lead to poor absorption of minerals and vitamins that are necessary to make blood. Avoiding gluten reverses the problem. The haemoglobin screening test will check that an individual is not significantly anaemic before a donation is taken.
<i>Reason for change</i>	'Additional Information' has been added.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 01.

Communication Difficulties

Obligatory

1. All donors must:

- a) Fully understand the donation process.
- b) Give their informed consent to the process and to the testing of their blood for diseases that may affect its suitability for use.

2. Third party interpreters:

If they are to be present at any part of the selection procedure where there is an exchange of confidential information between the donor and the qualified health professional, they must:

- a) Understand the requirements of the Blood Safety and Quality Regulations (BSQR) relevant to the donation process and provide an accurate and truthful translation and interpretation of all information provided to enable the Blood Service to comply with these regulations.
- b) Not be personally known to the donor.
- c) Fully understand their duty of confidentiality and the confidential nature of any information obtained from the donor.

See if Relevant

Central Nervous System Disease
Disabled Donor
Neurobehavioral Disorders

Additional Information

The Blood and Tissue Services are aware of their duties under Race Relations and Disability Discrimination Legislation and will, whenever and wherever reasonable, try to provide facilities for individuals whose first language is not English, or who have other difficulties in communicating. Potential donors with such difficulties are advised to seek advice from their local [Blood Service Help Line](#) before attending a donor session to see if their needs can be met. It is however important to note the following.

To comply with Part 2 of the Blood Safety and Quality Regulations 2005 (BSQR) every donor must:

- a) Be provided with accurate educational materials, which are written in terms which can be understood by members of the general public (Part A 1-13).
- b) Complete a health and medical history questionnaire and undergo a personal interview performed by a health professional (defined in the BSQR as a doctor, a nurse or a donor carer) trained and qualified in the requirements of the BSQR (Part B 15).
- c) Provide written informed consent to proceed with the donation process which must be countersigned by the qualified health professional responsible for obtaining the health history (Part B 16 (a) - (f)).

A qualified health professional may assist a donor in the completion of the health and medical history questionnaire and in understanding the consent statement and any other information provided by the Blood Service. To facilitate comprehension it is permissible to use alternative formats (e.g. a language other than English, audio, computer, Braille) for the donor information leaflets, the health and medical history questionnaire and consent statements. The donor must be able to clearly demonstrate they have understood this material. At present there is no standardized way of assessing comprehension so this will be a personal judgement made by the qualified health professional.

Use of third party interpreters.

It is permissible for any third party to act as an enabler by helping to reassure the donor and to assist in establishing effective communication between the donor and the qualified health professional. The third party **must not** however be present during any exchange of confidential information, unless they are **not** personally known to the donor, understand the requirements of that part of the BSQR relevant to the donation process and provide an accurate and truthful interpretation of all information, including personal and confidential information, provided to enable the Blood Service to comply with these regulations. Confidential parts of the process include the evaluation of the health and medical history questionnaire, the medical interview and the obtaining of valid consent. Any third party, with the permission of the donor, may accompany the donor through other parts of the donation process that do not include the exchange of confidential information.

Rationale.

There is concern that the use of third parties during any exchange of confidential information between the donor and the qualified health professional may compromise the confidentiality of the donor and the safety of the blood supply. Interpreters are often part of a close community, or a family member, and this may inhibit or embarrass the potential donor in any confidential exchange of information. This may result in the non-disclosure of sensitive information that could affect the individual's eligibility to donate. If a third party is not fully

aware of the relevant aspects of the BSQR and the need to provide an accurate and truthful interpretation of all information, including personal and confidential information, provided by the donor this may make the interpretation of information incomplete and potentially put both the donor and the blood supply at risk. There is also a requirement to communicate the results of any testing performed by the Blood Services that may be of relevance to the donor's health in a way that protects their confidentiality. The continuing availability of an independent interpreter, to maintain donor confidentiality, should be taken into account when deciding if an individual donor may be accepted.

To comply with both the BSQR and Health and Safety Regulations no donor can be accepted if it unnecessarily puts their own safety or the safety of others at risk.

<i>Information</i>	Part of this entry is a requirement of the Blood Safety and Quality Regulations 2005.
<i>Reason for change</i>	This has been updated to clarify the role of a third party interpreter and to bring it in line with the Tissue donor Guidelines.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 29.

Complementary Therapy

<i>Obligatory</i>	<p>Must not donate if:</p> <p>a) The condition for which treatment was given is not acceptable.</p> <p>b) It is less than four months from any treatment that involves:</p> <ul style="list-style-type: none"> • piercing the skin (e.g. acupuncture) • drawing blood (e.g. wet cupping) • an invasive procedure (e.g. colonic irrigation)
<i>Discretionary</i>	<p>a) If oral or topical complementary medicines only and reason for which treatment was given is acceptable, accept</p> <p>b) For all other therapies</p> <p>1. Performed within the NHS If performed by NHS staff on NHS premises including GP surgeries, accept.</p> <p>2. Performed outside of the NHS If performed by a Qualified Health Care Professional registered with the</p> <p>General Medical Council (GMC), Nursing and Midwifery Council (NMC), General Dental Council (GDC), The General Chiropractic Council (GCC), The General Optical Council (GOC), The General Osteopathic Council (GOsC), General Pharmaceutical Council (GPhC), Pharmaceutical Society of Northern Ireland (PSNI), The Health and Care Professions Council (HCPC) (which regulates Physiotherapists, Arts therapists, Biomedical Scientists, Chiropodists/ Podiatrists, Clinical Scientists, Dieticians, Hearing Aid Dispensers, Occupational Therapists, Operating Department Practitioners, Orthoptists, Paramedics, Practitioner Psychologists, Prosthetists and Orthotists, Radiographers and Speech and Language Therapists), accept.</p>
<i>Additional Information</i>	<p>Equipment that has been reused has passed infection from person to person. Therapists who are subject to discipline from statutorily constituted professional authorities are expected to follow safe practices.</p> <p>This guidance presumes that a validated NAT test for hepatitis C is negative. If this test is stopped the guidance will change.</p> <p>When there is any doubt about infection being passed on, waiting four months means infections are more likely to be picked up by the tests used by the blood services.</p>
<i>Information</i>	Part of this entry is a requirement of the Blood Safety and Quality Regulations 2005.
<i>Reason for change</i>	

The regulatory organisations for Pharmacists in the UK have been added.
The HCPC ceased to be the regulatory authority for Social Workers in England in 2019. The list of health and care professionals regulated by the HCPC has been amended.

Update Information This entry was last updated in:
DSG-WB Edition 203, Release 61.

Conn's Syndrome

<i>Excludes</i>	Secondary Hyperaldosteronism
<i>Obligatory</i>	Must not donate
<i>Discretionary</i>	If donor has undergone surgery for a benign adrenal tumour (adenoma), the donor is fully recovered, has been discharged from follow up and is not on medication for Conn's syndrome, accept.
<i>See if Relevant</i>	<u>Blood Pressure – high</u> <u>Diuretics</u>
<i>Additional Information</i>	<p>Conn's syndrome (primary hyperaldosteronism) is caused by overproduction of aldosterone in the adrenal cortex. Aldosterone is a hormone that regulates potassium and sodium levels as well as fluid balance in the body thereby maintaining blood volume and blood pressure, helping the body to control stress and maintain a steady metabolic state and normal electrolyte balance.</p> <p>Secondary hyperaldosteronism is caused by increased adrenal production by medical conditions outside the adrenal gland which stimulates the renin- angiotensin- aldosterone mechanism.</p>

Contraceptive Use

<i>Discretionary</i>	Accept.
<i>See if Relevant</i>	<u>Anaesthetic</u> <u>Pregnancy</u> <u>Surgery</u>
<i>Additional Information</i>	<p>If a female donor has been pregnant in the last nine months, see <u>Pregnancy</u>.</p> <p>The use of contraceptives should not normally be a reason to defer a donor. However if surgery (leaving a wound), a local anaesthetic or sedation was required to introduce a contraceptive within the last seven days, please see the entry on 'Surgery' or 'Anaesthesia' as appropriate.</p>
<i>Reason for change</i>	Advice in additional information has been brought in line with that for Surgery and Anaesthesia.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 04 Issue 02.

Coronavirus Infection (COVID-19)

<i>Includes</i>	COVID-19 disease due to infection with SARS-CoV-2 virus, previously known as Novel Coronavirus or 2019-nCoV.
<i>Definitions</i>	Testing: PCR (polymerase chain reaction) and rapid lateral flow tests (LFTs), usually by throat and/or nose swab, to detect the presence of SARS-CoV-2. This does not include testing for antibodies to SARS-CoV-2

1. Individuals with confirmed or suspected COVID-19 infection

Includes

- Individuals with confirmed COVID-19 infection, diagnosed by a positive LFT or PCR test.
- Individuals where the results of SARS-Cov-2 testing, if carried out at the request of a health care professional, are awaited.

Obligatory **Must not donate.**

Discretionary If it is at least seven days from the resolution of symptoms, and no further testing is required, accept.

2. Individuals with non-specific symptoms, not confirmed as COVID-19

Includes Individuals who have non-specific symptoms of a respiratory infection, including coughs and cold symptoms.

Excludes Individuals who are awaiting test results for SARS-CoV-2 infection, as requested by a health care professional.

Obligatory See Infection - Acute

3. Post-Covid Syndrome (Long Covid)

Obligatory **Must not donate**

Discretionary If it is at least 6 months since all symptoms, including fatigue, have resolved, accept.

4. Occupational and other routine surveillance

Discretionary Donors who have regular testing for the presence of SARS-CoV-2 (Coronavirus) can be accepted to donate provided they have not had a positive test for SARS-CoV-2 in the last seven days.

This includes donors who work in a Health, Social Care or Educational setting.

See if Relevant Clinical Trials
Immunisation – Non-live
Infection - Acute

Additional Information Common coronaviruses cause colds and respiratory tract infections but are not considered a risk for transfusion recipients. Since 2002 there have been outbreaks in humans of new strains of coronavirus, associated with severe pulmonary infections and mortality rates of 10-35% e.g. SARS and MERS.

COVID-19 is an illness caused by infection with SARS-CoV-2, a new coronavirus first identified in 2019. The guidance within this entry is focussed on COVID-19. Donors who report MERS or SARS, or contact with these infections, should be referred to a DCSO.

Many respiratory illnesses, including COVID-19, share common symptoms. As routine testing for SARS-CoV-2 infection is no longer recommended, most individuals will not have test results to confirm or exclude COVID-19. Where COVID-19 testing is not being undertaken, assessing donors using the Infection – Acute entry ensures that individuals are deferred for the appropriate time based on their symptoms.

Individuals affected by COVID-19 may experience longer term symptoms. Post-Covid Syndrome (PCS), which may also be known as Long Covid, is recognised in individuals who have persistent symptoms for 12 weeks or more. PCS is a multisystem disease; common symptoms include fatigue, breathlessness and 'brain fog'. Affected individuals may also experience cardiac, musculoskeletal, gastrointestinal and neurological symptoms. As PCS may follow a relapsing course, it is important individuals have fully recovered before being accepted to donate.

Post Donation Information There is no evidence at present that coronaviruses can be transmitted by blood transfusion and therefore these measures are precautionary.

Donors must be provided with information about contacting the blood service if they develop any illness after blood or component donation.

If a donor reports post-donation respiratory illness, refer to [Appendix 4 – Management of post donation illness](#)

Reason for change The deferral after COVID-19 vaccination has been removed.

Update Information This entry was last updated in:
WB-DSG Edition 203 Release 74

Decompression Illness

Obligatory **Must not donate if:**
a) Undergoing investigation or treatment or are still symptomatic.
b) The illness has been complicated by conditions that exclude the donor from donation.

Discretionary If recompression treatment ended more than 24 hours previously, the donor feels well enough to have returned to work / normal daily activities, neither steroid nor anticoagulant drugs have been taken within the previous 7 days, and:

a) Muscle (e.g. limb pain), skin (e.g. lymphatic swelling), or mild neurological symptoms (such as weakness or numbness) have stabilised and the donor has been discharged, accept.

b) Arterial gas embolism has responded fully to recompression treatment, with no evidence for myocardial or cerebral ischaemic event (heart attack/stroke), accept.

See if Relevant [Anticoagulant Therapy](#)
[Cardiovascular Disease](#)
[Cerebrovascular Disease and Intracranial Haemorrhage](#)
[Disabled Donor](#)
[Epilepsy](#)
[Investigations](#)
[Nonsteroidal Anti-Inflammatory Drugs](#)
[Urinary Catheterisation](#)
[Steroid Therapy](#)
[Vertigo](#)

Additional Information Decompression illness incorporates "Decompression sickness" (the bends) and arterial gas embolism. Most events reported by potential donors are likely to relate to diving incidents. The symptoms are caused by bubbles of inert gas (either nitrogen or helium) forming within the tissues (skin, muscle, nerves), or within the circulation, due to inappropriately rapid ascent from depth. This can lead to a broad spectrum of symptoms from mild muscle cramps at one end, to paralysis, heart attack or stroke at the other.

Treatment is a combination of re-pressurising the patient, and increasing the inspired partial pressure of oxygen, which facilitates the gradual removal of the retained inert gas. Additional treatment with nonsteroidal anti-inflammatory drugs (NSAIDs), steroids and anticoagulants may sometimes be used.

Complete relief of symptoms occurs in 50 to 98% of individuals depending on the severity, and period of time between development of symptoms and treatment. Donors who have suffered significant medical problems (heart attack, stroke, paralysis etc.) would be deferred on the basis of this outcome.

Donors with milder symptoms which have either resolved completely, or are considered by the treating physician to have improved as much as they are going to, can be accepted as long as they meet the above criteria, and they have felt well enough to return to normal activities of daily life (housework, employment, driving etc.).

Reason for change Link updated from 'Central Nervous System Disease' to 'Cerebrovascular Disease and Intracranial Haemorrhage' in the 'See if Relevant' section.

Update Information

This entry was last updated in:
WB-DSG Edition 203 Release 73

Dental Treatment

Obligatory

Must not donate if:

- a) Less than seven days since root canal treatment, dental capping (crown or veneer), dental implants or having a tooth removed.
- b) Less than 24 hours since a filling, scale and polish or other superficial treatments.
- c) All wounds are not healed.
- d) There is any infection or the donor has been on antibiotics within the last seven days.
- e) Allogeneic human tissue (bone) has been used.
- f) Less than three months since any invasive dental treatment outside of the UK and Republic of Ireland (ROI).
- g) Receiving, or waiting for, treatment for periodontal gum disease.

Discretionary

- a) If inspection, dental impressions or re-cementing of an existing crown or veneer only, with no requirement for further drilling or local anaesthetic, accept.
- b) If the donor has received an autologous bone graft within the UK or ROI, accept. An autologous graft is derived from the donor's own bone.
- c) If the donor has been treated within the UK or ROI with graft material derived from a non-biological or approved non-human source, accept.
- d) If donor has received graft materials during dental treatment outside the UK and ROI, refer to a **Designated Clinical Support Officer (DCSO)**.
- e) If the donor has completed any treatment for periodontal gum disease, is well with no symptoms of gum disease, and they are only attending a dentist or dental hygienist for cleaning to maintain gum health no more than once every three months, accept.

See if Relevant

Infection - General
Surgery
Tissue and Organ Recipients

Additional Information

Dental extractions, other treatments and active gum disease can result in bacteria getting into the blood stream. The waiting times after treatment are to allow healing and for any bacteria that have entered the blood stream to be cleared.

Gum disease is common. Symptoms include pain, swelling, ulcers and difficulty eating. Donors may declare that they have gingivitis which is inflammation of the gums and which may be acute or chronic, but more chronic or extensive inflammation is usually called periodontal disease. If left untreated, disease can spread to the underlying bones causing teeth to become loose and/or require removal.

Referral to a periodontal specialist may be required, but active significant gum disease will usually be managed to prevent progression, by frequent visits to a dentist or dental hygienist for a course of intensive deep cleaning. Once this course of periodontal treatment is completed then the usual regular schedule of dental hygienist cleaning three-monthly to maintain gum health will resume and continue. Gum disease can be associated with e.g. diabetes or pregnancy, and so care must be taken to ensure a donor is also otherwise eligible.

As there may be uncertainty about infection risks for invasive dental treatment performed outside the UK and ROI, a deferral period of three months is required. Invasive treatments include root canal treatments, dental capping, dental implants and tooth extractions.

Graft materials used in dental procedures are highly processed products, derived from autologous bone, other human bone (allogeneic), animal bone or non-biological materials.

In the UK and ROI, any animal-derived graft material used in dental treatment are approved by regulatory authorities and can be regarded as free from known infection risks. If the donor knows that they received a product derived from an animal, or non-biological) source, the donor can be accepted. If the donor is unsure, advise them to check with their dentist.

For dental surgery performed elsewhere in the world, it may be necessary to request more information about any graft products which were used.

Donors who have had more extensive surgery on their jaw may have received a standard human bone graft. If in doubt, refer to a **DCSO**.

Reason for change Guidance for donors receiving treatment for periodontal disease has been added.

Update Information This entry was last updated in:
WB-DSG Edition 203 Release 72

Dermatitis

Includes Eczema

Obligatory **Must not donate if:**

- a) The venepuncture site is affected.
- b) Large areas of skin are affected.
- c) Taking steroid tablets, injections, or applying steroid, tacrolimus (Protopic[®]) or pimecrolimus (Elidel[®]) creams over large areas.
- d) The donor has needed long term (six months or more) steroid treatment within the last 12 months.
- e) The affected areas are infected.
- f) Less than four weeks from the last dose of Alitretinoin (Toctino[®]).

Discretionary a) If the area affected is small, the venepuncture site is not affected and using topical treatment only, accept.

b) If the donor:

- has been established on oral treatment for their skin disease with only one of the following drugs: Methotrexate, Sulfasalazine, Hydroxychloroquine or Azathioprine, and
- the dose of the drug has not increased in the previous 6 months, and
- their skin disease is controlled by medication, and
- the venepuncture site is not affected, and
- the donor is well,

accept.

c) If there is any uncertainty about the diagnosis or the nature of treatment, refer to a DCSO.

See if Relevant

Allergy
Autoimmune Disease
Infection - General
Monoclonal antibody therapy or other Biological Modalities
Steroid Therapy

Additional Information Dermatitis refers to a group of skin conditions characterised by epidermal change. It may involve both allergic and non-allergic processes. Because of damage to the skin, local

infection is a common problem. For this reason the venepuncture site must not be affected.

Steroid therapy in high doses causes immunosuppression. This may mask infective and inflammatory conditions that would otherwise prevent donation. Long term steroid therapy may also cause temporary adrenal dysfunction. A waiting period of 12 months from the last dose allows time for the adrenal glands to recover.

Some of the treatments used to treat eczema can affect the immune system (e.g. azathioprine (Imuran[®]), ciclosporin, hydroxycarbamide (hydroxyurea, Hydrea[®]), mycophenolate (CellCept[®])) and so can mask signs of infection. This is why systemic treatments (taken by mouth or injection and so affecting the whole body) requires a 12 month deferral period from the time the treatment stops. Under normal circumstances the use of topical treatment with steroid, tacrolimus (Protopic[®]) or pimecrolimus (Elide[®]) will not result in blood levels which cause systemic suppression of the immune response. Systemic suppression is more likely if there is a skin barrier defect or high doses are used over large areas for extended periods. A large area of skin is defined as >9% (Wallace Rule of Nines). 1% is equal to the area of the closed digits and palm of the donor's hand.

Reason for change A discretion to accept donors on oral medication has been added and the text has been updated to ensure consistency with other DSG references to immunosuppression.

Update Information This entry was last updated in:
WB-DSG Edition 203 Release 71

Diabetes Insipidus

Obligatory **Must not donate.**

Additional Information Diabetes insipidus is an unusual condition (about 1 in 25,000 people are affected) where the body cannot retain enough water. It is very different from diabetes mellitus (sugar diabetes). Because of the difficulty in maintaining a normal fluid balance it is considered unwise for a person with this condition to be a donor.

Reason for change 'Additional Information' has been added.

Update Information This entry was last updated in:
DSG-WB Edition 203, Release 01.

Diabetes Mellitus

Also Known As Sugar diabetes and type I (1) and II (2) diabetes.

Obligatory **Must not donate if:**

- a) Requires treatment with insulin.
- b) Diabetes medication has been altered in the last four weeks.
- c) Is having problems with feeling faint, fainting or giddiness.
- d) Has suffered from heart failure.
- e) Has renal impairment requiring dialysis, the use of erythropoietin or similar drugs, or is either under active investigation or continued follow up for renal impairment.
- f) Has required surgery for a blocked or narrowed artery including any type of amputation.
- g) Has or has had gangrene.
- h) Has or has had ulcers or wounds related to a loss of sensation.

i) Has had a transplant of pancreatic tissue.

Discretionary

a) If diagnosed with pre-diabetes or gestational diabetes but not requiring treatment, accept.

b) If controlled by diet or oral medication or injectable medication other than insulin, e.g. Exenatide (Byetta[®]) or Liraglutide (Victoza[®]), that has not been changed in type or dose in the last four weeks, accept.

c) If previous treatment with insulin (including bovine insulin) was stopped more than four weeks ago, accept.

d) If gangrene was not related to diabetes or peripheral vascular disease (e.g. it was due to hypothermia or meningococcal meningitis) and all wounds are fully healed, even if amputation was required, accept.

e) If a donor has a glucose monitoring device to manage their diabetes and is otherwise eligible according to the above criteria, then as long as there is no inflammation or infection at or around the site of the skin sensor, accept.

See if Relevant

Cardiovascular Disease
Cerebrovascular Disease and Intracranial Haemorrhage
Chiropody
Infection - General
Pregnancy
Shunts, Stents and Devices
Tissue and Organ Recipients
Wounds, Mouth and Skin Ulcers

Additional Information

In the UK about one in twenty individuals has diabetes. The majority of cases do not require treatment with insulin. Many people with this type of diabetes (often called type II (2)) are in good health and are fit to donate blood.

It is however important that complications due to diabetes are carefully assessed and, where necessary, donors are excluded from donating (e.g. those at risk of postural hypotension due to autonomic neuropathy, or those at risk of bacteraemia due to unhealed ulcers).

The rationale for not accepting donors on oral medication for diabetes mellitus was reviewed by the Standing Advisory Committee for the Care and Selection of Donors in 2008. It was decided that available data did not support the deferral of all individuals with diabetes that required treatment.

It is a requirement of the Blood Safety and Quality Regulations not to accept donors who are being treated with insulin, or who have received a transplant of human tissue.

Diabetic donors should be informed that blood donation will lower their HbA1c (glycated haemoglobin) levels. This blood test is used to monitor their diabetic control. Donors should inform their diabetic team that they are blood donors so this can be taken into account when reviewing HbA1c levels. Blood donation should preferably be performed after HbA1c testing.

HbA1c decreases under conditions which shorten the life-span of red blood cells (RBC). HbA1c is made when the glucose (sugar) in the body sticks to the RBC. As the body can't use the sugar properly more of it sticks to the RBC and builds up in the blood. RBC are active for around 3 months. By measuring HbA1c, clinicians are able to get an overall picture of what a patient's average blood sugar levels have been over a period of weeks /months. For people with diabetes this is important as the higher the HbA1c, the greater the risk of developing diabetes-related complications.

According to national guidelines people with type 1 diabetes should be offered a continuous glucose monitor (CGM) or flash glucose monitor (e.g. FreeStyle Libre). Some people with type 2 diabetes may also be offered a CGM or flash glucose monitor. Care should be taken to ensure donors are eligible in regard to their diabetes as most will be using insulin, often via a pump, which would preclude them from donation, and/or will be having difficulties monitoring or controlling their blood glucose levels and donation at that time would not be advisable.

Glucose monitoring devices are available to buy and donors with stable diabetes not treated with insulin may have chosen to buy a device to use.

<i>Information</i>	Part of this entry is a requirement of the Blood Safety and Quality Regulations 2005.
<i>Reason for change</i>	Link updated from 'Central Nervous System Disease' to 'Cerebrovascular Disease and Intracranial Haemorrhage' in the 'See if Relevant' section.
<i>Update Information</i>	This entry was last updated in: WB-DSG Edition 203 Release 73

Disabled Donor

- Obligatory*
- 1. All donors must:**
- Fully understand the donation process.
 - Give their informed consent to the process and to the testing of their blood for infections that may affect its suitability for use.
 - Be able to use the bleed facilities provided without risking their own safety or the safety of others (donors must not be bled in a wheelchair).

2. Third party interpreters:

If they are to be present at any part of the selection procedure where there is an exchange of confidential information between the donor and the qualified health professional, they must:

- Understand the requirements of the Blood Safety and Quality Regulations (BSQR) relevant to the donation process.
- Not be personally known to the donor.

- Discretionary*
- Donors with difficulty in reading:**
Ensure by questioning the donor that they:
- Understand and fully complete the tick-box questionnaire.
 - Give valid consent to donation and to the testing of their blood for diseases that may affect its suitability for use.

- See if Relevant*
- Central Nervous System Disease
Urinary Catheterisation
Neurobehavioral Disorders
Spina Bifida

- Additional Information*
- The Services are aware of their duties under Disability Discrimination Legislation and will, whenever and wherever reasonable, try to provide facilities for disabled individuals. Potential donors with a disability are advised to seek advice from their local Blood Service Help Line before attending a donor session to see if their needs can be met. It is however important to note the following.

Some donors, especially those with spinal cord injuries can have significant problems with regulating their blood pressure and as such may be at a greater risk of vasovagal events following blood donation. People who are in wheelchairs are more at risk if they suffer a delayed vasovagal event in the chair, and are alone, as they could remain upright and may suffer prolonged cerebral hypoxia. This can result in permanent brain injury or in extreme circumstances death. For this reason donors must not donate from a wheelchair. Some potential donors may have indwelling shunts and/or catheters in situ which will mean that they are not eligible to donate.

To comply with Part 2 of the Blood Safety and Quality Regulations 2005 (BSQR) every donor must:

be provided with accurate educational materials, which are written in terms which can be understood by members of the general public (Part A 1-13)

complete a health and medical history questionnaire and undergo a personal interview

performed by a health professional (defined in the BSQR as a doctor, a nurse or a donor carer) trained and qualified in the requirements of the BSQR (Part B 15)

provide written informed consent to proceed with the donation process which must be countersigned by the qualified health professional responsible for obtaining the health history (Part B 16 (a) - (f)).

A qualified health professional may assist a donor in the completion of the health and medical history questionnaire and in understanding the consent statement and any other information provided by the Blood Service. To facilitate comprehension it is permissible to use alternative formats (e.g. audio, Braille, computer or alternative language) for the donor information leaflets, the health and medical history questionnaire and consent statements. The donor must be able to clearly demonstrate they have understood this material. At present there is no standardized way of assessing comprehension so this will be a personal judgement made by the qualified health professional.

Use of third party interpreters.

It is permissible for any third party to act as an enabler by helping to reassure the donor and to assist in establishing effective communication between the donor and the qualified health professional. The third party **must not** however be present during any exchange of confidential information, unless they are **not** personally known to the donor and understand the requirements of that part of the BSQR relevant to the donation process. Confidential parts of the process include the evaluation of the health and medical history questionnaire, the medical interview and the obtaining of valid consent. Any third party, with the permission of the donor, may accompany the donor through other parts of the donation process that do not include the exchange of confidential information.

Rationale.

There is concern that the use of third parties during any exchange of confidential information between the donor and the qualified health professional may compromise the confidentiality of the donor and the safety of the blood supply. Interpreters are often part of a close community, or a family member, and this may inhibit or embarrass the potential donor in any confidential exchange of information. This may result in the non-disclosure of sensitive information that could affect the individual's eligibility to donate. If a third party is not fully aware of the relevant aspects of the BSQR this may make the interpretation of information incomplete and potentially put both the donor and the blood supply at risk. There is also a requirement to communicate the results of any testing performed by the Blood Services that may be of relevance to the donor's health in a way that protects their confidentiality. The continuing availability of an independent interpreter, to maintain donor confidentiality, should be taken into account when deciding if an individual donor may be accepted.

To comply with both the BSQR and Health and Safety Regulations no donor can be accepted if it unnecessarily puts their own safety or the safety of others at risk.

<i>Information</i>	Part of this entry is a requirement of the Blood Safety and Quality Regulations 2005.
<i>Reason for change</i>	The See if Relevant section has been revised.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 61.

Diuretics

<i>Also Known As</i>	Water tablets.
<i>Obligatory</i>	Must not donate if: a) Taken for heart failure. b) Taken for kidney failure.
<i>Discretionary</i>	a) If taken for pre-menstrual syndrome, accept. b) If taken to treat hypertension as either the only drug or with other anti-hypertensive medication, accept.
<i>See if Relevant</i>	<u>Blood Pressure - High</u> <u>Cardiovascular Disease</u> <u>Kidney and Bladder Disease</u>

<i>Additional Information</i>	Diuretics (water tablets) are used for many different reasons. If they are taken for a serious condition such as heart or kidney failure the donor should not be accepted.
<i>Reason for change</i>	'Obligatory' and 'Additional Information' entries have been added together with links to 'Cardiovascular Disease' and 'Kidney Disease'. The 'Discretionary' entry has been amended to be consistent with the change to 'Blood Pressure - High'.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 22.

Diverticular Disease

<i>Obligatory</i>	Must not donate if: a) Has symptoms of diverticulitis. b) Less than seven days from completing systemic antibiotic treatment.
<i>Discretionary</i>	If the donor has no symptoms other than mild abdominal pain or constipation, accept.
<i>See if Relevant</i>	<u>Endoscopy</u> <u>Infection - General</u> <u>Investigations</u> <u>Stoma</u> <u>Surgery</u>
<i>Additional Information</i>	Diverticula are pouches sticking out of the side of the large bowel (colon). They become more common as a person ages (50% of people have them by the age of 50, and 70% by the age of 80). Often they are an incidental finding when the large bowel is examined. This is known as diverticulosis and is not a problem. About a quarter of people who have diverticula have symptoms and this is known as diverticular disease. Symptoms are commonly related to pain and constipation but the condition can lead to infection (diverticulitis) and bleeding. Some people may require surgery.
<i>Reason for change</i>	This is a new entry. The previous entry on diverticulosis did not deal with any of the complications of diverticular disease.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 01.

Donor Weight

<i>Definitions</i>	EBV – Estimated Blood Volume. This is calculated using the Nadler formula (Ref: Chapter 3.7 Guidelines for the Blood Transfusion Services in the UK). ECV – Extra Corporeal Volume. This is the total volume outside the donor's circulation at any time during a donation procedure. It includes all blood, plasma and components in the collection packs, the machine harness and testing samples.
<i>Obligatory</i>	1. Must not donate if: a) Under 65 kg (10 stone 3 pounds) b) The donor weight means that they have difficulty in getting onto or off the donation couch. c) Venous access is very difficult. d) The safe weight limit of the bleeding couch/chair is exceeded. e) They are a double red cell donor and weigh under 70 kg (11 stone).
<i>Discretionary</i>	a) If male and over 50kg of weight (7 stone 12 pounds), accept.

- b) If female, 20 years of age or older and over 50kg of weight (7 stone 12 pounds), accept.
- c) If female, less than 20 years of age with an EBV of 3500ml or greater (as per Appendix 1), accept.
- d) Treatment with anti-obesity drugs, accept. This can include treatment with injectable drugs such as liraglutide (Saxenda[®]), provided they have been prescribed by a health care professional.

Component Donation During any planned component donation procedure, the donor's ECV must not exceed 16% of their EBV at any point in the procedure.

Careful consideration should be taken when calculating the EBV for transgender donors to ensure the most appropriate chart is selected.

See if Relevant [Appendix 1 - Estimated Blood Volume for Female donors \(after Nadler\) by height and weight](#)
[Appendix 3 – Maximum permitted ECV for component donation](#)
[Sleep Apnoea](#)

Additional Information Limits on donation volume are in place to protect the donor from adverse effects such as fainting.

There is a minimum legal donor weight of 50kg at which a donation can be accepted. In young women there is a significant risk of fainting if their donation exceeds 15% of their EBV thus a minimum EBV of 3500ml is needed.

For individuals with a body mass index greater than 40, there is a risk that the formula used to calculate blood volume may result in an overestimation of EBV.

The 50kg lower weight limit is not appropriate for double red cell donations because of the increased volume and iron that is being taken from the donor.

Obesity is associated with several medical conditions, including cardiovascular disease, type 2 diabetes, fatty liver disease, gallstones, and gastro-oesophageal reflux disease. Drugs used for obesity treatment include oral or injectable medications. Donors who have been prescribed these can be accepted provided they otherwise meet the donor selection criteria.

Liraglutide for the treatment of obesity is usually marketed as Saxenda[®]. It is provided as a pre-filled pen device with single-use needles. Donors who have been prescribed liraglutide for obesity can be accepted if they are using the equipment provided as per the manufacturer's instructions and are not sharing pens or needles with others.

Information Part of this entry is a requirement of the Blood Safety and Quality Regulations 2005. Other parts are related to research in to the reasons why donors faint.

Reason for change Guidance and information have been added for weight loss drugs, including injectable liraglutide.

Update Information This entry was last updated in:
DSG-WB Edition 203, Release 67

Drug Index

See [Drug Index - preparations which may affect platelet function](#)

Reason for change This is a new entry to link with the Drug Index.

Update Information This entry was last updated in:
DSG-WB Edition 203, Release 01.

Drug Treatment

<i>Obligatory</i>	The taking of some drugs may make a donor ineligible. This could be due to the underlying disease or to the medication. See: Any A-Z index entry for the disease being treated or the drug taken.
<i>Discretionary</i>	Self-medication with some drugs e.g. vitamins, aspirin, sleeping tablets, need not prevent a donation being accepted, providing the donor meets all other criteria. The number of different drugs taken should not of itself make a donor ineligible.
<i>See if Relevant</i>	<u>Acne</u> <u>Alopecia</u> <u>Anti-Androgens</u> <u>Antibiotic Therapy</u> <u>Autoimmune Disease</u> <u>Drug Index - preparations which may affect platelet function</u> <u>Immunodeficiency</u> <u>Immunoglobulin Therapy</u> <u>Lichen Planus</u> <u>Nonsteroidal Anti-Inflammatory Drugs</u> <u>Prostate Problems</u> <u>Psoriasis</u> <u>Steroid Therapy</u>
<i>Additional Information</i>	In most circumstances it is the condition that a drug is being taken for, rather than the drug itself, that will lead to deferral. This is because the amount of drug that will be transfused will be very small. Some drugs are however known to cause birth defects even in tiny amounts. As we do not know who may receive donated blood (it may be transfused directly into an unborn baby) people taking these drugs must be deferred. It is also important to be certain that a particular drug will not stop platelets from working properly. The blood of anyone who has taken drugs in the last seven days that can interfere with platelet function can be used for red cells but may not be suitable for preparing platelets. If a specific drug is not indexed individually, or as a group (e.g. Nonsteroidal Anti-Inflammatory Drugs and Steroids), and the reason for treatment is not a cause for deferral, the donor should be accepted. If in doubt contact a 'Designated Clinical Support Officer' .
<i>Reason for change</i>	Link updated from 'Immunosuppression' to 'Immunodeficiency' in the 'See if Relevant' section.
<i>Update Information</i>	This entry was last updated in: WB-DSG Edition 203 Release 73

Ehlers Danlos Syndrome

<i>Obligatory</i>	Must not donate if: a. there is history of excessive bleeding or bruising b. there is a history of repeated joint dislocation involving the upper limbs c. there are complications due to effects on the heart, cardiovascular system and other organs, e.g. heart valve disease or aortic root involvement, or the donor is under active investigation, treatment or follow up by a specialist d. there is active periodontal disease
<i>Discretionary</i>	If the condition is mild and donor is not prone to excessive bruises or bleeding, even if taking analgesics for joint pain, accept
<i>See if Relevant</i>	<u>Autoimmune Disease</u> <u>Cardiovascular Disease</u> <u>Drug Index</u>
<i>Additional Information</i>	Ehlers Danlos syndrome is usually a mild condition which typically presents with hypermobility of joints, joint pains and tendency to bruising. Most donors are treated

symptomatically, usually with analgesics for joint pains.

If there is doubt about the diagnosis or severity, refer to a Designated Clinical Support Officer.

Endocarditis

<i>Includes</i>	Subacute bacterial endocarditis (SBE).
<i>Obligatory</i>	<p>Must not donate if:</p> <p>a) Has active infection.</p> <p>b) Has a heart defect that limits activity.</p>
<i>See if Relevant</i>	<p><u>Cardiac Surgery</u></p> <p><u>Cardiovascular Disease</u></p> <p><u>Infection - General</u></p> <p><u>Transfusion</u></p>
<i>Additional Information</i>	People with heart problems that may lead to endocarditis (inflammation of the heart lining, heart muscles and heart valves) may not be fit to donate because of either their heart defect or because of treatment for it. This may have included surgery and transfusion.
<i>Information</i>	This is a requirement of the Blood Safety and Quality Regulations 2005.
<i>Reason for change</i>	<p>This new entry replaces the previous entry for 'Subacute Bacterial Endocarditis'. It recognizes that the cause of endocarditis is not always bacterial and the course is not always subacute.</p> <p>The entry has also been changed from the previous entry for 'Subacute Bacterial Endocarditis'. In particular, any risk of developing endocarditis as a result of venepuncture is now considered insignificant. There has also been a NICE review of when antibiotics are needed for prophylaxis against endocarditis. This has resulted in the guideline no longer referring to taking 'antibiotics when having dental treatment' as this is no longer advised.</p>
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 01.

Endometriosis

<i>Discretionary</i>	Accept.
<i>See if Relevant</i>	<p><u>Drug Index - preparations which may affect platelet function</u></p> <p><u>Endoscopy</u></p> <p><u>Nonsteroidal Anti-Inflammatory Drugs</u></p> <p><u>Surgery</u></p>
<i>Additional Information</i>	Endometriosis is a common condition affecting women in their reproductive years. It is caused by the type of cells that usually line the womb occurring elsewhere in the body - usually in the pelvis, outside of the uterus. The cells outside of the womb undergo the same cyclical changes as the ones lining the womb. The commonest symptom is pain and discomfort around the time that a period would be expected. Endometriosis should not normally affect donation but it is important to check if the donor is taking pain killers that might prevent a donation being used for platelet production.
<i>Reason for change</i>	Links and 'Additional Information' have been added.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 01.

Endoscopy

Obligatory

Must not donate if:

Less than four months from an examination with a flexible endoscope.

Discretionary If the examination has been carried out with a rigid endoscope (e.g. colposcopies and most arthroscopies and proctoscopies), the donor is well and not waiting for further tests or results, accept.

See if Relevant Malignancy
Surgery

Additional Information Flexible endoscopes can be difficult to disinfect. There have been cases where infection has been passed from person to person by examination and biopsy using this type of instrument.

This guidance presumes that a validated NAT test for hepatitis C is negative. If this test is stopped, the guidance will change.

A colposcope is an instrument used to view more easily the neck of the womb (cervix). It is not a flexible endoscope so in itself is not a reason for deferral. Care should be taken to ensure that donor is well and not waiting for further tests or results.

The procedure of 'Virtual endoscopy' is a radiographic technique that does not involve the use of an endoscope just a disposable tube it is not in itself a reason for deferral.

Information This is a requirement of the Blood Safety and Quality Regulations 2005.

Reason for change This entry was updated in line with the recommendations of the SaBTO Donor Selection Criteria Review Report published on 23rd July 2017,

Update Information This entry was last updated in:
DSG-WB Edition 203, Release 40

Epilepsy

Obligatory **Must not donate if:**
a) Requiring treatment for epilepsy.
b) Has had an epileptic episode in the last three years.

Discretionary **Previous epilepsy:**
If a person with a past history of epilepsy has, for the past three years, neither required anticonvulsant therapy, nor been subject to fits, accept.

See if Relevant Malignancy
Neurosurgery

Additional Information Faints following donation can lead to epileptiform convulsions. This is caused by a lack of oxygen reaching the brain. This could lead to a true epileptic fit in a person with a recent history of epilepsy.
It may also cause difficulties with the DVLA and/or employment in a person who has been free from fits for some time.

Information This is a requirement of the Blood Safety and Quality Regulations 2005.

Reason for change The 'Discretionary' entry has been modified and further 'Additional Information' has been added.

Update Information This entry was last updated in:
DSG-WB Edition 203, Release 01.

Erectile Dysfunction

Obligatory **Must not donate if:**
Oral or injectable therapy has been obtained abroad or from the Internet from a non-UK registered medical practitioner.

See if Relevant Blood Pressure - High
Cerebrovascular Disease and Intracranial Haemorrhage

Diabetes Mellitus
Prostate Problems

<i>Additional Information</i>	Treatment for erectile dysfunction (including self-injection of UK prescribed drugs) should not normally prevent donation but the underlying cause of the erectile problem might. Drugs obtained while travelling abroad or from non-regulated sources may pose unknown health risks.
<i>Reason for change</i>	Link updated from 'Central Nervous System Disease' to 'Cerebrovascular Disease and Intracranial Haemorrhage' in the 'See if Relevant' section.
<i>Update Information</i>	This entry was last updated in: WB-DSG Edition 203 Release 73

Etretinate

<i>Obligatory</i>	Must not donate if: Has ever taken Etretinate (Tigason®).
<i>See if Relevant</i>	<u>Acne</u> <u>Lichen Planus</u> <u>Psoriasis</u> <u>Skin Disease</u>
<i>Additional Information</i>	Etretinate (Tigason®) is no longer prescribed in many countries because it is highly teratogenic (causes birth deformities) and stays in the body for an extremely long time. It has largely been replaced by acitretin (Neotigason®) which also has restrictions - please follow the relevant link in the index. As it is not possible to know if a donation may be given to a woman in the early stages of pregnancy, individuals who have ever been exposed to this drug cannot donate.
<i>Reason for change</i>	A link has been added to Skin Disease.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 10 Issue 01

Exercise

<i>Discretionary</i>	Providing the donor is well hydrated, recovered from recent exercise and appropriate advice is given concerning post-donation exercise, accept.
<i>Additional Information</i>	People who are planning to undertake exercise after giving blood should be advised that donation may affect their performance and may also increase the risk of bleeding from the venepuncture (needle entry) site and of other adverse events such as fainting. They may wish to wait until the following day so as to avoid any problems. Individuals who undertake sport at high levels of performance should be aware of both the short term affect of blood donation on performance and the possible long term affects if they should become short of iron. They may wish to seek specialist advice on how to avoid adverse affects on their performance from donation.
<i>Reason for change</i>	This is a new entry.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 01.

Eye Disease

<i>Obligatory</i>	Must not donate if: a) Active ocular inflammation or infection (including conjunctivitis, acute glaucoma, iritis or scleritis). b) History of malignancy.
-------------------	--

- c) Ocular tissue transplanted.
- d) Within seven days of receiving injected treatment for age-related macular degeneration (AMD).
- e) The donor declares a history of optic neuritis.

- Discretionary*
- a) If chronic glaucoma treatment is with tablets or drops only, accept.
 - b) Non-injection treatment for age-related macular degeneration (AMD), accept.
 - c) Most donors with poor vision can be accepted but see 'Disabled Donor' if they are not able to read.
 - d) If more than seven days from intravitreal injection treatment for age-related macular degeneration (AMD), accept.
 - e) If completed course of any eye drops following surgery for a benign condition not requiring ocular tissue transplant and there is no active infection or inflammation, accept.

See if Relevant

Autoimmune Disease
Diabetes Mellitus
Disabled Donor
Indwelling Shunts and Stents and Implanted Devices
Infection - General
Laser Treatment
Malignancy
Steroid Therapy
Thrombosis and Thrombophilia
Tissue and Organ Recipients
Central Nervous System Disease

Additional Information

Allogeneic (from another person) ocular tissue may be transplanted in operations other than corneal transplants, including surgery for glaucoma. If surgery was performed after 1997 and any transplanted ocular material was supplied through UK Transplant, this information will be stored on the National Transplant Database held by NHS Blood and Transplant.

There is a risk of bacterial infection and other complications following injection treatment for age-related macular degeneration (AMD). This is why a seven day deferral is required.

Intravitreal injection treatment for wet/age-related macular degeneration is with anti-vascular endothelial growth factor (anti-VEGF) therapy, these include Bevacizumab (Avastin[®]), Ranibizumab (Lucentis[®]) and Pegaptanib sodium (Macugen[®]).

Information

Part of this entry is a requirement of the Blood Safety and Quality Regulations 2005.

Reason for change

An entry re Optic Neuritis has been added as well as a link to Central Nervous System Disease

Update Information

This entry was last updated in:
 DSG-WB Edition 203, Release 38 Issue 01

Faints

Definitions

Delayed Faint:
 Is a faint that occurs after the donor has left the donation venue.

Obligatory

Must not donate if:

- a) History of an unexplained delayed faint.
- b) Two consecutive faints following or during donation.

Discretionary

A donor with a history of a feeling faint on more than one occasion may be accepted following a thorough assessment by a Registered Health Care Professional.

If a donor with a history of fainting is accepted, careful observation is required.

Additional Information

Vasovagal events can range in severity.

When assessing a donor's eligibility to continue to donate following a previous vasovagal event, the Registered Health Care Professional should consider:

- Any contributory factors before and after the donation?
- Was there prolonged recovery after the previous vasovagal event?
- Did the donor sustain an injury?
- Did the donor require treatment from a Health Care Professional outside the blood service?

An unexplained delayed faint occurs when there is no obvious reason for the faint, other than the history of donation. Events that might contribute to a delayed faint would be exertion, dehydration, exposure to an unpleasant situation, or standing for prolonged periods.

A previous history of faints increases the likelihood of a severe adverse reaction to donation.

<i>Reason for change</i>	Updated to include guidance on assessing donors who report a previous vasovagal reaction during or after blood donation.
<i>Update Information</i>	This entry was last updated in: WB-DSG Edition 203 Release 72

Familial Pseudohyperkalaemia

Obligatory **Must not donate**

Discretionary If the donation will not be used to manufacture red cell components for Intrauterine Transfusion (IUT) or transfusion to neonates and infants, accept

Additional Information Familial pseudohyperkalaemia (FP) is a red cell disorder characterised by altered permeability of the red cell membrane. This is an asymptomatic condition which causes increased leakage of potassium from red cells when stored at refrigerated temperatures. High levels of potassium in stored red cell units are a particular risk for transfusion recipients less than one year of age.

The most common FP variant affects about 1 in 400 of the UK donor population (although it is less common in donors from non-European ethnic groups).

Reason for change New entry

Update Information This entry was last updated in:
DSG-WB Edition 203, Release 44

Fertility

Obligatory **Must not donate if:**

- a) Under investigation for infertility.
- b) Any underlying cause or reason for fertility treatment precludes donation.
- c) Undergoing current fertility treatment and monitoring for pregnancy.
- d) Has ever been given human gonadotrophin of pituitary origin.
- e) The donor knows that they have ever been treated with Metrodin HP®.

Discretionary a) If the donor:

- is not undergoing fertility investigation or treatment, and
- is fully recovered from any fertility-related procedure as applicable, and

- is not being monitored for pregnancy, and
- does not have any underlying or associated condition that precludes donation, and
- has never received treatment with human gonadotrophin of pituitary origin, and
- has never received any other treatment that precludes donation,

accept.

b) If a recipient of donated egg(s), embryo(s) and/or sperm is otherwise eligible in regard to fertility investigation and treatment and any underlying or associated condition, accept.

See if Relevant Prion Associated Diseases
Surgery

Additional Information The use of human gonadotrophin of pituitary origin (follicle-stimulating hormone (FSH) and luteinizing hormone (LH)) had stopped in the UK by 1986. The situation in other countries varied so specific dates cannot be given.

Metrodin HP[®] was withdrawn by the Committee on Safety of Medicines in 2003 and following advice from the Medicines and Healthcare products Regulatory Agency the precautionary principle has been applied to withdraw donors who have been treated with this product. Donors treated for infertility after 2003 in the UK will not have been treated with this product. There is no need to confirm that the donor was not treated with Metrodin HP[®]. Donors may have received other non-pituitary derived gonadotrophins which are acceptable.

Recipients of donated eggs, embryos and sperm are eligible to donate. This is in accordance with the SaBTO Microbiological Safety Guidelines.

Donors who have undergone egg donation, egg collection for fertility preservation, and surgical sperm retrieval should be assessed regarding any hormone treatment they have received and using the Surgery entry.

Fertility preservation is available for:

- Individuals undergoing treatment (for cancer but also for some benign conditions) that puts them at risk of becoming infertile.
- Individuals considering gender transitioning.
- Elective social egg freezing; for individuals who wish to delay fertility.

Reasons that donors may be undergoing treatments and procedures for fertility include Polycystic Ovary Syndrome (PCOS), Endometriosis, testicular problems, genetic conditions. For one in four individuals the cause of their fertility problems may not be known. Sensitive questioning may be needed to establish any underlying cause or associations that may be relevant to the donor's health and eligibility.

Donors with inherited conditions e.g. mitochondrial disease may undergo fertility-related diagnostic and assistance treatments and procedures. Unaffected carriers may be eligible to donate if they otherwise fulfil the criteria.

Reason for change The eligibility of recipients of donated eggs or embryos is now specified. Inclusion of criteria and additional information regarding any underlying or associated conditions.

Replacement of reference to specific drugs with criteria to preclude donation whilst being monitored for pregnancy.

Addition of link to Surgery entry to facilitate assessment of donors who have had fertility-related procedures.

Inclusion of additional information for donors who have had a fertility-related procedure not due to a primary fertility problem.

Update Information This entry was last updated in:
WB-DSG Edition 203 Release 72

Fibromyalgia

<i>Also Known As</i>	Fibromyositis or fibrositis.
<i>Discretionary</i>	Accept.
<i>See if Relevant</i>	<u>Disabled Donor</u> <u>Drug Index - preparations which may affect platelet function</u> <u>Nonsteroidal Anti-Inflammatory Drugs</u> <u>Steroid Therapy</u>
<i>Additional Information</i>	Fibromyalgia is a common problem affecting soft tissues (muscles, tendons and ligaments) rather than bones or joints. The cause is not known but it is often linked to sleep disorders.
<i>Reason for change</i>	The link to 'Inflammation' has been replaced with more appropriate links. 'Additional Information' has been added.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 01.

Frequency of Donation

Discretionary **1. Whole Blood:**

A minimum interval of 12 weeks between donations should normally be observed.

Donors who regularly attend at intervals of less than 16 weeks should be informed that they are at increased risk of iron deficiency. They should be advised to reduce their frequency of donation to an average of 16 weeks or more.

Donors with genetic haemochromatosis may donate at intervals of less than 12 weeks.

2. Components:

a) Double Red Cells:

A minimum interval of 26 weeks between donations should normally be observed.

Donors who attend at intervals of less than 32 weeks should be informed that they are at increased risk of iron deficiency. They should be advised to reduce their frequency of donation to an average of 32 weeks or more.

Donors with genetic haemochromatosis may donate at intervals of less than 26 weeks.

b) Apheresis Platelets and/or Plasma:

A minimum interval of two weeks between donations should normally be observed. The combined total of platelet and plasma donations in any 12-month period should not be more than 26.

Donors of convalescent plasma can donate at weekly intervals, provided they meet all other requirements for plasma donation. They should not donate more than 26 donations in any 12-month period.

Donors who attend at intervals of less than four weeks may be at increased risk of iron deficiency.

c) Stem Cell Donors:

A donor should not give any routine donations for six months following bone marrow harvest and for three months following peripheral blood stem cell harvest or lymphocyte donation.

d) Donors who change donation type:

Care must be taken to ensure that limits on the frequency of donation are maintained for donors who move between donation types.

The following deferral periods should be applied:

Donors moving from whole blood to component donation (except double red cells):

4 weeks

Donors moving from platelet or plasma component donation to whole blood:

4 weeks since last component donation (and at least 12 weeks since the most recent whole blood donation)

Donors moving from whole blood to double red cell donation:

12 weeks.

Donors moving from double red cell donation to other component donation:

8 weeks.

<i>Additional Information</i>	The various intervals are to minimise the risk of developing iron deficiency, except for the deferral periods following stem cell or lymphocyte donation «which are in place to allow the donor to be available for further stem cell or lymphocyte donations should this be required. Stem cells and lymphocytes are collected by apheresis.
<i>Reason for change</i>	The deferral periods after bone marrow or stem cell donation have been reduced in keeping with WMDA guidelines.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 68

Gall Bladder Disease

<i>Obligatory</i>	Must not donate if: a) Symptomatic. b) Associated with an inherited haemolytic anaemia e.g. spherocytosis.
<i>Discretionary</i>	If recovered from symptomatic disease or has asymptomatic gallstones not associated with an inherited haemolytic anaemia, accept.
<i>See if Relevant</i>	<u>Endoscopy</u> <u>Haemolytic Anaemia</u> <u>Infection - General</u> <u>Malignancy</u> <u>Surgery</u>
<i>Reason for change</i>	Links have been added for 'Endoscopy', 'Haemolytic Anaemia' and for 'Malignancy'.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 01.

Gastroenteritis

<i>Definitions</i>	<p>Acute: Lasting for a limited duration only.</p> <p>Chronic: Continuing or prolonged.</p>
<i>Obligatory</i>	<p>Must not donate if:</p> <p>a) Chronic or associated with inflammatory bowel disease.</p> <p>b) Less than two weeks since full recovery.</p>
<i>Discretionary</i>	If due to irritable bowel syndrome, accept.
<i>See if Relevant</i>	<p><u>Diverticular Disease</u> <u>Infection - General</u> <u>Inflammatory Bowel Disease</u> <u>Irritable Bowel Syndrome</u></p>
<i>Additional Information</i>	Acute gastroenteritis is usually caused by an infection. The Blood Safety and Quality Regulations 2005 require a two week deferral from the time of recovery. Chronic gastroenteritis is most likely to be caused by inflammatory bowel disease or irritable bowel syndrome.
<i>Information</i>	This is a requirement of the Blood Safety and Quality Regulations 2005.
<i>Reason for change</i>	This is an updated entry which encompasses several previous entries, including 'Diarrhoea' and 'Gastric Flu'.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 01.

Gastrointestinal Disease

<i>Obligatory</i>	<p>Must not donate if:</p> <p>a) Ulcerative colitis</p> <p>b) Crohn's disease.</p> <p>c) Malignant.</p>
<i>Discretionary</i>	<p>a) Other conditions may be acceptable but carefully consider the suitability of individuals liable to iron deficiency through impaired iron absorption or blood loss.</p> <p>b) Coeliac disease, accept.</p>
<i>See if Relevant</i>	<p><u>Anaemia - 1. Iron Deficiency</u> <u>Diverticular Disease</u> <u>Gastroenteritis</u> <u>Indwelling Shunts and Stents and Implanted Devices</u> <u>Infection - General</u> <u>Inflammatory Bowel Disease</u> <u>Irritable Bowel Syndrome</u> <u>Surgery</u> <u>Transfusion</u></p>
<i>Information</i>	Part of this entry is a requirement of the Blood Safety and Quality Regulations 2005.
<i>Reason for change</i>	Relevant links have been added.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 01.

Giardiasis

<i>Discretionary</i>	Accept.
<i>Additional Information</i>	This is a local intestinal infection that does not affect donation.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 202, Release 02.

Glycogen Storage Disease

<i>Obligatory</i>	Must not donate if: Suffers from a Glycogen Storage Disease.
<i>Discretionary</i>	If the potential donor suffers from type 0 (glycogen synthase deficiency), type V (McArdle disease), type XI (Fanconi-Bickel syndrome), type XII (Red cell aldolase deficiency), or type XIII Glycogen Storage Disease (Beta-enolase deficiency), accept.
<i>Additional Information</i>	Glycogen storage disease (GSD) is the result of defects in the processing of glycogen synthesis or breakdown within muscles, liver, and other cell types. GSD in humans is genetic caused by an inborn error of metabolism (genetically defective enzymes) involved in these processes. <u>A position statement on Glycogen storage disorders</u> is available in the JPAC Document Library
<i>Update Information</i>	This entry was last updated in: WBDSG-CB Edition 203, Release 36

Gout

<i>Obligatory</i>	See: Is there an entry for any underlying condition? Must not donate if: Related to malignancy.
<i>Discretionary</i>	If any underlying condition is not of itself a reason to defer, even if on treatment, accept.
<i>See if Relevant</i>	<u>Drug Index - preparations which may affect platelet function</u> <u>Nonsteroidal Anti-Inflammatory Drugs</u> <u>Malignancy</u>
<i>Additional Information</i>	Gout is due to high levels of uric acid. This will not affect the quality of the blood but taking nonsteroidal anti-inflammatory drugs may affect the suitability of the donation for platelet production. Gout may be secondary to malignancy or its treatment.
<i>Reason for change</i>	The entry has been modified to include an 'Obligatory' entry, additional links and 'Additional Information'.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 01.

Growth Hormone

<i>Obligatory</i>	Must not donate if: Has ever received human pituitary derived growth hormone.
<i>Discretionary</i>	If treated exclusively with recombinant-derived growth hormone, accept.
<i>See if Relevant</i>	<u>Prion Associated Diseases</u>
<i>Additional Information</i>	The use of human growth hormone of pituitary origin had stopped in the UK by 1986. The situation in other countries varied so specific dates cannot be given.
<i>Information</i>	This is a requirement of the Blood Safety and Quality Regulations 2005.

Reason for change 'Additional Information' on the use of human growth hormone of pituitary origin has been added. The date that this ceased to be used in the UK has been revised from 1987 to 1986.

Update Information This entry was last updated in:
DSG-WB Edition 203, Release 01.

Guillain-Barre Syndrome

Obligatory **Must not donate if:**

- a) Less than 24 months from resolution.
- b) There has been any recurrence of symptoms.
- c) The doctor who managed the donor cannot confirm a typical monophasic Guillain-Barre syndrome that recovered completely within 12 months.
- d) **Refer to a 'Designated Clinical Support Officer'** before accepting a donor.

See if Relevant **If treated with immunoglobulin or plasma exchange:**
Transfusion

Additional Information The cause of Guillain-Barre syndrome is not known but it often follows an infection or immunization. It probably is associated with auto-antibodies to parts of the peripheral nervous system. This guideline is intended to prevent transferring antibody to a person receiving a transfusion that could then affect their nervous system.

Update Information This entry was last updated in:
DSG-WB Edition 202, Release 02.

Haematological Disease

Obligatory **Must not donate if:**

- a) Malignant.
- b) A clonal disorder, e.g. primary polycythaemia (rubra vera), essential thrombocythaemia or monoclonal gammopathy of unknown significance (MGUS).

Discretionary

- a) If following specialist investigation a polycythaemia is not diagnosed as Polycythaemia Rubra Vera, or another myeloproliferative neoplasm, and no treatment or further investigation is planned, accept
- b) If following specialist investigation a thrombocythaemia, or another myeloproliferative neoplasm, is not diagnosed as Essential Thrombocythaemia and no treatment or further investigation is planned, accept

See if Relevant Anaemia
Haemochromatosis
Haemoglobin Disorders
Haemolytic Anaemia
Immune Thrombocytopenia
Malignancy
Polycythaemia and Raised Haemoglobin

Additional Information Clonal disorders result from the proliferation of a single cell. Because they have the potential to become malignant they are treated in the same way as malignancy.

Information This is a requirement of the Blood Safety and Quality Regulations 2005.

Reason for change The discretionary and see if relevant sections have been updated to include the revised Polycythaemia and Raised Haemoglobin entry.

Update Information

This entry was last updated in:
DSG-WB Edition 203, Release 58

Haematuria

<i>Obligatory</i>	Must not donate if: a) Due to infection. b) Due to malignancy. c) Not fully investigated.
<i>See if Relevant</i>	<u>Kidney and Bladder Disease</u> <u>Infection - General</u> <u>Malignancy</u> <u>Prostate Problems</u>
<i>Additional Information</i>	Haematuria has many causes and most will require an individual to be deferred. If a person has not been fully investigated for the cause of their haematuria, they may have an underlying problem that would lead to deferral.
<i>Information</i>	Part of this entry is a requirement of the Blood Safety and Quality Regulations 2005.
<i>Reason for change</i>	The need to be fully investigated has been added under 'Obligatory'. Links have been added for 'Infection - General', 'Malignancy' and 'Prostate Problems'. 'Additional Information' has been added.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 22.

Haemochromatosis

<i>Obligatory</i>	Refer to a 'Designated Clinical Support Officer' if: Therapeutic venesection has been required or is planned.
<i>Discretionary</i>	<ol style="list-style-type: none"> 1. If the donor does not currently require therapeutic venesection, and has not been venesected in the past, accept. Previous blood donation is not considered to be the same as clinical venesection. 2. If the donor has been approved to donate by a DCSSO and is otherwise eligible to donate, accept. Minimum intervals between donations for approved donors will be determined by individual blood services.
<i>See if Relevant</i>	<u>Cardiovascular Disease</u> <u>Liver Disease</u> <u>Diabetes Mellitus</u> <u>Hormone Replacement Therapy</u>
<i>Additional Information</i>	<p>Genetic Haemochromatosis (GH) is an inherited condition that can cause the body to accumulate too much iron. The standard treatment for GH is removal of blood through venesection. Individuals with GH will usually be monitored for iron overload through their GP or hospital clinic, and will be offered venesection if required.</p> <p>Blood from an individual with GH is safe for transfusion as long as the donor meets all other donor selection criteria. However, it is important that GH patients are not under any additional pressure to donate blood. They must be under the care of an appropriate physician who can offer alternative venesection facilities if the donor is unable to donate. For this reason any patient with GH who has been venesected or who currently requires venesection must be approved by a 'Designated Clinical Support Officer' prior to acceptance.</p>

Someone who has a diagnosis of GH following genetic testing but who has no iron overloading may be advised by their physician to donate blood, as this will reduce the likelihood of venesection being needed in the future. Donors in this situation can be accepted without DCSO referral, as long as they have not been venesected in clinic.

Reason for change Clarification of when referral to DCSO is required prior to donation.

Update Information This entry was last updated in:
DSG-WB Edition 203, Release 55.

Haemoglobin Disorders

Obligatory **Must not donate if:**
a) Sickle cell syndrome.
b) Thalassaemia syndrome.
c) Has a high affinity haemoglobin.

Discretionary a) Donors with symptomless traits for abnormal haemoglobin, accept. **Note**, there is special guidance for donors with sickle trait.
b) Donors with thalassaemia trait, accept but advise they may fail the haemoglobin screening test.

See if Relevant Anaemia
Polycythaemia and Raised Haemoglobin
Sickle Cell Trait
Transfusion

Additional Information People with traits for abnormal haemoglobin and thalassaemia may be able to donate if they pass the haemoglobin screening test at the session and have no other problems associated with the trait.

Some individuals with thalassaemia trait have levels of haemoglobin lower than that required to pass the screening test required by the Blood Services. Although this is normal for them, they may never be able to donate.

Individuals with certain 'high affinity' haemoglobins develop polycythaemia because of the reduced oxygen carrying capacity of their blood. This would be detrimental to a recipient of their blood and donation may be harmful to the donor. For these reasons they should not be accepted.

Information This is a requirement of the Blood Safety and Quality Regulations 2005.

Reason for change The see if relevant section has been updated.

Update Information This entry was last updated in:
DSG-WB Edition 203, Release 58

Haemoglobin Estimation

Obligatory The haemoglobin concentration should be estimated each time a potential donor presents.

Lower limits

1. Whole Blood Donors

Must not donate if the haemoglobin concentration is less than:

- a) Female donors: 125 g/L
- b) Male donors: 135 g/L
- c) Not disclosed*: 125 g/L

2. Double Red Cell Donors**Must not donate if the haemoglobin concentration is less than:**

All donors: 140 g/L

3. Component Donors who will only donate plasma**Must not donate if the haemoglobin concentration is less than:**

a) Female donors: 120 g/L

b) Male donors: 130 g/L

c) Not disclosed*: 125 g/L

4. All other Component Donors**Must not donate if the haemoglobin concentration is less than:**

a) Female donors: 125 g/L

b) Male donors: 135 g/L

c) Not disclosed*: 125 g/L

Upper limits**All Donors****Must not donate if the haemoglobin concentration is greater than:**

a) Female donors: 165 g/L

b) Male donors: 180 g/L

c) Not disclosed*: 180 g/L

If a donor is not accepted, the reason why must be explained to them and, if appropriate, advice given to see their own GP.

Discretionary a) Potential donors whose haemoglobin concentration is estimated to be below the acceptable level may be asked to give a further sample of blood for testing by alternative means. If the haemoglobin concentration is not less than the levels shown above, accept.

b) If the haemoglobin concentration is above the upper limits listed above, refer to the Polycythaemia and Raised Haemoglobin entry.

See if Relevant Hormone Replacement and Sex Hormone Therapy
Polycythaemia and Raised Haemoglobin
Transgender and Non-Binary Individuals

Additional Information * Blood Services should have donor selection processes that are inclusive of transgender and non-binary individuals. These may be based on asking donors their Sex Assigned at Birth or asking donors about their gender identity and transgender history. For the purposes of this entry, 'not disclosed' criteria apply to donors who are not comfortable to answer these details. It is important to ensure donors understand the rationale for asking these questions.

Transgender and Non-binary donors may take gender affirming hormone therapy to support their transition. This may change the haemoglobin level in their blood and consideration can be given to changing the Haemoglobin criteria used to assess the donor, based on the therapy the donor is taking. See Transgender and Non-binary Individuals.

A 500 ml donation of whole blood contains about 250 mg of iron. It can take months for the average donor to replace this loss of iron from the diet. Taking a donation from a person with a haemoglobin concentration below the recommended value may make them anaemic.

The lower haemoglobin acceptance limits apply only to plasmapheresis donors who will only donate plasma by apheresis. If it is anticipated that red cells or platelets will be collected during the procedure the donor must be assessed against the limits for 'all other component donor's above.

Component donors giving double units of red cells lose twice as much iron and so it is even more important that they start with a good haemoglobin concentration.

Information Part of this entry is a requirement of the Blood Safety and Quality Regulations 2005.

Reason for change The entry has been revised to include guidance on assessing transgender and non-binary donors. The See if Relevant section has been revised.

Update Information This entry was last updated in:
DSG-WB Edition 203 Release 70

Haemolytic Anaemia

Includes Red cell disorders:

- Enzyme abnormalities e.g. G6PD deficiency, pyruvate kinase deficiency
- Membrane abnormalities e.g. hereditary spherocytosis, hereditary elliptocytosis
- Paroxysmal nocturnal haemoglobinuria

Immune causes:

- Transfusion-related
- Drug-induced
- Autoimmune conditions

Other causes:

- Infection
- Toxins
- Venom
- Trauma e.g. march haemoglobinuria
- Liver disease – e.g. cirrhosis, Wilson's disease, pregnancy-induced including HELLP syndrome
- Malignancy

Excludes For sickle cell syndrome, thalassemia syndrome – see Haemoglobin Disorders

Obligatory **Must not donate.**

Discretionary a) If there is a known cause for the haemolysis that does not otherwise preclude donation (e.g. an adverse reaction to a medicine, march haemoglobinuria or a venomous bite) and the individual is completely recovered, accept.

b) Hereditary elliptocytosis not causing haemolysis or requiring splenectomy, accept.

See if Relevant Autoimmune Disease
Haemoglobin Disorders
Liver Disease
Monoclonal antibody therapy and other Biological Modalities
Splenectomy
Steroid Therapy
Transfusion

Additional Information Causes of haemolytic anaemia include red cell and haemoglobin disorders.

Affected red cells are more likely to break down after collection. This could make the stored blood dangerous to transfuse.

Most cases of hereditary elliptocytosis do not affect red cell survival and may be accepted

Haemolytic anaemia can also be caused by immune reactions and other triggers. Care should be taken to establish the cause or associated condition and any treatment which may preclude donation or affect eligibility e.g. malignancy, splenectomy, transfusion, steroid or monoclonal antibody therapy. Only individuals who have otherwise had a distinct episode from which they have fully recovered, with no ongoing problems, no risk of recurrence and no ongoing specialist follow-up will usually be eligible.

Reason for change Addition of causes and associated conditions in a new 'Includes' section and into the 'See if Relevant' and 'Additional Information' sections with reference to treatments that may also affect eligibility. Addition of 'Excludes' section for clarification.

Update Information This entry was last updated in:
WB-DSG Edition 203 Release 74

Haemorrhoids

Obligatory **Must not donate if:**

	Regular or severe bleeding is reported.
<i>Discretionary</i>	If asymptomatic or occasional mild bleeding only, accept.
<i>See if Relevant</i>	<u>Anaemia - 1. Iron Deficiency</u> <u>Endoscopy</u> <u>Surgery</u>
<i>Additional Information</i>	Regular bleeding from haemorrhoids can lead to a shortage of iron. This would be made worse by donation and is likely to cause anaemia.
<i>Reason for change</i>	The 'Discretionary' entry has been modified, a link has been added to 'Endoscopy' and 'Additional Information' has been added.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 01.

Hair Removal

<i>Discretionary</i>	Unless the technique has lead to wounds or infection, accept.
<i>See if Relevant</i>	<u>Infection - General</u> <u>Wounds, Mouth and Skin Ulcers</u>
<i>Additional Information</i>	There are many different ways of removing hair, including creams, waxing, electrolysis and the use of co-cyprindiol (Dianette®). Providing there are no wounds or infection, the donor may be accepted.
<i>Reason for change</i>	This is a new entry.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 01.

Hazardous Activity

<i>Definitions</i>	<p>Hazardous Activity: Is something that may put either the donor or others at high risk of serious injury or death if the donor were to suffer a delayed faint following donation. Such events are uncommon but not unknown.</p> <p>This is of necessity a risk reduction exercise rather than an elimination of risk. As an example, the consequences of a driver losing control of a large goods vehicle is likely to be worse than if they were at the controls of a car or light van. Some occupations have a requirement not to return to duty until a defined period of time has passed. For others it is sensible to recommend a night's rest before undertaking something that may be considered a hazardous activity.</p>
<i>Obligatory</i>	<p>Must not donate if: a) Required to undertake a hazardous activity, following donation, on the same working day - donors must be advised of the risks of delayed faints and advised not to perform a hazardous occupation or hobby on the same day.</p>
<i>Discretionary</i>	<p>Hazardous occupation: If going off duty, accept.</p> <p>Exposure to hazardous material: If the donor is well and has not been exposed by inoculation or mucous membrane exposure to potentially infective biological material, accept.</p>
<i>See if Relevant</i>	<u>Air Crew and Air Traffic Controllers</u> <u>Health Care Worker</u> <u>Non-Consented Exposure to Human Body Fluids</u>
<i>Additional Information</i>	<p>Examples of hazardous activities include but are not limited to: climbing, diving (all types), flying, motor sport, parachuting.</p> <p>Examples of hazardous occupations include but are not limited to: air traffic controller, climbing ladders or scaffolding, crane or heavy machine operator, diver, emergency response vehicle driver, fire crew, flying, large goods vehicle driver (LGV, HGV</p>

over 7.5 tonnes maximum authorised mass), miner working underground, public service vehicle driver (excluding vehicles with less than eight passenger seats), train driver.

The suggested driving restrictions would not normally apply to drivers restricted to a category B or C1 licence.

Many occupations expose individuals to hazardous materials. In some cases this may require statutory monitoring e.g. exposure to certain types of radiation or to high levels of lead. Provided the individual is well and they have not been directly exposed by inoculation or mucous membrane exposure to potentially infective biological material they should be accepted.

Reason for change The See if Relevant section has been revised.

Update Information This entry was last updated in:
DSG-WB Edition 203, Release 66

Headache

1. Occasional

Discretionary Accept.

See if Relevant Drug Index - preparations which may affect platelet function
Migraine
Nonsteroidal Anti-Inflammatory Drugs

Update Information This entry was last updated in:
DSG-WB Edition 203, Release 01.

Reason for change Links have been added to 'Drug Index' and 'Nonsteroidal Anti-Inflammatory Drugs'.

2. Regular

Obligatory **Must not donate if:**
Not investigated.

Discretionary If investigated and diagnosis does not contra-indicate donation, accept.

See if Relevant Drug Index - preparations which may affect platelet function
Migraine
Nonsteroidal Anti-Inflammatory Drugs

Additional Information Headache has many causes and some will require an individual to be deferred. If a person has not been fully investigated for the cause of their headache, they may have an underlying problem that would lead to deferral.

Update Information This entry was last updated in:
DSG-WB Edition 203, Release 01.

Reason for change Links have been added to 'Drug Index' and 'Nonsteroidal Anti-Inflammatory Drugs'.
'Additional Information' has been added.

Update Information This entry was last updated in:
DSG-WB Edition 203, Release 01.

Update Information This entry was last updated in:
DSG-WB Edition 203, Release 01.

Health Care Worker

Definitions Non-Consented Exposure to Human Body Fluids:
A non-consented injury or assault in which an individual is exposed to potentially infective material that could be transferred through donation. The causes may range from a sharps

injury to bites, punches and abrasions or sexual assault where mucous membranes have been contaminated with human blood or other body fluids. It also applies to any inoculation injury with abnormal prions from any species.

1. History of Non-Consented Exposure to Human Body Fluids

See Non-Consented Exposure to Human Body Fluids

2. No History of Non-Consented Exposure to Human Body Fluids

Discretionary Accept.

See if Relevant Infectious Diseases - Contact With
Non-Contagious Diseases - Contact With

Additional Information Health care workers should normally be accepted. It is however important to ensure that they have not suffered any relevant events that might put them at risk of infection.

It is also important to ensure that they have not been put at significant risk of infectious diseases through patient or sample contact that may prevent them from donating. Such contact would be exceptional and they should be aware of any potential threat to their own health.

Contact with MRSA and other common hospital acquired infections should not normally prevent donation.

Reason for change The 'Definitions' section was updated as part of the implementation of recommendations from the FAIR study.

Update Information This entry was last updated in:
DSG-WB Edition 203, Release 57.

Henna Painting

Also Known As Hina and mehndi.

Discretionary Accept.

See if Relevant Body Piercing

Additional Information Traditional henna painting (also known as mehndi or hina) is sometimes referred to as tattooing but it does not involve skin piercing and so does not represent a transfusion hazard. The dye binds permanently with proteins in the skin and so the effect can last several months.

Reason for change 'Additional Information' has been added.

Update Information This entry was last updated in:
DSG-WB Edition 203, Release 01.

Hepatitis

Obligatory **See:**
Any specific A-Z index entry for the cause of the hepatitis.

Discretionary If fully recovered from non-viral hepatitis, accept.

See if Relevant Addiction and Drug Abuse
Autoimmune Disease
Hepatitis A
Hepatitis B
Hepatitis C

Hepatitis E
Hepatitis of Unknown Cause

<i>Additional Information</i>	Hepatitis has many causes, including alcohol abuse, autoimmune disease, infection (viral, bacterial and parasitic) and inflammation caused by drugs and toxins. The major concern is with viral hepatitis that can be transmitted by transfusion. Individuals who have fully recovered from non-viral hepatitis may donate if they fully comply with all other selection criteria.
<i>Reason for change</i>	The entry has been re-written to make it clear that there are many different causes of hepatitis.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 01.

Hepatitis A

1. Affected Individual

Obligatory **Must not donate if:**

- less than 6 months from recovery of symptoms, and
- less than 6 months since the donor was diagnosed with hepatitis A infection following laboratory testing.

See if Relevant Travel

2. Current or Former Sexual Partner of Affected Individual

Obligatory **Must not donate if less than 6 months:**

- since a current sexual partner has recovered from symptoms of hepatitis A, or
- since a current sexual partner tested positive for hepatitis A RNA, or
- since last sexual contact with a former sexual partner who had hepatitis A.

3. Person Currently or Formerly Sharing a Home with an Affected Individual

Obligatory **Must not donate if less than 6 months:**

- from recovery of the last affected person in the home, or
- since a person sharing a home tested positive for hepatitis A RNA.

4. Immunisation

Obligatory **Known exposure:**
Must not donate if:
 Less than six months post the last known contact with the affected individual even if vaccine or intramuscular immunoglobulin was given.

Discretionary **No known exposure:**
 Accept.

See if Relevant Hepatitis B - 4. Immunisation
Travel

Additional Information Hepatitis A is a viral infection of the liver, usually spread by the faecal -oral route or by sewage-contaminated food and water. It is rare in the UK with most cases occurring in people returning from travel to endemic countries. Household contacts of cases are at risk of

infection. It can also be spread sexually. Transfusion-transmitted infection is known to occur.

Hepatitis A usually presents with malaise, fever and abdominal symptoms followed by the onset of jaundice, although some individuals may be asymptomatic. Most people recover after a few weeks but in a small number of cases, infection can lead to more severe liver disease and death. Hepatitis A does not cause long term infection. People who have recovered from hepatitis A have life-long immunity.

Blood services may screen for hepatitis A infection using a test for hepatitis A virus RNA. Donors who are diagnosed with hepatitis A infection following blood donation screening or as part of an outbreak investigation must be deferred for 6 months, even if they do not have any symptoms of the disease. After 6 months, they can return to donate without further testing.

Hepatitis A immunisation is often given before travel to parts of the world which also have a risk of infections such as malaria or tropical viruses. The donor's travel history should be checked if they have had hepatitis A vaccine.

Hepatitis A immunisation is sometimes given in combination with hepatitis B immunisation. Refer to the Hepatitis B entry if necessary.

Reason for change The entry has been updated to include testing by blood services for hepatitis A virus. The layout has also been reorganised.

Update Information This entry was last updated in:
WB-DSG Edition 203 Release 70

Hepatitis B

Definitions **HBV:** Hepatitis B virus
HBsAg: Hepatitis B surface antigen
Anti-HBs: Antibody against hepatitis B surface antigen
Anti-HBc: Antibody against hepatitis B core antigen

Active hepatitis B infection refers to an individual with circulating HBsAg and/or HBV DNA. This term includes acute and chronic hepatitis B infection.

Recovered hepatitis B infection refers to an individual who was previously diagnosed with hepatitis B infection but has subsequently cleared HBV from their circulation.

Current exposure refers to an individual who has recent household or sexual contact with an individual with active or recovered hepatitis B infection. 'Recent' is defined as the last 4 months for household contacts or the last 3 months for sexual contacts.

Previous exposure refers to an individual who has been a household contact more than 4 months ago, or a sexual contact more than 3 months ago, of someone with active or recovered hepatitis B infection.

The longer time period specified for household contact is a requirement of the Blood Safety and Quality Regulations (BSQR).

Hepatitis B (HBV) Testing: Unless otherwise stated, HBV testing refers to routine HBsAg and pooled HBV DNA (NAT) testing.

1. Active or recovered hepatitis B infection

Obligatory **Must not donate**

This includes donors who are identified as anti-HBc positive by Blood Transfusion Service testing.

Discretionary a) If:

- The history of HBV infection is not certain, and
- It is more than 12 months from recovery, and
- A test for Anti-HBc will be performed,

accept for donation or take samples for testing, as directed by local procedures.

b) If the donor has been tested previously by the Blood Transfusion Service and no new risks are disclosed, accept.

*Post-session
review of results*

a) **anti-HBc positive donors.**
Must not donate

b) **anti-HBc negative donors who reported hepatitis B infection. If:**

- more than 12 months from recovery, and
- The donor is negative for all markers (HBsAg, screening HBV DNA and anti-HBc negative)

accept.

*Additional
Information*

SaBTO have recommended that all donors are tested at least once for anti-HBc, which is a marker of hepatitis B infection. Individuals who have recovered from hepatitis B will remain anti-HBc positive. There is a risk that reactivation of hepatitis B virus in such an individual could give rise to occult hepatitis B infection which is not detected by routine testing. For this reason, donors who are anti-HBc positive are deferred from donation.

SaBTO included in their recommendations a discretion that anti-HBc positive donors could be accepted if (1) they have adequate immunity to HBV, as demonstrated by an anti-HBs result of greater than 100 iu/l in a validated assay within 24 months of donation and (2) all of their donations will be tested by individual HBV DNA testing. UK Blood Transfusion Services are not currently implementing this approach. These guidelines will be revised should any service implement this additional testing at a future date.

It is likely that donors who are anti-HBc negative have not had hepatitis B in the past. They do not require any additional testing once the diagnosis of hepatitis B has been excluded.

2. Individuals with current exposure to someone with active HBV infection

Obligatory **Must not donate.**

*Additional
Information*

Donors who have a sexual partner or household contact with active HBV infection may be at risk of acquiring HBV. Current guidelines do not allow a donor with a recent or ongoing risk of exposure to HBV to give blood, even if the donor is fully vaccinated against HBV.

3. Individuals with current exposure to someone with recovered HBV infection

Obligatory **Must not donate**

Discretionary

a) If:

- it is at least 3 months after a sexual partner recovered from hepatitis B, and
- it is at least 4 months after a household contact recovered from hepatitis B, and
- a test for Anti-HBc will be performed,

accept for donation or take samples for testing, as directed by local procedures.

b) If the donor has been previously tested by the Blood Transfusion Service, and no new risk is disclosed, accept.

*Post-session
review of results*

a) If the donor is negative for all hepatitis B markers, including anti-HBc, the donor can be accepted. Additional hepatitis B testing is not required for future donations unless the donor discloses a new risk.

b) If any of HBsAg, anti-HBc or HBV DNA are positive, refer to Section 1: Active or recovered hepatitis B infection.

Additional Information

The risk of acquiring hepatitis B infection from someone who has recovered from hepatitis B is very low. Testing for anti-HBc will rule out the possibility that the donor picked up HBV at an earlier stage when their sexual or household contact may have been infectious.

If the sexual or household contact has no history of hepatitis B but has been told they are anti-HBc positive, indicating previous infection only, it is likely that several months have elapsed since the contact cleared the virus from their circulation. The donor can be accepted for donation if anti-HBc testing will be undertaken on the donation, as long as their contact has not had an unexplained illness consistent with hepatitis B in the previous 3 months (sexual partner) or 4 months (household contact). There is no requirement to test the sexual partner or household contact.

4. Individuals with previous exposure to someone with active or recovered hepatitis B

Obligatory **Must not donate**

Discretionary

a) If:

- it is at least 3 months since sexual contact, and
- it is at least 4 months since household contact, and
- a test for anti-HBc will be performed on the donation,

accept for donation or take samples for testing, as directed by local procedures.

b) If the donor has been tested previously by the Blood Transfusion Service and no new risks are disclosed, accept.

Post-session review of results

a) If the donor is negative for HBV and Anti-HBc all hepatitis B markers, including anti-HBc, the donor can be accepted. Additional hepatitis B testing is not required for future donations unless the donor discloses a new risk.

b) If any of HBsAg, anti-HBc or HBV DNA are positive, refer to Section 1: Active or recovered hepatitis B infection

5. Individuals undergoing Hepatitis B immunisation

Obligatory **a) Known Exposure:**
Must not donate.

b) No Known Exposure:
Must not donate if:
Less than seven days since the most recent dose of vaccine was given.

Discretionary

a) Known Exposure:
If it is more than 4 months from the date of exposure, samples can be taken for HBV and anti-HBc testing. No donation should be taken.

b) No Known Exposure:
If it is more than 7 days from the date of the most recent dose of vaccine, accept

Post-session review of results

a) If the donor is negative for HBV and Anti-HBc, the donor can be accepted. Additional hepatitis B testing is not required for future donations unless the donor discloses a new risk.

b) If any of HBsAg, anti-HBc or HBV DNA are positive, refer to Section 1: Active or recovered hepatitis B infection

See if Relevant

Hepatitis A – 4. Immunisation
Immunoglobulin Therapy

<i>Additional Information</i>	Specific HBV immunoglobulin may be used in the management of individuals who have been exposed to hepatitis B. Administration of hepatitis B vaccine can lead to low level reactivity in HBsAg screening assays. For this reason, donors must be deferred until at least 7 days after receiving a dose of vaccine, even if they have not been exposed to HBV. Hepatitis B vaccine is sometimes given in a combined vaccine with hepatitis A vaccine.
<i>Reason for change</i>	The instruction to take samples rather than a donation in sections 1, 3 and 4 has been removed. These donors can be accepted for donation or samples in line with local policies and procedures.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 68

Hepatitis C

1. Person with current Hepatitis C infection

<i>Obligatory</i>	Must not donate.
<i>See if Relevant</i>	<u>Blood Safety Entry</u>
<i>Additional Information</i>	<p>Hepatitis C Virus (HCV) is a serious infection that can lead to chronic liver disease, liver cancer (hepatoma) and chronic fatigue syndrome. It has also been linked with malignant lymphomas and autoimmune disease. The infection is very easily spread by transfusion.</p> <p>Individuals who are chronically infected are sometimes referred to as 'carriers'. They often have no, or minimal, symptoms associated with their infection.</p> <p>Many cases are linked to previous drug use and, before the introduction of HCV screening of blood donations, to transfusion.</p>
<i>Reason for change</i>	See below

2. Person with treated Hepatitis C infection

<i>Includes</i>	Individuals who have received successful treatment for HCV.
<i>Obligatory</i>	Must not donate.
<i>Discretionary</i>	If six months has elapsed from the completion of therapy and the individual has been told their treatment has cleared their HCV infection, accept. Samples should be taken to confirm that the donor is HCV antibody negative before a full donation is taken.
<i>See if Relevant</i>	<u>Blood Safety Entry</u>
<i>Additional Information</i>	Individuals who have been successfully treated will usually remain HCV antibody positive for many years. As a negative HCV antibody screening test is required before blood can be issued, their blood cannot be used.
<i>Reason for change</i>	See below

3. Current or Former Sexual Partner of Affected Individual

Obligatory

Must not donate if

Less than three months from the last sexual contact.

Discretionary Donors who have a current sexual partner with a history of previous HCV infection may be able to donate, depending on the status of their partner:

1. If the partner has been treated for HCV infection, and has been free of therapy for six months, and is in sustained remission, accept.

2. If the partner has evidence of previous HCV infection (HCV RNA negative, anti-HCV positive), and has been fully assessed by an appropriate clinician who has confirmed that there is no current HCV infection, accept.

See if Relevant Blood Safety Leaflet

Additional Information Confirmation of the success of treatment of the HCV positive partner is not required.

Individuals who remain HCV RNA negative six months after completing treatment are likely to have been 'cured', with a risk of relapse of less than 1%

In the United Kingdom the risk of sexual transmission of HCV from an infected individual to a sexual partner is low, but not zero.

As the treated individual would have a very low (<1%) risk of relapse of infection and sexual transmission of the hepatitis C virus is rare, the transmission of hepatitis C from a successfully treated individual to a sexual partner is most unlikely.

All donations in the UK undergo HCV NAT screening so that the chance of a window period donation escaping detection is also exceedingly low (estimated residual risk for HCV transmission from a UK blood donation for 2014-2016 is 1 in 95.8 million donations).

Sexual Partners of anti-HCV positive, PCR negative donors

Individuals who have cleared an acute HCV infection naturally are sometimes identified through HCV testing, including testing of a blood donation. Such individuals will be HCV RNA negative but HCV antibody positive. Unlike people who have been treated for HCV infection, they may not have received appropriate clinical follow up, including repeat HCV RNA testing several months after the original negative result. Unless the individual has undergone specialist assessment and been given the assurance that they are not currently infected, their partner is not eligible to donate.

Reason for change See below

4. Person Currently or Formerly Sharing a Home with an Affected Individual

Discretionary Accept.

See if Relevant Current or Former Sexual Partner of Affected Individual, above.

Additional Information Hepatitis C is neither contagious nor spread by the faecal-oral route. It is usually only spread through a direct blood to blood route. For these reasons household contacts do not need to be deferred.

Reason for change See below

Reason for change This entry was updated to remove the reference to a separate entry for Northern Ireland. This is to reflect changes in donor selection criteria for donors in Northern Ireland (1st June 2020) which are in line with the other UK Blood Services and the SaBTO Donor Selection Criteria Review Report (2017).

Update Information This entry was last updated in:
DSG-WB Edition 203, Release 50

Hepatitis E

<i>Obligatory</i>	Must not donate if: Less than 6 months from recovery
<i>Discretionary</i>	If less than 6 months from recovery and documented HEV RNA negative and anti HEV IgG positive, accept.
<i>See if Relevant</i>	<u>Travel</u>
<i>Additional Information</i>	Hepatitis E is an infectious hepatitis that is usually spread through contaminated food or water. Infection may be associated with travel to countries with poor hygiene/sewage conditions but increasingly, cases of hepatitis E are being identified in the UK usually due to consumption of undercooked contaminated meat. Hepatitis E can affect non-human animals and has been found in pigs in the UK. There have been reports of transmission by transfusion. Infection in healthy individuals is often symptom free but in people with underlying problems in their immune systems it can be serious and occasionally fatal.
<i>Reason for change</i>	The deferral for household and sexual contacts has been removed.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 41

Hepatitis of Unknown Cause

<i>Definitions</i>	HBV: Hepatitis B virus HBsAg: Hepatitis B surface antigen Anti-HBs: Antibody against hepatitis B surface antigen Anti-HBc: Antibody against hepatitis B core antigen
--------------------	---

1. Person with Hepatitis of Unknown Cause

Obligatory **Must not donate.**

2. Person with previous (recovered) Hepatitis of Unknown Cause

Obligatory **Must not donate if:**
Less than 24 months from recovery.

Discretionary a) If:

- it is more than 12 months from recovery, and
- a test for Anti-HBc will be performed,

accept for donation or take samples for testing, as directed by local procedures.

b) If the donor has undergone previous testing by the Blood Transfusion Service, accept.

c) If more than 24 months from recovery, accept.

Post-session review of results a) anti-HBc positive donors
Must not donate

b) If the donor is negative for all HBV markers (HBsAg, screening HBV DNA and anti-HBc, accept.

Additional Information Most hepatitis of unknown origin will have been due to hepatitis A or hepatitis E (or non-viral causes).

Additional testing for those who give a history of hepatitis between 12 and 24 months previously ensures anti-HBc testing is carried out, to exclude hepatitis B infection.

After 24 months, donation testing for anti-HBc will be carried out routinely, as it will be at least two years since the donor last gave a blood donation or samples for testing by the transfusion service. These guidelines will be revised if policy for anti-HBc testing changes.

3. Household or Sexual Contact of someone with Hepatitis of Unknown Cause

- Obligatory* **a) Must not donate if:**
- Less than 12 months from last household or sexual contact; or
 - if ongoing household or sexual contact, less than 12 months from recovery of the sexual or household contact
- b) If the household or sexual contact is thought to have had hepatitis B infection, refer to the Hepatitis B entry

See if Relevant Hepatitis B

Additional Information The 12-month deferral period is to avoid transmission of any infection through transfusion.

There are different rules for contact with someone who has had hepatitis B. These should be followed if the donor reports that a household or sexual contact was most likely to have had hepatitis B.

Reason for change The entry has been updated to use a similar structure to the entry for HBV.

Update Information This entry was last updated in:
DSG-WB Edition 203, Release 68

Herpes Simplex

Includes Genital and oral herpes.

Obligatory **Must not donate if:**
Fresh lesions.

Discretionary a) If lesions are healing (scabbing over) and there is no tingling, accept.
b) If the donor is not immunosuppressed but taking long term prophylaxis with oral antiviral agents, accept.

See if Relevant Immunodeficiency
If there is a history of other sexually transmitted infections, see:
Sexually Transmitted Disease

Additional Information The herpes simplex viruses (HSV 1 and 2) can cause both cold sores and genital herpes. When the virus is actively multiplying it can cause tingling in the affected area and sores. There is a theoretical risk that the virus, or any secondary infection, could be passed on through transfusion. This is why donors with an active infection are not allowed to donate.

There is no need to defer donors who have a sexual partner with Herpes.

Reason for change Link updated from 'Immunosuppression' to 'Immunodeficiency' in the 'See if Relevant' section.

Update Information This entry was last updated in:
WB-DSG Edition 203 Release 73

HIV

Includes AIDS.

1. Affected Individual

Obligatory **Must not donate.**

See if Relevant Blood Safety Entry

Additional Information HIV (Human Immunodeficiency Virus) infection can destroy the immune system and lead to AIDS (Acquired Immunodeficiency Syndrome). It is known to be transmitted by transfusion. In the early stages of infection the testing used by the Blood Services may not detect the virus allowing it to be passed on by transfusion.

Information This is a requirement of the Blood Safety and Quality Regulations 2005.

Reason for change See below

2. Current or Former Sexual Partner of Affected Individual

Obligatory **Must not donate if:**
Less than three months from the last sexual contact.

See if Relevant Blood Safety Entry

Additional Information HIV infection can be spread through sexual activity, including oral and anal sex. It may however not be transmitted for a long time into a relationship. This could be because the infection becomes more active in the infected partner, the uninfected partner acquires another infection or injury to a mucous membrane, or there is a change in the use of, or failure of, barrier contraceptives (condoms etc.). In the early stages of infection the testing used by the Blood Services may not detect the virus allowing it to be passed on by transfusion.

Waiting three months from the last sexual contact will ensure that any infection is picked up by the tests used by the Blood Services.

Information This is a requirement of the Blood Safety and Quality Regulations 2005.

Reason for change See below

3. Person Currently or Formerly Sharing a Home with an Affected Individual

Discretionary Accept.

See if Relevant 2. Current or Former Sexual Partner of Affected Individual above.

Additional Information HIV is neither contagious nor spread by the faecal-oral route. It is usually only spread through a direct blood to blood or sexual route. For these reasons household contacts do not need to be deferred.

Reason for change See below

Reason for change This entry was updated to remove the reference to a separate entry for Northern Ireland. This is to reflect changes in donor selection criteria for donors in Northern Ireland (1st June 2020) which are in line with the other UK Blood Services and the SaBTO Donor Selection Criteria Review Report (2017).

Update Information This entry was last updated in:
DSG-WB Edition 203, Release 50.

Hormone Replacement and Sex Hormone Therapy

Includes **Hormone Replacement Therapy (HRT):**
Includes any form of sex hormones, such as:

- Tablets, patches or topical gels) as treatment for menopausal symptoms
- Testosterone replacement therapy

Gender Affirming Hormone Therapy:
Masculinising or feminising hormones taken to support transition

Obligatory **See:**
Is there an entry for the condition for which the hormones are being given?

Must not donate if:

- a) Used for malignancy.
- b) A recipient of human gonadotrophin of pituitary origin.
- c) A recipient of human pituitary growth hormone.
- d) A recipient of replacement adrenal steroid hormones.

Discretionary

- a) If treatment is for the menopause, its symptoms, or for osteoporosis prevention, accept.
- b) If treatment is for a shortage of sex hormones, e.g. in some cases of erectile dysfunction and is not related to the treatment of malignancy, accept.
- c) If treated with growth hormone that was exclusively recombinant, accept.
- d) If treated with gonadotrophins that were exclusively non-pituitary derived, accept.

See if Relevant Adrenal Failure
Anti-Androgens
Haemochromatosis
Malignancy
Prion Associated Diseases
Steroid Therapy
Thyroid Disease
Transgender and Non-Binary Individuals

Additional Information There are many reasons why an individual may be deficient in a specific hormone. If this is related directly to malignancy, or to the treatment of malignancy, or to the use of pituitary derived hormones (these have been linked with prion associated diseases), the donor cannot donate in order to protect any person who may receive a donation from that individual.

As well as hormones, donors may take other medication to modify the effect of sex hormones as part of gender-affirming treatment. This may include hormone blockers, such as anti-androgens, which could affect the donor's eligibility.

If there is a risk to the safety of the donor, as may be the case with a deficiency of adrenal steroid hormones, then a donation should not be taken.

Information Part of this entry is a requirement of the Blood Safety and Quality Regulations 2005.

Reason for change The title and scope have been changed to include cross sex hormone therapy.

Update Information This entry was last updated in:
WB-DSG Edition 203 Release 70

1. Affected Individual

<i>Obligatory</i>	Must not donate.
<i>See if Relevant</i>	<u>Blood Safety Entry</u>
<i>Additional Information</i>	HTLV (Human T Cell Lymphotropic Virus I and II) infection can cause serious blood and nervous system disease. It is known to be transmitted by transfusion. In the early stages of infection the testing used by the Blood Services may not detect the virus allowing it to be passed on by transfusion.
<i>Information</i>	This is a requirement of the Blood Safety and Quality Regulations 2005.
<i>Reason for change</i>	See below

2. Current or Former Sexual Partner of Affected Individual

<i>Obligatory</i>	Must not donate
<i>Discretionary</i>	1. If it is more than three months since last sexual contact and a validated test for anti-HTLV antibodies is to be undertaken on the donated component(s), accept 2. If it is more than three months since last sexual contact, the donor has been previously tested for anti-HTLV antibodies by the blood service and this test was performed at least three months after the last sexual contact, accept
<i>See if Relevant</i>	<u>Blood Safety Entry</u>
<i>Additional Information</i>	HTLV infection can be spread through sexual activity. It may however not be transmitted for a long time into a relationship. This could be because the infection becomes more active in the infected partner, the uninfected partner acquires another infection or an injury to a mucous membrane, or there is a change in the use of, or failure of, barrier contraceptives (condoms etc.). In the early stages of infection the testing used by the Blood Services may not detect the virus allowing it to be passed on by transfusion. Waiting three months from the last sexual contact will ensure that any infection is picked up by the tests used by the Blood Services. Blood services in the UK are not required to test all donations for anti-HTLV antibodies. Blood services will need to identify at risk donors at health screening and consider options for discretionary HTLV testing. Otherwise, donors who report sexual contact with an affected individual must be deferred.
<i>Information</i>	This is a requirement of the Blood Safety and Quality Regulations 2005.
<i>Reason for change</i>	See below

3. Person Currently or Formerly Sharing a Home with an Affected Individual

<i>Discretionary</i>	Accept.
<i>See if Relevant</i>	2. Current or Former Sexual Partner of Affected Individual above.
<i>Additional Information</i>	HTLV is neither contagious nor spread by the faecal-oral route. It is usually only spread through a direct blood to blood or sexual route. For these reasons household contacts do not need to be deferred.
<i>Reason for change</i>	See below
<i>Reason for change</i>	This entry was updated to remove the reference to a separate entry for Northern Ireland. This is to reflect changes in donor selection criteria for donors in Northern Ireland (1st June 2020) which are in line with the other UK Blood Services and the SaBTO Donor Selection Criteria Review Report (2017).
<i>Update Information</i>	

This entry was last updated in:
DSG-WB Edition 203, Release 50

Huntington's Disease

<i>Also Known As</i>	Huntington's chorea.
<i>Obligatory</i>	Must not donate if: Symptomatic.
<i>Discretionary</i>	Asymptomatic carriers, accept.
<i>Additional Information</i>	Huntington's disease (HD), is an inherited disorder of the central nervous system. It used to be known as Huntington's chorea or HC. Huntington's disease usually develops in adulthood and can cause a very wide range of symptoms including involuntary movements and memory problems. Involuntary movements could cause problems during the donation process and memory problems could interfere with the selection process.
<i>Reason for change</i>	'Additional Information' has been added.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 01.

Hydrocephalus

<i>Obligatory</i>	Must not donate if: Has an indwelling shunt.
<i>See if Relevant</i>	<u>Neurosurgery</u> <u>Spina Bifida</u>
<i>Additional Information</i>	Indwelling shunts can be a source of bacterial infection. This can be present without symptoms. Bacteria can be a serious threat to anybody receiving blood or blood components. This is because bacteria can multiply to dangerous levels after collection.
<i>Reason for change</i>	'Additional Information' has been added.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 01.

Hypercholesterolaemia

<i>Obligatory</i>	Must not donate if: <ol style="list-style-type: none"> Has caused symptomatic disease. Associated with cardiovascular disease. Is currently being treated with systemic monoclonal antibody therapy e.g. Evolocumab (Repatha[®]), Alirocumab (Praluent[®]). Has been treated with Evolocumab (Repatha[®]) or Alirocumab (Praluent[®]) in the last 4 months. Has been treated with any other monoclonal antibody therapy or biologic treatments in the last 6 months.
<i>Discretionary</i>	<ol style="list-style-type: none"> If has not led to symptomatic disease, even if currently on treatment (other than monoclonal antibody therapy), accept. If it is more than 4 months since cessation of treatment with Evolocumab (Repatha[®]) or Alirocumab (Praluent[®]), accept. If it is more than 6 months since cessation of treatment with any other monoclonal antibody therapy or biologic treatment, accept.

See if Relevant Cardiovascular Disease
Cerebrovascular Disease and Intracranial Haemorrhage

Additional Information Hypercholesterolaemia occurs when the level of cholesterol in the blood is outside of the reference range for the donor's age and sex. Usually this is managed by modifying the diet and often by the use of oral drugs.

Treatment may be with monoclonal antibodies which are administered by subcutaneous injection; these can affect the immune system making individuals more susceptible to infections and/or masking the usual symptoms of an infection, thus increasing the chance that a donor may unknowingly have an infection present at the time of donation. This effect may last until the drug is cleared from the body. Observing a deferral period after cessation of treatment with monoclonal antibody therapy will minimise this risk. The deferral periods advised take into account the characteristics of these drugs, including the time it takes for them to be cleared once treatment stops.

High levels of cholesterol are of themselves not a reason to defer a donor. If the hypercholesterolaemia has led to symptomatic disease, such as cardiovascular problems or transient visual or other neurological problems the donor should not be accepted, even if their cholesterol has returned to normal levels.

It is important to ensure that donors on treatment for hypercholesterolaemia do not have any associated cardiovascular disease.

Reason for change The deferral period following treatment with other monoclonal antibody therapy has been modified to be consistent with the Monoclonal antibody therapy and other Biological Modalities entry. Reference to other biologic treatments has also been included for the same reason.

Update Information This entry was last updated in:
WB-DSG Edition 203 Release 74

Hypnotics

Also Known As Sleeping tablets or sedatives.

Discretionary Accept.

Additional Information Many people take various preparations to aid sleep. This should not normally be a reason not to accept a donor, provided they are otherwise well.

Reason for change 'Additional Information' has been added.

Update Information This entry was last updated in:
DSG-WB Edition 203, Release 01.

Immune Thrombocytopenia

Obligatory **Must not donate if:**
a) Symptomatic.

b) Donor reports platelet count below $120 \times 10^9/l$.

c) Recovered but less than five years from recovery.

This applies to both adult and childhood disease.

Discretionary Individuals who have had a splenectomy and fulfil the other requirements, even if on prophylactic antibiotics, accept.

See if Relevant **If treated with immunoglobulin or plasma exchange:**
Transfusion

If treated with immunosuppressive therapy:
Autoimmune Disease

<i>Additional Information</i>	Donors with reduced platelet counts may suffer from increased bleeding and bruising following a donation. This may have serious consequences. Individuals who do not have problems with bleeding or bruising but know that their platelet count is less than $120 \times 10^9/l$ should not donate, as they also may have problems following venepuncture. There is no need to check the platelet count before whole blood donation if the potential donor has been asymptomatic for more than five years and has been told that their platelet count has recovered to greater than $120 \times 10^9/l$.
<i>Information</i>	This is a requirement of the Blood Safety and Quality Regulations 2005.
<i>Reason for change</i>	The links have been revised. The phrase 'Recovered but has ever had a recurrence' has been removed as this was considered too restrictive. This means individuals who have been splenectomised may be acceptable. The term 'Chronic' has been changed to a numerical value of $120 \times 10^9/l$. 'Additional Information' has been added.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 01.

Immunisation

1. Non-Exposed

<i>See if Relevant</i>	<u>Immunisation - Live</u> <u>Immunisation - Non-Live</u> <u>Smallpox Immunisation</u> If you do not know if an immunisation is live or not, see the A-Z index entry for the type of immunisation or: Refer to a 'Designated Clinical Support Officer'
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 01.
<i>Reason for change</i>	A link has been added to 'Smallpox Immunisation'.

2. Post Exposure

<i>Obligatory</i>	1. BCG: See <u>Immunisation - Live</u> 2. Hepatitis A: See <u>Hepatitis A - 4. Immunisation</u> 3. Hepatitis B: See <u>Hepatitis B - 4. Immunisation</u> 4. Rabies: See <u>Rabies - 2. Immunisation - Post Exposure</u> 5. Smallpox: See <u>Smallpox Immunisation</u> 6. Tetanus:
-------------------	---

See
Tetanus - 2. Immunisation

<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 01.
<i>Reason for change</i>	There have been changes to the layout but not to the actions required.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 01.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 01.

Immunisation - Live

<i>Obligatory</i>	Must not donate if: a) Less than eight weeks from administration. b) The inoculation site has not yet healed.
<i>Discretionary</i>	If more than four weeks from administration of a live immunisation other than smallpox immunisation and the inoculation site has healed, accept.
<i>See if Relevant</i>	<u>Smallpox Immunisation</u> <u>Tuberculosis</u>
<i>Additional Information</i>	Live immunisations use living viruses or living bacteria that will stimulate the immune system but do not normally cause a severe illness. They may however cause severe illness in people who are already unwell and have a weakened immune system. By four weeks, any infection caused by the immunisation should have been controlled and so should not be passed on through donated material. There are special rules for smallpox immunisations.
<i>Information</i>	This entry is compliant with the Blood Safety and Quality Regulations 2005.
<i>Reason for change</i>	Advice has been given from SACTTI that a period of four weeks is sufficient to ensure that there would be no circulating virus at time of blood or component donation for live immunisations other than smallpox.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 05.

Immunisation - Non-Live

<i>Obligatory</i>	1. Post Exposure: See: <u>Immunisation - 2. Post Exposure</u> 2. Hepatitis B: Must not donate if: Less than seven days after administration.
<i>Discretionary</i>	If not exposed, for non-live immunisations other than hepatitis B, if well on the day, accept.
<i>See if Relevant</i>	<u>Hepatitis B</u>
<i>Additional Information</i>	Sensitive assays for HBsAg may be positive following recent immunisation. A positive result can lead to the donation being wasted, unnecessary tests and the need to contact the donor. Note, hepatitis A immunisation may be combined with hepatitis B immunisation.

'Non-Live' immunisations do not use material that can cause infection. This means there is no risk to people receiving donated material from a recently immunised non-exposed donor.

<i>Information</i>	This entry is compliant with the Blood Safety and Quality Regulations 2005.
<i>Reason for change</i>	Guidance updated following the removal of deferral after COVID-19 vaccination.
<i>Update Information</i>	This entry was last updated in: WB-DSG Edition 203 Release 74

Immunodeficiency

<i>Includes</i>	Immunosuppression
<i>Obligatory</i>	<p>Must not donate if:</p> <p>a) Diagnosed with a congenital or acquired condition causing immunodeficiency with increased susceptibility to infection.</p> <p>b) Immunosuppressed due to drug treatment.</p>
<i>Discretionary</i>	<p>1. Donors taking immunosuppressive or immunomodulatory therapy to treat autoimmune disease Refer to the Autoimmune Disease entry.</p> <p>2. Donors with recovered immunosuppression If the underlying cause does not preclude donation, refer to the DCSO.</p> <p>3. IgA deficiency If not experiencing frequent infections, accept.</p>
<i>See if Relevant</i>	<p>Autoimmune Disease Immunoglobulin Therapy Monoclonal antibody therapy and other Biological Modalities Steroid Therapy</p>
<i>Additional Information</i>	<p>Immunodeficiency can mask the body's normal response to some infectious and inflammatory conditions. This could result in diseases that may be transmitted by donation from being missed by the Blood Services. If a donor reports recovery from immunodeficiency or, if the underlying cause was unclear, refer to a 'Designated Clinical Support Officer'.</p> <p>IgA deficiency is relatively common. Most people with this condition are healthy but some individuals may experience frequent infections, especially of the ears, sinuses, gut and lungs. Some blood services may screen donors for IgA deficiency to provide a supply of IgA-deficient blood components.</p>
<i>Reason for change</i>	Entry reworded, with addition of a discretionary section to improve clarity and provide guidance for donors with IgA deficiency. New links added.
<i>Update Information</i>	This entry was last updated in: WB-DSG Edition 203 Release 71

Immunoglobulin Therapy

<i>Obligatory</i>	<p>1. Must not donate if:</p> <p>a) After January 1st 1980 the donor has been treated with intravenous or subcutaneous human immunoglobulin.</p> <p>b) The donor has received multiple intramuscular injections of high dose immunoglobulin.</p>
-------------------	--

c) Immunosuppressed.

2. Donors with recovered immunodeficiency:
Refer to a '**Designated Clinical Support Officer**'.

- Discretionary* a) If the intravenous or subcutaneous human immunoglobulin was given before 1980, accept.
- b) If given routine ante- or post-natal anti-D immunoglobulin only (even if received more than one dose), accept.
- c) If single dose prophylactic immunoglobulin has been given, accept.

See **If treated with intravenous or subcutaneous human immunoglobulin:**
Transfusion

See if Relevant Hepatitis A - 4. Immunisation
Hepatitis B - 4. Immunisation
Immunodeficiency
Prion Associated Diseases
Rabies - 2. Immunisation - Post Exposure
Tetanus - 2. Immunisation

Additional Information Immunoglobulin used before 1980 is unlikely to be affected by vCJD (a prion associated disease).

Single dose intramuscular immunoglobulin is unlikely to pose a significant risk of transmitting vCJD.

Information This entry reflects guidance from the former Committee on the Microbiological Safety of Blood Tissues and Organs of the Department of Health.

Reason for change Link updated from 'Immunosuppression' to 'Immunodeficiency' in the 'See if Relevant' section.

Update Information This entry was last updated in:
WB-DSG Edition 203 Release 73

Infection - Acute

Definitions **Acute:**
Lasting for a limited duration only, with no long lasting carrier stage

Systemic:
Any medicine taken by mouth, injection or suppository. It does not include local skin or nail treatments, or drops or creams used in the eye, ear or nose.

Obligatory **See:**
Is there is a specific A-Z index entry for the condition you are concerned about?

Must not donate if:

a) Infected.

b) Less than two weeks from recovery.

c) Less than seven days from completing systemic antibiotic, anti-fungal or antiviral treatment.

Contact with:
See:
Infectious Diseases - Contact With
Or
Non-Contagious Diseases - Contact With

Discretionary Cold sores, genital herpes and common upper respiratory tract infections such as colds and sore throats but **not** influenza, if recovering, accept.

See if Relevant Chikungunya Virus

Endocarditis
Giardiasis
Herpes Simplex
Malaria
Rabies
Rheumatic Fever
SARS
Sexually Transmitted Disease
Steroid Therapy
Surgery
Tetanus
Thrush
Viral Haemorrhagic Fever
West Nile Virus

<i>Additional Information</i>	<p>Many infections can be spread by donated material. It is important that the donor does not pose a risk of giving an infection to a recipient. Waiting two weeks from when the infection is better and seven days from completing systemic antibiotic, anti-fungal or antiviral treatment makes it much less likely that there will still be a risk of the infection being passed on.</p> <p>There is no evidence that cold sores, genital herpes and common upper respiratory infections such as colds and sore throats can be passed on by transfusion but it is still necessary to wait until any such infection is obviously getting better before allowing anyone to donate.</p> <p>In some situations, although the infection may not be transmissible by donation, there is a duty of care to prevent infection passing to other donors or staff, e.g. an infestation of head lice.</p>
<i>Information</i>	This is a requirement of the Blood Safety and Quality Regulations 2005.
<i>Reason for change</i>	<p>Definitions of 'Acute' and 'Systemic' have been added.</p> <p>A reference to 'contact with' has been added under 'Obligatory'.</p> <p>The links in 'See if Relevant' have been extended.</p>
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 01.

Infection - Chronic

<i>Definitions</i>	<p>Chronic: Continuing, or possibly continuing, infection, even without symptoms or signs of infection.</p> <p>Systemic: Any medicine taken by mouth, injection or suppository. It does not include local skin or nail treatments, or drops or creams used in the eye, ear or nose.</p>
<i>Obligatory</i>	<p>Must not donate.</p> <p>Contact with: See: <u>Infectious Diseases - Contact With</u> Or <u>Non-Contagious Diseases - Contact With</u></p>
<i>Discretionary</i>	<p>1. Acne: Most donors with acne can be accepted but this depends on the type of treatment and lack of any secondary infection.</p> <p>2. Chronic superficial fungal infections: a) If on local therapy only, accept. b) If more than seven days from completing systemic antifungal therapy, accept.</p> <p>3. Typhoid and Paratyphoid If more than seven days from completion of antibiotic course and last symptoms, accept</p>
<i>See if Relevant</i>	<u>Acne</u> <u>Endocarditis</u> <u>Hepatitis</u> <u>Hepatitis A</u>

Hepatitis B
Hepatitis C
Hepatitis E
Herpes Simplex
HIV
HTLV
Malaria
Osteomyelitis
Prion Associated Diseases
Sexually Transmitted Disease
Skin Disease
South American Trypanosomiasis
Steroid Therapy
Surgery
Syphilis
Thrush
Toxoplasmosis
Tuberculosis

<i>Additional Information</i>	Many infections can be spread by donated material. It is important that the donor does not pose a risk of giving an infection to a recipient. Some infections may appear to have resolved but are only controlled by the person's immune system. If material from them is given to a recipient without immunity, severe infection may result. Typhoid and Paratyphoid are gastrointestinal infections which rarely have a chronic carrier state. It is usually caught while travelling. It is passed by the faecal oral route and is not transfusion transmitted.
<i>Reason for change</i>	To add entry for Typhoid and Paratyphoid.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 18.

Infection - General

<i>Definitions</i>	<p>Acute: Lasting for a limited duration only with no long lasting carrier stage</p> <p>Chronic: Continuing, or possibly continuing, infection, even without symptoms or signs of infection.</p> <p>Infectious Diseases: Are infections that can easily be passed from person to person, either through casual or intimate contact.</p> <p>Non Contagious Disease: Is a disease which is not transmitted person to person without the aid of a vector (e.g. a mosquito) or is a disease that is the result of an environmental issue which may be shared e.g. food poisoning.</p>
<i>Obligatory</i>	<p>See: Is there a specific A-Z index entry for the condition?</p> <p>If not, see as appropriate: <u>Infection - Acute</u> <u>Infection - Chronic</u> <u>Pyrexia</u></p> <p>Contact with: See: <u>Infectious Diseases - Contact With</u> <u>Non-Contagious Diseases - Contact With</u></p>
<i>Discretionary</i>	Symptomless carriers of Staphylococcus aureus (including methicillin resistant Staphylococcus aureus (MRSA)), accept.
<i>See if Relevant</i>	<u>Coronavirus Infection (COVID-19)</u> <u>Endocarditis</u> <u>Giardiasis</u>

[Hepatitis](#)
[Hepatitis A](#)
[Hepatitis B](#)
[Hepatitis C](#)
[Hepatitis E](#)
[Herpes Simplex](#)
[HIV](#)
[HTLV](#)
[Malaria](#)
[Osteomyelitis](#)
[Parvovirus B19](#)
[Prion Associated Diseases](#)
[Pyrexia](#)
[Rabies](#)
[Rheumatic Fever](#)
[Sexually Transmitted Disease](#)
[South American Trypanosomiasis](#)
[Syphilis](#)
[Tetanus](#)
[Thrush](#)
[Toxoplasmosis](#)
[Tropical Viruses](#)
[Tuberculosis](#)
[Viral Haemorrhagic Fever](#)
[West Nile Virus](#)

<i>Reason for change</i>	A link to the Pyrexia entry has been added and the See if Relevant section has been updated.
<i>Update Information</i>	This entry was last updated in: WB-DSG Edition 203 Release 72

Infectious Diseases - Contact With

<i>Definitions</i>	<p>Infectious Diseases: Are infections that can easily be passed from person to person, either through casual or intimate contact.</p>
<i>Obligatory</i>	<p>See: Is there a specific A-Z index entry for the condition with which there has been contact.</p> <p>Must not donate if: Within the incubation period for the condition or, if this is not known, less than four weeks from last contact.</p>
<i>Discretionary</i>	<p>a) If the infection is known to lead to permanent immunity (e.g. chickenpox, measles, mumps, rubella, whooping cough) and there is a definite history of past infection with the disease with which contact has occurred, accept.</p> <p>b) Contact with common upper respiratory tract infections such as colds, sore throats, influenza, norovirus and other causes of diarrhoea and vomiting, provided the donor is symptom free, accept.</p> <p>c) Contact with skin conditions which are not transmissible by donated material (e.g. scabies, ringworm, tinea) if no signs of infection, accept.</p> <p>d) Individuals who have been prescribed prophylactic antibiotics after contact with meningitis, anthrax or chlamydia, provided they are symptom free, accept.</p>
<i>See if Relevant</i>	<p> Hepatitis Hepatitis A Hepatitis B Hepatitis C Hepatitis E HIV HTLV Non-Contagious Diseases - Contact With SARS Sexually Transmitted Disease Smallpox Immunization </p>

Syphilis
Tuberculosis

<i>Additional Information</i>	<p>Many infectious diseases can be passed on through donated material, even before a potential donor develops any symptoms of the infection. This may lead to serious infection in the person receiving a donation.</p> <p>Many diseases are not infectious and so are not normally a risk.</p> <p>Contacts with meningitis or anthrax are often prescribed prophylactic antibiotics. These should prevent the disease from developing, so provided the potential donor is well, they may be accepted.</p> <p>If in doubt contact a 'Designated Clinical Support Officer'.</p>
<i>Reason for change</i>	A discretion has been added for contact with norovirus and other causes of diarrhoea and vomiting.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 10 Issue 01

Inflammatory Bowel Disease

<i>Also Known As</i>	IBD.
<i>Includes</i>	Crohn's disease, ulcerative colitis, microscopic colitis, collagenous colitis, lymphocytic colitis.
<i>Obligatory</i>	Must not donate.
<i>Discretionary</i>	<p>If diagnosed with microscopic colitis, collagenous colitis, lymphocytic colitis only, accept if:</p> <ul style="list-style-type: none"> • asymptomatic for more than four months, and • any deferral required for steroid or immunosuppressive therapy has passed.
<i>See if Relevant</i>	<p><u>Autoimmune Disease</u> <u>Immunodeficiency</u> <u>Steroid Therapy</u></p>
<i>Additional Information</i>	<p>Crohn's disease and ulcerative colitis usually have a chronic, relapsing course and require long-term treatment. These diseases will often have systemic effects, e.g. Fatigue or anaemia, and there can be association with other disorders, e.g. arthritis, which effects the donor's general health.</p> <p>Lesions in the gastrointestinal tract of individuals with Crohn's disease and ulcerative colitis caused by the disease can increase the risk of bacteria entering the blood stream. Bacteria in donated material can multiply to dangerous levels during storage.</p> <p>Microscopic colitis and its subtypes (collagenous colitis and lymphocytic colitis) are classified as inflammatory bowel diseases. However, there is a lower likelihood of relapsing disease and longer-term complications. Individuals with microscopic colitis can donate once their symptoms have settled and they feel well. A four-month deferral ensures the donor's recovery is maintained. Care should be taken to ensure that, as well as complete cessation of gastrointestinal symptoms, donors are not experiencing any extra-intestinal symptoms such as fatigue, arthralgia or myalgia that affect normal activities of daily living.</p>
<i>Information</i>	Part of this entry is a requirement of the Blood Safety and Quality Regulations 2005.
<i>Reason for change</i>	Link updated from 'Immunosuppression' to 'Immunodeficiency' in the 'See if Relevant' section.
<i>Update Information</i>	This entry was last updated in: WB-DSG Edition 203 Release 73

Inherited Diseases

<i>Obligatory</i>	See: Is there an A-Z index entry for the condition?
<i>Additional Information</i>	If there is not an index entry for the condition and neither the symptoms nor any treatment are a reason for deferral, the donor is probably acceptable. If in doubt contact a 'Designated Clinical Support Officer' .
<i>Reason for change</i>	The need to refer a to a 'Designated Clinical Support Officer' when there is not a specific entry for the condition has been removed. 'Additional Information' has been added.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 01.

Investigations

<i>Definitions</i>	Radionuclides: These are unstable materials that emit radioactivity when they decay. They are used in some special investigations carried out in radiology (X-ray) and medical physics departments. They may be breathed in, taken by mouth or given by injection.
<i>Obligatory</i>	Must not donate if: Waiting for investigation or the results of investigations for an undiagnosed condition which might lead to deferral.
<i>Discretionary</i>	If for 'routine' investigations, such as attending for a cervical smear, mammogram, a well person clinic when no abnormality is expected, or for the routine monitoring of a condition, such as diabetes controlled by diet or oral medication, which of itself would not be a cause for deferral, accept.
<i>See if Relevant</i>	<u>Endoscopy</u> <u>Prostate Problems</u> <u>Radionuclides</u>
<i>Additional Information</i>	Investigations may lead to the diagnosis of a condition that would lead to deferral. For this reason any investigations for an undiagnosed condition must lead to deferral until the results are known by the potential donor. A decision can then be made as to if the person can be accepted as a donor.
<i>Reason for change</i>	A 'Definition' of 'Radionuclides' has been added. Entries have been added under 'Discretionary', 'See if Relevant' and 'Additional Information'.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 01.

Irritable Bowel Syndrome

<i>Also Known As</i>	IBS.
<i>Discretionary</i>	If the condition has been diagnosed as irritable bowel disease, even if on medication, accept.
<i>See if Relevant</i>	<u>Endoscopy</u> <u>Inflammatory Bowel Disease</u>
<i>Additional Information</i>	Irritable bowel syndrome is due to hyper-activity/sensitivity of the large bowel. It should not be confused with 'Inflammatory Bowel Disease' which would not allow donation.
<i>Reason for change</i>	There has been a change to the wording of 'Discretionary' to improve clarity, a link has been added to 'Inflammatory Bowel Disease' and 'Additional Information' has been added.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 01.

Jaundice

<i>Obligatory</i>	Must not donate if: a) Jaundiced or has a history of jaundice.
-------------------	--

- b) If the cause of the jaundice was viral see the specific A-Z index entry for that condition.
- c) If the jaundice was related to malignancy or to its treatment.
- d) If the cause of the jaundice was not known, treat as Hepatitis of Unknown Cause.

Discretionary a) If fully recovered from a non-viral cause of jaundice (this includes, but is not limited to, physiological jaundice of the newborn, gall stones and drug reactions), accept.

b) If due to Gilbert's syndrome and not visibly jaundiced, accept.

See if Relevant Gall Bladder Disease
Hepatitis A
Hepatitis B
Hepatitis C
Hepatitis E
Hepatitis of Unknown Cause
Liver Disease
Malignancy

Additional Information Transfusion laboratories are unlikely to use blood that appears jaundiced. This means any visibly jaundiced donation is likely to be wasted.

Many things can cause jaundice. The concern is with infectious causes that might be passed on by a transfusion.

Reason for change A new 'Obligatory' entry for jaundice related to 'Malignancy' has been added together with links to 'Hepatitis of Unknown Cause' and to 'Malignancy'.

Update Information This entry was last updated in:
 DSG-WB Edition 203, Release 01.

Kidney and Bladder Disease

1. Acute Nephritis (to include Pyelonephritis, Acute tubular interstitial nephritis and Glomerulonephritis)

Definitions **Pyelonephritis:** acute nephritis due to ascending infection.

Acute tubular interstitial nephritis: acute nephritis caused by an 'allergic reaction' to medication, rarely as part of a systemic often autoimmune disease.

Glomerulonephritis: May be primary intrinsic to the kidney or secondary, associated with certain infections, drugs, systemic disorders (SLE, vasculitis), or diabetes.

Obligatory **Must not donate if:**
 If under active investigation, treatment or specialist follow-up by a specialist.

Discretionary If well, on no treatment and is discharged from follow-up, accept.

See if Relevant Autoimmune Disease

Additional Information Self-limiting renal disease e.g. single attacks of glomerulonephritis or pyelitis, from which recovery has been complete, do not necessarily disqualify the donor.

If there is doubt about the diagnosis refer to a '**Designated Clinical Support Officer**'.

2. Chronic Nephritis

Obligatory **Must not donate.**

3. Infection

<i>Obligatory</i>	Must not donate if: a) Has active infection b) Under investigation c) On antibiotics to prevent urinary tract infection
<i>Discretionary</i>	If the donor has taken a single dose of an antibiotic after sexual intercourse to prevent urinary tract infection, is symptom-free on the day of donation, and does not have an underlying condition that prevents donation, accept.
<i>See</i>	<u>Infection - General</u>
<i>See if Relevant</i>	<u>Antibiotic Therapy</u>
<i>Additional Information</i>	Donors may be taking prophylactic antibiotics long-term or as single post-coital doses. Anatomical anomalies within the urinary tract can make individuals more prone to recurrent infections. If a donor is accepted in accordance with the Discretionary guidance above, the importance of informing the relevant blood service of symptoms suggestive of urinary tract infection occurring within 14 days of donation must be emphasised with the donor.

Donors on long-term antibiotics may have ongoing urinary tract infection without having any symptoms. As an active infection at the time of donation cannot be ruled out, these donors must not be accepted.

4. Kidney Failure

<i>Obligatory</i>	Must not donate if: a) Has renal impairment requiring dialysis. b) Using erythropoietin or similar drugs to increase the haemoglobin concentration. c) Is either under active investigation, or continued follow up by a specialist for renal impairment, or has any associated cardiovascular complications. d) Has had a kidney transplant.
<i>Discretionary</i>	If a kidney transplant was of a non stored autologous organ, accept.
<i>See if Relevant</i>	<u>Autoimmune Disease</u> <u>Blood Pressure - High</u> <u>Diabetes Mellitus</u> <u>Immunodeficiency</u> <u>Tissue and Organ Recipients</u> If treated with blood or blood products, immunoglobulin, plasma exchange or filtration: <u>Transfusion</u>
<i>Additional Information</i>	People with significant kidney failure usually have a high risk of anaemia. This, together with other factors, make them unsuitable as donors.
<i>Information</i>	This is a requirement of the Blood Safety and Quality Regulations 2005.

5. Polycystic Kidney Disease

<i>Discretionary</i>	A diagnosis of polycystic kidney disease does not necessarily prevent donation. If otherwise well, accept.
<i>See if Relevant</i>	<u>Blood Pressure - High</u> <u>Infection - General</u> Kidney Failure above.
<i>Additional Information</i>	Polycystic kidney disease is usually genetic. It varies markedly in its severity and many people will not run into problems until later in their lives. Before this

happens, provided they are otherwise well, there is no reason why affected individuals should not donate. Often they will have higher haemoglobin concentrations than normal.

6. Renal Colic, Kidney and Bladder Stones

<i>Obligatory</i>	Must not donate if: a) Symptomatic. b) Under investigation.
<i>See if Relevant</i>	<u>Infection - General</u>
<i>Additional Information</i>	Renal colic is most commonly caused by solid material (crystals or a stone) passing through the tube that connects the kidney to the bladder (the ureter). It is commonly associated with infection. It is important to wait until the donor is fully recovered and any investigations have been completed. This should avoid a donation being taken from an individual with infection. Infection can lead to bacteria contaminating any donated material. This can be dangerous because bacteria can multiply to dangerous levels in the stored donation. Kidney and bladder stones have many causes and may be associated with infection. It is important to ensure that there is not an underlying cause that would prevent donation.

7. Interstitial Cystitis

<i>Obligatory</i>	Must not donate if: a) Under investigation b) Has an associated condition which would prevent donation c) Has required catheterisation within the last 7 days. d) On treatment with Pentosan polysulfate sodium (Elmiron)
<i>Discretionary</i>	If investigations are complete, there are no associated conditions or treatments which would prevent donation, symptoms are controlled even if on medication other than Pentosan polysulfate sodium (Elmiron), the potential donor has not required catheterisation within the last 7 days and any treatment with Pentosan polysulfate sodium was completed more than seven days ago, accept.
<i>See if Relevant</i>	<u>Autoimmune Disease</u> <u>Chronic Fatigue Syndrome</u> <u>Endoscopy</u> <u>Infection -General</u> <u>Surgery</u> <u>Urinary Catheterisation</u>
<i>Additional Information</i>	Interstitial Cystitis or Painful Bladder Syndrome is a condition which causes chronic or recurrent pain in the bladder and in the pelvic region due to damaged bladder lining or urothelium. The cause is unknown but may be associated with other conditions such as Irritable Bowel Syndrome, Fibromyalgia, Chronic Fatigue Syndrome, Autoimmune Disease and Anxiety Disorder It may also be caused by traumatic injury to the bladder and precipitated by infection. The diagnosis of IC or PBS is one of exclusion. Treatment can be through diet modification, bladder training techniques, exercise and stress management. It can include oral medication with analgesics, antidepressants, and Cimetidine. Treatment can also be with Pentosan polysulfate sodium (Elmiron) which can be associated with increased bleeding and bruising. The condition can also be treated by interventional methods including catheterisation, surgery and botulinum toxin injections.

Investigation and treatment can involve cystoscopy. Use of neuromodulation techniques with a transcutaneous electrical nerve stimulation (TENS) machine does not prevent donation.

<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 61.
<i>Reason for change</i>	Obligatory and discretionary guidance has been added for Infection, including for donors taking antibiotics to prevent urinary tract infection. Relevant links have been included. For Interstitial Cystitis, Obligatory and Discretionary have been amended to avoid repetition, the relevant links have been updated, and reference to the use of cystoscopy has been added to Additional Information.
<i>See if Relevant</i>	<u>Indwelling Shunts and Stents and Implanted Devices</u>
<i>Reason for change</i>	Link updated from 'Immunosuppression' to 'Immunodeficiency' in the 'See if Relevant' section of 4. Kidney Failure .
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 61.
<i>Update Information</i>	This entry was last updated in: WB-DSG Edition 203 Release 73

Klinefelter's Syndrome

<i>Discretionary</i>	Accept.
<i>Additional Information</i>	Klinefelter's syndrome is caused by a chromosomal abnormality that affects males. It may cause low levels of testosterone so that affected men, not on replacement therapy, may have haemoglobin levels in the female range. This may lead to them failing the haemoglobin screening test.
<i>Reason for change</i>	'Additional Information' has been added.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 01.

Laser Treatment

<i>Obligatory</i>	Must not donate if: a) For malignancy. b) Any wounds are not healed.
<i>Discretionary</i>	a) If for basal cell carcinoma (a type of malignancy), treatment is completed and fully recovered, accept. b) If for Cervical Carcinoma in Situ, treatment is completed and a follow up smear did not show abnormal cells, accept. c) If for cosmetic purposes, when healed, accept. d) If for varicose veins, when healed, accept. e) If laser treatment to the eye, when healed, accept.
<i>See if Relevant</i>	<u>Cervical Carcinoma in Situ</u> <u>Diabetes Mellitus</u> <u>Eye Disease</u> <u>Malignancy</u> <u>Surgery</u> <u>Wounds, Mouth and Skin Ulcers</u>
<i>Additional Information</i>	Medical lasers can be used in many different situations. Their action is through heating and burning. The concern is when they are used for treating malignancies and when they leave

areas of tissue damaged and susceptible to infection. Provided the reason that the laser was used is not of itself a reason to defer the donor, once all wounds are healed, so that there is no further infection risk, the donor may be accepted.

If used for diabetic retinopathy it is likely that the donor will need to be deferred and reference should be made to Diabetes Mellitus.

<i>Information</i>	This is a requirement of the Blood Safety and Quality Regulations 2005.
<i>Reason for change</i>	The need for all wounds to be healed has been added under 'Obligatory'. Laser eye treatment has been added under 'Discretionary'. Additional links have been added under 'See if Relevant' together with 'Additional Information'.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 01.

Liver Disease

1. Non-Alcoholic Fatty Liver Disease (NAFLD)

<i>Excludes</i>	Alcoholic Fatty Liver Disease (AFLD)
<i>Obligatory</i>	Must not donate if diagnosed with: <ul style="list-style-type: none"> • Non-alcoholic steatohepatitis (NASH) • Cirrhosis
<i>Discretionary</i>	A diagnosis of non-alcoholic fatty liver disease does not necessarily prevent donation. If the donor is otherwise well and managed with diet and lifestyle changes such as exercise, accept.
<i>Additional Information</i>	NAFLD is a common medical condition, caused mainly by lifestyle factors such as weight, type 2 diabetes, high blood pressure and high cholesterol. There is no drug treatment for this condition. It is usually managed with diet and lifestyle changes along with treatment of any associated medical conditions. Regular monitoring of the condition, e.g. blood tests and liver scans, should not preclude donation. NASH is an advanced form of NAFLD. It is caused by an excessive accumulation of fat in the liver. This can progress to chronic liver inflammation and can result in cirrhosis if untreated.

2. Alcohol-Related Liver Disease

<i>Obligatory</i>	Must not donate.
<i>Discretionary</i>	If the donor is well, and <ul style="list-style-type: none"> • not under specialist follow up, and • has not been diagnosed with alcohol related hepatitis or cirrhosis, accept. Refer to a Designated Clinical Support Officer (DCSO) if there is uncertainty about the diagnosis or the extent of liver damage.
<i>See if Relevant</i>	<u>Addiction and Drug Abuse</u>
<i>Additional Information</i>	Alcohol-related liver disease is common but preventable liver damage that is caused by drinking too much alcohol. It is reversible in the early stages when it is characterised mainly by fatty liver changes. In some individuals it may progress to alcoholic hepatitis and alcoholic cirrhosis.

3. Infective Liver Disease

<i>Includes</i>	Liver abscess, Glandular fever, Viral hepatitis
<i>Obligatory</i>	Refer to the WBDSG entry for the condition. If there is no specific entry, must not donate
<i>Discretionary</i>	If the donor is fully recovered and there is no specific guidance for the condition, refer to <u>Infection – General</u>
<i>See if Relevant</i>	For GlandularFever see <u>Infection - Acute</u> <u>Infection - General</u> <u>Hepatitis</u>

4. Autoimmune Liver Disease

<i>Includes</i>	Autoimmune Hepatitis (AIH), Primary Biliary Cholangitis (PBC) and Primary Sclerosing Cholangitis (PSC)
<i>Obligatory</i>	Must not donate if: <ul style="list-style-type: none"> • under active investigation or treatment, or • associated with Inflammatory Bowel Disease
<i>Discretionary</i>	a) If well, even if on treatment to control symptoms. e.g. Cholestyramine (Questran) or Ursodeoxycholic acid (also known as Ursodiol), accept. b) If well and taking treatment to suppress the condition, refer to the Autoimmune Disease entry
<i>See if Relevant</i>	<u>Autoimmune Disease</u> <u>Hepatitis</u> <u>Steroid Therapy</u>
<i>Additional Information</i>	Autoimmune liver disease in its early stages may be asymptomatic or present with mild symptoms such as itchy skin (pruritis) and fatigue. The donor may require no treatment or treatment for symptom control only for an extended period. Autoimmune liver disease may be diagnosed during investigation for other conditions, especially other autoimmune conditions. Treatment to suppress these diseases may include steroids, Azathioprine and other immunosuppressants. If there is doubt about the diagnosis and treatment, refer to a DCSO .

5. Drug or Pregnancy Induced Liver Disease

<i>Includes</i>	Acute Liver Failure
<i>Obligatory</i>	Must not donate if <ul style="list-style-type: none"> • Under active investigation, treatment or follow up by a specialist • Has received a liver transplant • Has chronic liver failure
<i>Discretionary</i>	If the donor has recovered, is not on treatment and has been discharged from follow up, accept. If there is doubt about the diagnosis, refer to a DCSO .
<i>See if Relevant</i>	<u>Addiction and Drug Abuse</u> <u>Tissue and Organ Recipients</u>
<i>Additional Information</i>	Liver failure may be acute or chronic. Acute liver failure (also known as fulminant liver failure) can be caused by drugs, such as paracetamol overdose, prescription medications, herbal preparations and ingestion of toxins. Liver problems can also occur during pregnancy e.g acute fatty liver of pregnancy (AFLP) and intrahepatic

cholestasis of pregnancy (ICP). Acute liver failure can occur in an individual with no pre-existing liver disease. It is often reversible with full recovery if adequately treated.

Chronic liver failure is caused by longstanding liver disease such as autoimmune liver disease, hepatitis, alcohol related liver disease, liver cirrhosis, haemochromatosis and Wilson's disease.

6. Liver Cirrhosis

Obligatory **Must not donate**

Additional Information Cirrhosis can be caused by many different conditions and by several different liver conditions in combination. Transmissible viruses, some of which are not detected in transfusion service testing, can cause some cases. Because cirrhosis is a sign of worsening or progressive liver disease, it is considered safest not to accept individuals with cirrhosis.

7. Liver Tumours

Includes Liver Cancer, Hepatocellular Carcinoma, Bile Duct Cancer

Obligatory **Must not donate.**

Discretionary Donors with benign liver cysts or adenomas who are fit and well, even if regularly monitored, accept.

See if Relevant Malignancy

Additional Information If in doubt about the diagnosis, refer to a **DCSO**.

8. Inherited Diseases Affecting the Liver

Obligatory Refer to WBDSG entry for the condition.
If there is no specific entry, **must not donate**

Discretionary a) If the donor is well and stable on treatment for Wilson's Disease, accept.
b) If the donor has Gilbert's Syndrome and is not visibly jaundiced, accept
c) For other conditions, see the Inherited Diseases entry

See if Relevant Inherited Diseases
Haemochromatosis

Additional Information Wilson's disease is caused by an excessive accumulation of copper in the liver and other organs. e.g. brain. If diagnosed and treated early with chelating agents, such as Penicillamine and Trientine, and avoidance of high copper foods, the prognosis is good and individuals can lead a normal life. If there is uncertainty about the donor's health or treatment, refer to a **Designated Clinical Support Officer**.

Alpha-1-antitrypsin deficiency can occasionally cause liver disease in adults. This may lead to liver failure and the need for liver transplantation.

Gilbert's syndrome is an inherited defect in bilirubin metabolism. It is harmless but can cause jaundice (yellowing of the whites of the eyes). Blood banks are unlikely to use blood that appears jaundiced. This means any visibly jaundiced donation is likely to be wasted.

Update Information This entry was last updated in:
DSG-WB Edition 203, Release 66

Malaria

<i>Excludes</i>	Donors who will only donate plasma for fractionation. See <u>Malaria - plasmapheresis donors</u>
<i>Obligatory</i>	<p>Must not donate if:</p> <p>a) The donor has ever had malaria.</p> <p>b) The donor has had an undiagnosed fever (which could have been malaria) while abroad or within four months of leaving a malaria endemic area.</p> <p>c) The donor has lived in any malarial endemic area for a continuous period of six months or more at any time of life.</p> <p>d) Less than 12 months after last leaving a malaria endemic area.</p>
<i>Discretionary</i>	<p>a) Donors who have had malaria diagnosed in the past: If more than three years have passed since anti-malarial therapy has been completed and symptoms caused by malaria have resolved and a validated test for malarial antibody is to be performed, accept.</p> <p>If the donor (with a history of malaria) has revisited a malaria endemic area and at least four months have passed since return and a validated test for malarial antibody is to be performed, accept.</p> <p>b) Donors who have EVER had an undiagnosed fever that could have been malaria while in a malarial area or within four months of leaving a malaria endemic area: If at least four months have passed since the donor returned from the malarial endemic area, or from the date of recovery from symptoms (undiagnosed fever) that may have been caused by malaria, whichever is later, and a validated test for malarial antibody is to be performed, accept.</p> <p>NB. this may have to be increased to six months if the area is also identified as a risk area for T.cruzi or a tropical virus; the longest stipulated deferral period must be applied</p> <p>c) Donors who have EVER been resident in a malarial endemic area for 6 months or more: If at least four months has passed since the date of the last potential exposure to malaria, and a validated test for malaria antibody is to be performed, accept.</p> <p>d) For all other donors: If at least four months and less than 12 months have passed since return from a malaria endemic area, and a validated test for malarial antibody is to be performed, accept.</p>
<i>See if Relevant</i>	<u>The 'Geographical Disease Risk Index'</u> for countries with a current endemic malaria risk.
<i>Additional Information</i>	Cases of transfusion transmitted malaria have occurred many years after the donor was last at risk of becoming infected with malaria. This is mainly a problem in people who have had repeated episodes of infection with malaria. Although this is uncommon, before allowing someone who has had, or may have had, malaria to donate, it is safer to test for malaria antibodies rather than to wait a specific length of time. Transfusion transmitted malaria is often fatal.
<i>Information</i>	This is a requirement of the Blood Safety and Quality Regulations 2005.
<i>Reason for change</i>	Entry updated to exclude donors who will only donate plasma for fractionation.
<i>Update Information</i>	This entry was last updated in: WB-DSG Edition 203 Release 71

Malaria - plasmapheresis donors

For donors who will donate whole blood, platelets and other cellular components see Malaria

<i>Excludes</i>	This entry only applies for donors who will only donate plasma for fractionation. It should not be used for donors who will donate whole blood, platelets and other cellular components.
<i>Obligatory</i>	Must not donate if: The donor has been diagnosed with malaria and the donor has not fully recovered from their illness.
<i>Discretionary</i>	In all other cases, the donor may be accepted after their return from malaria risk area if they are well.
<i>See if Relevant</i>	The ' <u>Geographical Disease Risk Index</u> ' for countries with a current endemic malaria risk.
<i>Additional Information</i>	Cases of transfusion transmitted malaria have occurred. This is mainly a problem in people who have had repeated episodes of infection with malaria. Transfusion transmitted malaria is often fatal. The processes used to fractionate plasma include several measures that inactivate or remove malarial parasites. This means that malarial risks described in the GDRl do not need to be applied for donors who will only donate plasma for fractionation. Malarial antibody testing is not required for these donors.
<i>Information</i>	This entry is compliant with the Blood Safety and Quality Regulations 2005.
<i>Reason for change</i>	This is a new entry.
<i>Update Information</i>	This entry was last updated in: WB-DSG Edition 203 Release 71

Malignancy

<i>Obligatory</i>	Must not donate.
<i>Discretionary</i>	a) If this was a non metastasized basal cell carcinoma (rodent ulcer) and local treatment is completed and all wounds are healed, accept. If any systemic medical treatment was required and has been completed at least 24 months previously, accept. b) If the potential donor has a non haematological (non-clonal) premalignant condition (e.g. polyposis coli or Barrett's oesophagus) that is being regularly monitored, or has had a similar condition cured and has been discharged from follow-up, accept. c) If the potential donor has been cured of a carcinoma in situ (CIS) and discharged from follow-up, accept. Donors who have been returned to screening following treatment for CIS can be accepted. Examples of CIS include cervical or vulval CIS, ductal CIS of the breast (DCIS), prostatic intraepithelial neoplasia (PIN) and squamous cell CIS of the skin (also known as intraepithelial squamous cell carcinoma or Bowen's disease). d) If the potential donor has had a diagnosis of melanoma in situ (including Lentigo Maligna), refer to DCSO to confirm they have not had an invasive melanoma (eg Lentigo Maligna Melanoma). Donors who have already been cleared by a DCSO can be accepted. e) Potential donors with a high risk of cancer due to family history or following genetic tests, even if had or having prophylactic surgery, or on prophylactic medication (e.g. Tamoxifen), or on routine follow up, accept.
<i>See if Relevant</i>	<u>Haematological Disease</u> <u>Surgery</u> <u>Cervical Dysplasia</u>
<i>Additional Information</i>	

Many malignancies spread through the blood stream and by invading surrounding tissues. Viruses that can be spread by blood and tissue donation can also cause some malignancies. For these reasons it is considered safer not to accept blood from people who have had a malignancy.

Basal cell carcinoma (rodent ulcer) does not spread through the blood, therefore people who have had successful treatment may donate.

The term carcinoma in situ (CIS) refers to a group of abnormal cells which have not invaded deeper tissue or spread to another part of the body. Donors who have been cured and discharged from follow up may donate. For cervical CIS, donors can be accepted if treatment is complete and any follow up smear, if performed, did not show abnormal cells. Regular screening smears are not defined as follow up.

Premalignant conditions are very common, particularly in older donors. Regular monitoring should prevent donors with invasive malignancy from being accepted. Clonal blood disorders are dealt with differently - see [Haematological Disease](#).

Melanoma in situ which has been cured by excision is not associated with a risk of metastasis. Patients with a confirmed diagnosis of melanoma in situ (ie Breslow thickness of 0 and no regression) do not require ongoing follow up beyond the initial post-operative appointment.

Lentigo Maligna is a form of melanoma in situ found on the head and neck. It should be distinguished from Lentigo Maligna Melanoma which is a true malignant melanoma.

<i>Information</i>	This is a requirement of the Blood Safety and Quality Regulations 2005.
<i>Reason for change</i>	Addition of the terms squamous cell carcinoma in situ of the skin and intraepidermal squamous cell carcinoma.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 61.

Marfan's Syndrome

<i>Obligatory</i>	Must not donate if: Has heart or blood vessel involvement.
<i>Discretionary</i>	If there is no heart or blood vessel involvement, accept.
<i>See if Relevant</i>	<u>Cardiac Surgery</u> <u>Cardiovascular Disease</u>
<i>Additional Information</i>	This a genetic disorder of connective tissues. Some individuals with Marfan's syndrome have heart and blood vessel problems that can be serious. These are screened for routinely in people who have been diagnosed with this condition. Donations should not be taken from people with heart or blood vessel problems as there may be an increased risk of serious adverse events.
<i>Reason for change</i>	The 'Obligatory' deferral has been changed from 'Cardiac involvement' to 'Heart or blood vessel involvement.'
<i>Update Information</i>	Relevant links have been added together with 'Additional Information'. This entry was last updated in: DSG-WB Edition 203, Release 01.

Meniere's Disease

<i>Discretionary</i>	If well on the day, even if on treatment to prevent attacks, accept.
<i>Additional Information</i>	Meniere's disease affects about one in a thousand people and is due to middle ear damage. Attacks can be accompanied by vertigo (a feeling of dizziness and of things spinning around). It is not thought that donation causes attacks. Providing a person with Ménière's disease is well at the time of donation, and there are no other factors that would lead to their deferral, they should be accepted.

<i>Reason for change</i>	The statement, 'even if on treatment to prevent attacks,' has been added to 'Discretionary'. 'Additional Information' has been added.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 01.

Menopause

<i>Discretionary</i>	Even if on hormone replacement therapy (HRT) or other treatment to control menopausal symptoms, accept.
<i>See if Relevant</i>	<u>Malignancy</u>
<i>Reason for change</i>	The 'Discretionary' entry has been extended to include all therapies taken to control menopausal symptoms. A link has been added to 'Malignancy' as the menopause can be secondary to treatment for cancer.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 01.

Mental Health Problems

<i>Obligatory</i>	Must not donate if: 1. Not able to fully understand and consent to the donation process and to the testing of their blood for diseases that may affect its suitability for use. 2. On treatment with sodium valproate or valproic acid.
<i>See if Relevant</i>	<u>Communication Difficulties</u> <u>Valproate and Topiramate</u>
<i>Additional Information</i>	Many people have mental health problems that are controlled with regular medication. Providing individuals are well on the day of donation and have the mental capacity to give full informed consent, there is no reason why they cannot donate whether on medication or not. Individuals who are over anxious, depressed, manic or psychotic cannot always give valid consent, or fully understand why they are being asked certain questions. Occasionally donors are on medication such as Clozapine and are under close monitoring. They should be accepted as long as they pass the Hb test.
<i>Reason for change</i>	Link updated from 'Valproate' to 'Valproate and Topiramate' in the 'See if Relevant' section.
<i>Update Information</i>	This entry was last updated in: WB-DSG Edition 203 Release 74

Migraine

<i>Obligatory</i>	Must not donate if: 1. Migraine is severe. 2. Migraine occurs more than once a week. 3. On prophylactic treatment with sodium valproate (valproic acid), topiramate or a monoclonal antibody therapy.
<i>Discretionary</i>	If the donor's migraine is not severe, and

- it occurs less than once per week, and
- any treatment does not preclude donation,

accept.

See if Relevant Headache
Monoclonal antibody therapy and other Biological Modalities
Nonsteroidal anti-inflammatory drugs
Valproate and Topiramate

Additional Information Migraine is caused by a disturbance in the normal blood flow to parts of the brain. Severe forms can be associated with focal neurological signs and symptoms, such as hemiplegia, or result in the frequency and duration of symptoms limiting normal activities.

By not accepting people with the more severe forms of migraine we hope to prevent precipitating an attack through the process of donating blood.

Care should be taken to assess any medication the donor has taken, either as prophylaxis or as relief for a migraine. Such medication may include Valproate, Topiramate or new monoclonal antibody treatments that may mean the donor is not eligible to donate.

Monoclonal antibody (MAb) therapy may be given to patients with frequent migraine where other preventative treatment has failed. MAb therapies are given by injection or by infusion. They include Eptinezumab (Vyepi[®]), Erenumab (Aimovig[®]), Fremanezumab (Ajovy[®]), Galcanezumab (Emgality[®]).

Any donor who has had migraine associated with giving blood on more than one occasion should be advised not to continue as a donor.

Reason for change The entry has been revised including the addition of an obligatory deferral for prophylaxis with topiramate or monoclonal antibody therapy. The 'See if Relevant' section has been updated.

Update Information This entry was last updated in:
WB-DSG Edition 203 Release 74

Monoclonal antibody therapy and other Biological Modalities

Includes Systemic treatment with monoclonal antibody (MAb) treatments

Obligatory **Must not donate.**

Discretionary a) If an individual monoclonal antibody treatment is listed in the entry for the underlying condition, apply guidance as per that entry.

b) If the underlying condition does not preclude donation, and it is more than 6 months from last treatment, accept.

See Is there is a specific A-Z index entry for the treatment and/or condition you are concerned about?

See if Relevant Asthma
Autoimmune Disease
Clinical Trials
Dermatitis
Eye Disease
Hypercholesterolemia
Migraine
Osteopenia
Psoriasis

Additional Information The use of monoclonal antibody and other biological therapies continues to expand. Care should be taken to understand the type of treatment and the indication, which may not be included in the 'See if Relevant' section.

Current scientific literature does not provide conclusive evidence to reject concerns that

individuals on these treatments are more prone to infection. Until further clarity is provided in the literature, donors on these medications are withdrawn, unless otherwise stated.

Reason for change Links to conditions that are currently commonly treated with monoclonal antibody therapies have been added to the 'See if Relevant' section, with reference to the expansion in types of treatment and use of these treatments within the 'Additional Information' section.

Update Information This entry was last updated in:
WB-DSG Edition 203 Release 74

Mpox (Monkeypox)

1. Affected Individuals

Obligatory **Must not donate**

Discretionary If the donor has recovered from confirmed or suspected Mpox infection and

- It is at least 28 days since the diagnosis of Mpox was made, and
- It is at least 14 days since recovery and the donor remains well, and
- It is at least 14 days since all skin lesions have healed, and
- It is more than seven days since completing any antiviral or antibiotic therapy, and
- The donor has been discharged from all follow up (including public health surveillance),

accept.

2. Contact with an individual with Mpox

Includes Individuals who have been identified by public health teams as a close contact of an individual with Mpox.

Obligatory **Must not donate**

Discretionary If it is more than 21 days since last contact and,

- the donor has no symptoms of Mpox and
- the donor has completed any isolation period, and
- the donor has been discharged from all follow-up (including surveillance by public health), and
- the donor fulfils the criteria in section 3 below regarding vaccination if applicable,

accept.

3. Immunisation for contact or risk

Excludes Individuals who have received vaccination because they work in a health care setting – see section 4 below.

Obligatory **Must not donate.**

Discretionary If the donor fulfils the criteria in section 2 above and

- it is more than four weeks since the most recent dose of a non- live or attenuated smallpox vaccination e.g. Imvanex, and
- the course of vaccination (if more than one dose) is complete,

accept.

4. Immunisation – no known contact

<i>Includes</i>	Individuals who have received vaccination because they work in a health care setting.
<i>Discretionary</i>	An individual who has received routine vaccination with Imvanex or another third-generation smallpox vaccination in an occupational setting, can be accepted provided that they are not deemed to be at risk due to an exposure episode.
<i>See if Relevant</i>	<u>Immunisation</u>
<i>Additional Information</i>	<p>Mpox was previously known as Monkeypox. In November 2022, WHO recommended Mpox as the new name for Monkeypox disease. Mpox is endemic in some African countries. During 2022 a multi-country outbreak was identified with cases in the UK, Europe, North America and other regions.</p> <p>The incubation period of Mpox is up to 21 days. The initial symptom are fever, myalgia, fatigue and headache. These symptoms are followed by a rash starting from the site of the primary infection, this rash develops into vesicles and pustule followed by scabs. Infectivity may start during initial symptoms and lasts until the rash clears and all scabs have dropped off.</p> <p>Staff should be alert for donors who report rashes and illnesses consistent with Mpox, regardless of sexual behaviour, travel history or other risk factors.</p> <p>Mpox does not spread easily between people. Human-to-human transmission occurs through contact with:</p> <ul style="list-style-type: none"> • infectious material from skin lesions • respiratory droplets in prolonged face-to-face contact • virus-contaminated objects such as bedding or clothing <p>During the 2022 multi-country outbreak, the predominance of cases among men who have sex with men and the distribution of the Mpox skin rash at presentation, suggests Mpox transmission is associated with direct contact during sex.</p> <p>Contacts may have received vaccination, to reduce the risk of serious illness. Usually vaccination will be with Imvanex or other third generation vaccine against smallpox. Contacts are eligible to donate once they satisfy the requirements of Sections 2 and 3 above.</p> <p>Health care workers may also have received vaccination to protect against Mpox in the event of possible exposure to monkeypox during their work. They will be working in accordance with Infection Prevention and Control policies and with suitable Personal Protective Equipment, which if not breached means they are eligible to donate.</p> <p>Other recipients of vaccination for Mpox must be assessed according to section 3 above.</p> <p>Imvanex is a live attenuated non-replicating third generation Smallpox vaccination. For donor selection purposes this can be assessed as a non-live vaccine but primarily donors must be assessed according to their individual risk of exposure to Mpox. The deferral of some donors for four weeks from the date of a non-live vaccination allows symptoms of Mpox from prior exposure to become evident (incubation period up to 21 days) and encompasses the time for maximum efficacy of the immunisation (up to four weeks). Donors should be deferred until completion of a course of vaccination.</p>
<i>Reason for change</i>	Inclusion of sections for donors who have received vaccination either because they are or could be a contact or because they work in a health care setting. Additional Information applicable for the whole entry contained within one section.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 67

Narcolepsy

<i>Obligatory</i>	Must not donate if: Sleep attacks or cataplexy in the previous 12 months On treatment with Modafinil (Provigil)
<i>Discretionary</i>	If free from sleep attacks or cataplexy for 12 months or more, and it is more than 7 days from the last dose of modafinil, accept.
<i>Additional Information</i>	<p>Narcolepsy is a rare neurological condition that affects the brain's ability to regulate the normal sleep-wake cycle. This can lead to symptoms such as disturbed night-time sleep, excessive daytime sleepiness and cataplexy. Consequently, narcolepsy is often thought of as a sleep disorder, but its underlying cause means that it is better classified as a disorder of the central nervous system.</p> <p>Some affected individuals may fall asleep without warning (sleep attacks) or lose muscular control that can result in falling to the ground (cataplexy). In some cases cataplexy may have the appearance of a stroke, though recovery usually occurs within minutes.</p> <p>Modafinil is a drug used to treat excessive daytime sleepiness in some patients with narcolepsy. It is associated with an increase in congenital abnormalities if taken during pregnancy. Individuals taking Modafinil are deferred to avoid the risk of components made from their donation being transfused to someone who is pregnant.</p> <p>Some individuals only have minor narcolepsy symptoms that should not interfere with donation. Sleep attacks and cataplexy may obviously cause problems during and after the donation process. However, some individuals have good control of symptoms through lifestyle adaptations and/or taking medication. If these problems are well controlled (no attacks for 12 or more months) the donor may be accepted. This includes individuals on medication to prevent attacks, with the exception of anyone taking Modafinil.</p>
<i>Reason for change</i>	A deferral for individuals treated with Modafinil has been added.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 63

National Help Lines

<i>Information</i>	<p>Donors requiring assistance who have donated, or intend to donate: In England please contact the:</p> <p>NHS Blood and Transplant (NHSBT) Telephone: 0300 123 23 23 Website: www.blood.co.uk</p> <p>In Northern Ireland, please contact the:</p> <p>Northern Ireland Blood Transfusion Service (NIBTS) Telephone: 028 9032 1414 Website: www.nibts.org</p> <p>In Scotland, please contact the:</p> <p>Scottish National Blood Transfusion Service (SNBTS) Telephone: 0345 90 90 999 Website: www.scotblood.co.uk</p> <p>In Wales, please contact the:</p> <p>Welsh Blood Service (WBS) Telephone: 0800 252266 Website: www.welsh-blood.org.uk</p>
<i>Reason for change</i>	Changes made to the National Help Lines after the move of North Wales from NHSBT to Wales
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 36.

Neurobehavioral Disorders

<i>Obligatory</i>	Must not donate if on treatment with Modafinil
<i>Discretionary</i>	If the potential donor is able to give valid consent; their disorder will not interfere with the collection process, and they have not taken Modafinil (Provigil) for 7 days, accept.
<i>See if Relevant</i>	<u>Communication Difficulties</u>
<i>Additional Information</i>	If the donor suffers from involuntary movements (tics) it is important to ensure that these will not interfere with the donation process. Donors can usually be accepted even if on medication to treat a neurobehavioral disorder. Modafinil is licensed for the treatment of narcolepsy but can be used off-licence for other disorders including ADHD. Modafinil is associated with an increase in congenital abnormalities if taken during pregnancy. Individuals taking Modafinil are deferred to avoid the risk of components made from their donation being transfused to someone who is pregnant.
<i>Reason for change</i>	A deferral has been added for donors on treatment with Modafinil.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 63

Neurofibromatosis

<i>Also Known As</i>	Von Recklinghausen's disease.
<i>Obligatory</i>	Must not donate if: History of malignant change.
<i>Discretionary</i>	Otherwise accept.
<i>See if Relevant</i>	<u>Blood Pressure - High</u> <u>Epilepsy</u> <u>Malignancy</u>
<i>Additional Information</i>	Neurofibromatosis is an inherited condition that causes tumours (swellings) on nerve tissue. These tumours are usually not cancerous but occasionally may become malignant. If they are in the brain they may cause epilepsy.
<i>Information</i>	Part of this entry is a requirement of the Blood Safety and Quality Regulations 2005.
<i>Reason for change</i>	Links have been added together with 'Additional Information'.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 01.

Neurosurgery

<i>Obligatory</i>	Must not donate.
<i>Discretionary</i>	a) If burr hole surgery only, accept. b) If carried out in the UK after 1992, providing the reason for the surgery is not itself a reason for exclusion, accept. c) If it can be shown that dura mater was not used during surgery in the UK prior to 1992 and there is no evidence of malignancy, the donor may be accepted by a ' Designated Clinical Support Officer '.

See if Relevant Cardiovascular Disease
Disabled Donor
Indwelling Shunts and Stents and Implanted Devices
Malignancy
Prion Associated Diseases
Urinary Catheterisation
Surgery

Additional Information Dura mater has led to the spread of prion related diseases (CJD). It should not have been used in the UK after 1992. The situation in other countries varied so specific dates cannot be given.

Information Part of this entry is a requirement of the Blood Safety and Quality Regulations 2005.

Reason for change The See if Relevant section has been revised.

Update Information This entry was last updated in:
 DSG-WB Edition 203, Release 61.

Night Sweats

Obligatory **Must not donate if:**
 Unexplained.

Discretionary If due to the menopause, accept.

See if Relevant Infection - General
Malignancy
Menopause

Additional Information Unexplained night sweats may be an indication of an undiagnosed infection or malignancy. Both would be a reason to defer a potential donor.

Reason for change Links and 'Additional Information' have been added.

Update Information This entry was last updated in:
 DSG-WB Edition 203, Release 01.

Non-Consented Exposure to Human Body Fluids

Definitions **Non-Consented Exposure to Human Body Fluids:**
 A non-consented injury or assault in which an individual is exposed to potentially infective material that could be transferred through donation. The causes may range from a sharps injury to bites, punches and abrasions or sexual assault where mucous membranes have been contaminated with human blood or other body fluids. It also applies to any inoculation injury with abnormal prions from any species.

1. Affected Individual

Obligatory **Must not donate if:**

- a) The incident involved any material containing abnormal prions.
- b) Less than four months after the date of an inoculation injury or contamination of mucosa or non-intact skin with human blood or body fluids.
- c) Under ongoing investigations following exposure.

See if Relevant Animal Bite (Non-Human)
Blood Safety Entry
Hepatitis
HIV
HTLV
Prion Associated Diseases
Xenotransplantation

<i>Additional Information</i>	<p>Prion related diseases can be symptom free for many years. During the incubation stage, infection may be passed on by donated material.</p> <p>Human blood or body fluids may be contaminated with infective material that may be passed on by donated material. Waiting four months (if a validated test for HBV, HCV HIV NAT is negative) helps to ensure that any infection is not passed on. This includes donors where the contact has Hepatitis B infection or is a recipient of blood derived coagulation factor concentrates.</p> <p>If an individual is undergoing further tests or follow up following an exposure, donation should be deferred until all follow up is complete and above criteria apply.</p> <p>This guidance presumes that a validated NAT test for hepatitis C is negative. If this test is stopped, the guidance will change.</p>
<i>Information</i>	This is a requirement of the Blood Safety and Quality Regulations 2005.

2. Current or Former Sexual Partner of Affected Individual

<i>Obligatory</i>	<p>Must not donate if: The donor's sexual partner is being monitored for evidence of transmitted infection following exposure to a known infected individual.</p>
<i>Discretionary</i>	<p>a) If the partner has not been exposed to known infective material, accept.</p> <p>b) If the partner was exposed to known infective material and has been told that they are not infected and no longer require to be monitored, accept.</p> <p>c) If a former sexual partner and it is more than three months since the last sexual contact, accept.</p>
<i>See if Relevant</i>	<p><u>Hepatitis</u> <u>HIV</u> <u>HTLV</u> <u>Prion Associated Diseases</u></p>
<i>Reason for change</i>	The wording of this entry has been revised to improve clarity.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 57

Non-Contagious Diseases - Contact With

<i>Definitions</i>	<p>Non Contagious Disease: This is a disease which is not easily transmitted from person to person. It may be a zoonotic infection or it may result from a shared environmental issue e.g. food poisoning or a common travel history.</p> <p>Zoonotic infection: The WHO defines this as any disease or infection that is naturally transmissible from vertebrate animals to humans.</p>
<i>Obligatory</i>	<p>Refer to the specific entry for the disease. If there is no specific entry, and/or no instruction for contacts, use the guidance below.</p> <p>Must not donate if:</p> <p>a) Diagnosed with, or showing symptoms of, the disease.</p> <p>b) Under investigation or monitoring for potential infection with the disease. This includes any deferral period recommended by Public Health.</p> <p>c) The donor has been exposed to the same circumstances that led to infection in another human. This might include a common travel history or, in the case of food poisoning, to the consumption of the same food.</p>

<i>Discretionary</i>	If the donor: <ul style="list-style-type: none"> • Is well and has no symptoms of infection, and • Is not being investigated or monitored, and • Is not considered to be at high risk of infection, accept.
<i>See if Relevant</i>	If there is a shared risk for a transfusion transmissible infection, see the index entry for that condition.
<i>Additional Information</i>	<p>Many infections are not easily spread from one person to another. In other situations, the infection with which there has been contact will not represent a risk to recipients and staff. Donors reporting such contact can normally be accepted but the nature of the contact should be assessed. For instance, has the potential donor had the same risk factors as an infected individual e.g. travelled to the same place or eaten the same food. If they have, the risk of the potential donor also being infected must be considered together with any appropriate deferral.</p> <p>For some zoonotic infections, people in contact with affected animals may be followed up by health protection teams. This can include a period of monitoring for potential transmission of the infection. Individuals in this situation should not be accepted to donate until further review or testing is no longer required, and any deferral period stipulated by the health protection team has elapsed.</p> <p>If in doubt: Contact a 'Designated Clinical Support Officer'.</p>
<i>Reason for change</i>	Guidance and information has been added for donors under investigation or monitoring following contact with a zoonotic infection.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 61.

Nonsteroidal Anti-Inflammatory Drugs

<i>Also Known As</i>	NSAIDs or nonsteroidals.
<i>Obligatory</i>	<p>Assess the reason for treatment and see if there is a relevant index entry.</p> <p>1. Must not donate if: Taken for a serious long-term illness including cardiovascular disease.</p> <p>2. Platelets: If the donor has taken a NSAID drug in the 48 hours before attending, their donation must not be used for the preparation of platelets and other blood components intended to treat thrombocytopenia and/or platelet dysfunction.</p>
<i>Discretionary</i>	If medication is self-prescribed, the donation will not be used to prepare platelets and other blood components intended to treat thrombocytopenia and/or platelet dysfunction, and the donor meets all other criteria, accept.
<i>See if Relevant</i>	<u>Cardiovascular Disease</u> <u>Drug Index</u> - preparations which may affect platelet function
<i>Additional Information</i>	<p>Nonsteroidal anti-inflammatory drugs can stop platelets (small fragments of cells that help control bleeding) from working properly. Platelets may be manufactured from whole blood or component (apheresis) donations. Blood Services may produce other components that included functional platelets, including some forms of whole blood. As these are used to control or prevent bleeding in patients, it is essential that they do not include platelets affected by nonsteroidal anti-inflammatory drugs.</p> <p>Taking these drugs will not affect the use of a donation for red cell transfusion (the commonest use) but, the reason they are being taken, might.</p>
<i>Reason for change</i>	The entry has been revised to add reference to other blood components that contain active platelets.

Update Information This entry was last updated in:
WB-DSG Edition 203 Release 72

Nose Bleeds

Discretionary If the potential donor passes the haemoglobin screening test, accept.

See if Relevant Anaemia - Discretionary 1. Iron deficiency
Bleeding Disorder
Transfusion

Additional Information Severe or regular nose bleeds lead to a loss of iron from the body and this can cause iron deficiency anaemia. Donating blood also causes the body to lose a substantial amount of iron. The combination of the two will make anaemia much more likely.

Reason for change Links to 'Anaemia - Discretionary 1. Iron deficiency' and 'Transfusion' have been added together with 'Additional Information'.

Update Information This entry was last updated in:
DSG-WB Edition 203, Release 01.

Oseltamivir

Also Known As Tamiflu®.

Obligatory **Must not donate if:**
a) Taking oseltamivir (Tamiflu®) as treatment for influenza.

b) At any time in the seven days prior to, or while taking oseltamivir, the donor has had symptoms of influenza, (a temperature of more than 38 degrees centigrade, or a history of fever and two or more of the following symptoms: cough, headache, runny nose, diarrhoea or vomiting).

Discretionary If the potential donor is taking oseltamivir as prophylaxis, they have not been advised to be confined to home and have not had any symptoms of influenza, accept.

See if Relevant Infection - Acute

Additional Information Oseltamivir is a viral neuraminidase inhibitor (neuraminidase is an enzyme that helps the virus spread from cell to cell). It is used to treat influenza and for post-exposure prophylaxis of influenza. It appears to be a very safe drug with little evidence for teratogenic (potential to cause birth defects) or mutagenic (potential to cause malignancy) effect.

Information Part of this entry is a requirement of the Blood Safety and Quality Regulations 2005.

Update Information This entry was last updated in:
DSG-WB Edition 202, Release 13.

Osteoarthritis

Obligatory Must not donate if it is less than four months from any invasive treatment or investigation including but not limited to joint injections.

Discretionary If the donor has been treated by joint injection, and

- treatment was performed within the NHS, or treatment was performed by Qualified Health Care Professional (as defined in the Complementary Therapy entry), and
- if it is at least 7 days from any joint injection with steroid, and
- it is at least 4 months from any joint injection with autologous platelet rich plasma (PRP)

accept.

<i>See if Relevant</i>	<u>Complementary Therapy</u> (for the approved list of Qualified Health Care Professionals) <u>Disabled Donor</u> <u>Drug Index - Drugs and Platelet Donations</u> <u>Endoscopy</u> <u>Nonsteroidal Anti-Inflammatory Drugs</u> <u>Steroid Therapy</u>
<i>Additional Information</i>	Donors who have severe pain or disability that makes it difficult for them to get on or off the bleed bed should not donate. Medicines taken for arthritis may affect platelet function. This can be checked in the <u>Drug Index</u> . Some individuals may undergo arthroscopy as part of the assessment and investigation of their condition; this is usually performed by a rigid endoscope (see <u>Endoscopy</u>). Some individuals may receive joint injections to relieve the symptoms of osteoarthritis. These may be intra-articular (into the joint) or periarticular (around the joint). Treatments may include steroids, hyaluronic acid or platelet rich plasma. Individuals who have received autologous platelet rich plasma must wait at least four months before donation. If the donor is uncertain about the treatment that was given, refer to a DCSO.
<i>Reason for change</i>	Guidance on the acceptance of donors treated with joint injection has been added. A link to Endoscopy has been added.
<i>Update Information</i>	This entry was last updated in: WB-DSG Edition 203 Release 72

Osteomyelitis

<i>Obligatory</i>	Must not donate if: Less than 24 months from completing treatment and cure.
<i>Additional Information</i>	Sometimes it is difficult to be certain that all infection has been eliminated. Waiting 24 months minimizes the risk of any infection being passed on through donation.
<i>Information</i>	This is a requirement of the Blood Safety and Quality Regulations 2005.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 202, Release 02.

Osteopenia

<i>Obligatory</i>	Must not donate if: The donor is being treated with systemic monoclonal antibody therapy e.g. Denosumab (Prolia®). See <u>Monoclonal antibody therapy and other Biological Modalities</u> .
<i>Discretionary</i>	If the cause is not of itself a reason to defer, even if on treatment to prevent or treat (other than Denosumab), accept.
<i>See if Relevant</i>	<u>Disabled Donor</u> <u>Malignancy</u> <u>Monoclonal antibody therapy and other Biological Modalities</u> <u>Steroid Therapy</u> <u>Vitamins and Other Nutritional Supplements</u>
<i>Additional Information</i>	Osteopenia occurs when there is decreased mineralization (mainly lack of calcium) of bone. It can occur for many reasons so it is important to ensure that it is not associated with a condition that would require a potential donor to be deferred.
<i>Reason for change</i>	The See if Relevant section has been updated.

Update Information This entry was last updated in:
WB-DSG Edition 203 Release 71

Paget's Disease of Bone

Also Known As Osteitis deformans.

Discretionary Even if on medication with painkillers or bisphosphonates, accept.

See if Relevant Disabled Donor
Drug Index - preparations which may affect platelet function
Nonsteroidal Anti-Inflammatory Drugs
Surgery

Additional Information Paget's disease of bone is very common in the UK, affecting about one in 20 adults aged over 50 years. The cause is not known. Many people with the condition have no symptoms and so will be accepted by the blood and tissue services. There is no evidence that it is spread by donation. It is most commonly treated with painkillers and bisphosphonates. The use of these drugs is accepted for other conditions so there seems no reason why individuals with Paget's disease of bone on treatment should not be accepted, provided that they are otherwise fit to donate.

Reason for change Links have been added to 'Drug Index' and 'Surgery'.

Update Information This entry was last updated in:
DSG-WB Edition 203, Release 01.

Pain Killers

Obligatory Assess the reason for treatment and see any relevant Index entry.

Discretionary If the donor is otherwise fit to donate, regardless of the type of medication, accept.

See if Relevant Disabled Donor
Drug Index - preparations which may affect platelet function
Indwelling Shunts and Stents and Implanted Devices
Nonsteroidal Anti-Inflammatory Drugs

Reason for change The entry has been revised to improve clarity.

Update Information This entry was last updated in:
DSG-WB Edition 203, Release 01.

Palpitations

Obligatory **Must not donate if:**

1. The donor has a history of palpitations.
2. A significant arrhythmia or other cardiac cause has been confirmed or is suspected as the cause of palpitations.

Discretionary

1. Donors with a history of palpitations where:
 - The donor has consulted their GP and a cardiac cause has been excluded, and
 - The donor is not undergoing any investigations, and,
 - The donor is symptom-free on the day of attendance;
Accept, even if on beta blockers to prevent symptoms.

2. Donors with a history of palpitations where a cardiac cause has not been excluded, refer to Arrhythmia entry.

See if Relevant Arrhythmias

Additional Information Donors with ongoing symptoms of palpitations are at risk of a donor adverse event during or after donation.

Many donors will have experienced palpitations at some time in their lives. Symptoms are often associated with anxiety or stress. As long as they are symptom-free on the day of donation and investigations have ruled out a cardiac cause, donors can be accepted, even if on treatment to prevent symptoms.

Reason for change New entry

Update Information This entry was last updated in:
DSG-WB Edition 202, Release 44

Parvovirus B19

Includes Human parvovirus; slapped cheek syndrome; erythema infectiosum; fifth disease

1. Affected Individual

Obligatory **Must not donate.**

Discretionary If it is more than 4 weeks from:

- recovery from systemic symptoms (low-grade fever, malaise, headache, nasal discharge, abdominal pain, sore throat), and
- a positive result for parvovirus B19 DNA, if testing has been carried out,

accept.

2. Close Contact with an Affected Individual

Obligatory **Must not donate if:**
Less than 3 weeks since contact with an individual with suspected or confirmed parvovirus B19 infection. This includes individuals diagnosed through blood donation screening.

- Discretionary*
- a) If the donor has a definite past history of parvovirus B19 infection, accept.
 - b) If more than 3 weeks since last contact with an individual with parvovirus B19 infection, accept.
 - c) If any contact in the previous 3 weeks has only been with someone in the post-infectious phase, i.e. after the rash has appeared, accept.

See if Relevant Infectious Diseases – Contact With

Additional Information Parvovirus B19 is a viral infection which occurs most commonly in children. The virus can be transmitted through droplet spread (respiratory), or from mother to baby, or via blood transfusion. Outbreaks of parvovirus B19 occur every 3 to 4 years during the late winter / early spring period. Following exposure, the incubation period is 14 to 21 days.

Infection may be asymptomatic or may result in a mild self-limiting illness presenting with fever, malaise, upper respiratory tract symptoms and abdominal pain. Children often develop a bright red facial rash ('slapped cheek') one to two weeks later, followed after a few days by a light pink rash on the chest, stomach, arms and thighs. Adults have similar symptoms but are less likely to have a facial rash. Adults are also more likely to have joint pain (polyarthropathy) especially in the hands, wrists, feet and ankles. Rashes and joint pain

can persist for several weeks or months after someone has recovered from the infection and do not prevent donation.

Individuals with parvovirus B19 are infectious at the early stage, before and while experiencing systemic and upper respiratory tract symptoms. Once a rash has appeared an individual is no longer infectious. People who have recovered from parvovirus B19 have lifelong immunity to the virus and cannot be reinfected.

Parvovirus B19 may cause more serious illnesses, including anaemia and bone marrow failure, in non-immune individuals who are also immunocompromised. If acquired during pregnancy, parvovirus B19 may result in severe fetal anaemia (hydrops fetalis), miscarriage and stillbirth.

<i>Reason for change</i>	This is a new entry.
<i>Update Information</i>	This entry was last updated in: WB-DSG Edition 203 Release 70

Peptic Ulcer

<i>Obligatory</i>	Must not donate if: a) Recent symptoms or on active treatment to heal an ulcer. b) Associated with malignant change.
<i>Discretionary</i>	If not affected by a) or b) above, even if on maintenance treatment, accept.
<i>See if Relevant</i>	<u>Anaemia - Discretionary 1. Iron deficiency Surgery Transfusion</u>
<i>Additional Information</i>	Bleeding is a common problem associated with peptic ulcers. This can be profuse and may require transfusion, or gradual, leading to iron deficiency. Taking blood from a person at risk of bleeding will reduce their ability to compensate for blood loss and may lead to treatment that would not otherwise have been needed.
<i>Reason for change</i>	Total gastrectomy is no longer a reason for withdrawal. Additional links have been added together with 'Additional Information'.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 01.

Periods

<i>Obligatory</i>	Must not donate if: a) Period has been missed. b) If under investigation for heavy and prolonged periods. c) Has uncontrolled period pain.
<i>Discretionary</i>	a) If a period has been missed, pregnancy can be excluded and the donor is well, accept. b) If the potential donor is taking supplemental iron to prevent anaemia, is not under investigation for heavy or prolonged periods and understands that donation will make anaemia more likely, accept. c) If taking medication to decrease blood loss (e.g. etamsylate (Dicycne®), tranexamic acid (Cyklokapron®)), accept.
<i>See if Relevant</i>	<u>Anaemia - Discretionary 1. Iron deficiency Drug Index - preparations which may affect platelet function Nonsteroidal Anti-Inflammatory Drugs</u>

Pregnancy
Surgery

<i>Additional Information</i>	It is OK to donate while having a period. However, the combination of blood loss from periods and donation will make iron deficiency anaemia more likely, particularly if the periods are heavy or prolonged. This affect can be minimised by taking supplemental iron.
	If the donor feels unwell because of their period, they should not donate but if period pain is well controlled by medication, they may be accepted. It is important that the type of medication taken, and its affect on platelet function is noted.
<i>Reason for change</i>	The entry has been rewritten for greater clarity.
	A 'Discretionary' entry has been added for donors taking drugs to reduce blood loss.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 01.

Perthes' Disease

<i>Discretionary</i>	Accept.
<i>See if Relevant</i>	<u>Drug Index - preparations which may affect platelet function</u> <u>Nonsteroidal Anti-Inflammatory Drugs</u> <u>Surgery</u>
<i>Additional Information</i>	Perthes' disease affects about one in 10,000 children between the ages of two and 15 years. It causes damage to the femoral head and will usually heal with conservative treatment. Surgery may be required and there is the possibility of chronic arthritis. This may require treatment with pain killers that might affect platelet function.
<i>Reason for change</i>	Relevant links and 'Additional Information' have been added.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 01.

Pituitary Disorders

<i>Obligatory</i>	<p>Must not donate if:</p> <ul style="list-style-type: none"> a) Malignant tumour. b) Part of a multiple endocrine neoplasia (MEN) syndrome. c) Has had open neurosurgery. d) Ever received injections of human pituitary extract. e) Acromegaly or growth hormone excess. f) Has adrenal failure, or requires treatment with oral steroids. g) Has cranial diabetes insipidus, or syndrome of inappropriate anti-diuretic hormone (SIADH), even if on treatment. h) Currently on injectable drug treatment e.g. pegvisomant (Somavert®).
<i>Discretionary</i>	<ul style="list-style-type: none"> a) If a non secretory or prolactin secreting pituitary tumour (adenoma) has been confirmed as non-malignant and the donor has no symptoms, even if on oral medication and/or underwent neurosurgery in the UK after 1992, accept. b) If open neurosurgery carried out in the UK after 1992, providing the reason for the surgery is not itself a reason for exclusion, accept. c) If it can be shown that dura mater was not used during open neurosurgery in the UK prior to 1992 and there is no evidence of malignancy, the donor may be accepted by a 'Designated Clinical Support Officer'. d) If treated, exclusively with recombinant-derived growth hormone, accept. (In the UK this has been since 1986).

e) If the donor has undergone trans-sphenoidal surgery for a pituitary tumour, all wounds are healed, accept.

f) If the donor has undergone radiation therapy for a benign tumour, even if on long term follow up, provided there are no complications related to either the treatment received or to the underlying condition, accept.

See if Relevant Central Nervous System Disease
Epilepsy
Growth Hormone
Neurosurgery
Prion Associated Diseases
Surgery

Additional Information Pituitary adenomas are quite common and the majority are benign, i.e. not able of spreading to other parts of the body (metastasizing). Two-thirds of pituitary adenomas remain completely confined to the pituitary gland and approximately one-third will expand into tissues in the immediate vicinity of the gland. Less than 1% of pituitary tumours are malignant.

Pituitary adenomas may be non-secretory (25%) or secrete hormones such as prolactin (30%), growth hormone (10-15%, leading to acromegaly), ACTH (leading to Cushing's disease), TSH (leading to thyroid dysfunction) or LH/FSH (leading to fertility problems).

Acromegaly, caused by growth hormone over secretion, is associated with an increased risk of cardiovascular complications, including cardiomyopathy, increase in left ventricular mass, arrhythmias and hypertriglyceridaemia.

Hypopituitarism, with a reduction in levels of one or more pituitary hormones, can result from either the underlying pituitary condition or its medical/surgical management. A deficiency of ACTH may result in adrenal failure. Pituitary hormones are replaced through medication as required.

Patients with posterior pituitary lesions may develop diabetes insipidus or hypothalamic problems, which require careful fluid balance. Donating a unit of blood may compromise this balance.

Sheehan's syndrome is post-partum (after the birth of a baby) pituitary necrosis. It is caused by hypovolaemia from post-partum blood loss. It is likely that the patient will have been transfused.

Reason for change This is a new entry.

Update Information This entry was last updated in:
DSG-WB Edition 203, Release 01.

Platelet Count

Obligatory **All donors:
Must not donate if:**
a) Under investigation for an abnormal platelet count.

b) The platelet count is known to be less than $150 \times 10^9/L$.

c) The platelet count is known to be more than $450 \times 10^9/L$.

d) Any underlying cause precludes donation.

**In addition, for Platelet Component Donors only:
Must not donate if:**

The predicted post-donation platelet count is less than $100 \times 10^9/L$.

Discretionary a) If a donor has been investigated for an abnormal platelet count and

- no underlying cause has been identified that would lead to deferral, and
- the donor does not require any monitoring or follow up,

accept

b) If testing by the blood transfusion service finds a donor to have a platelet count which is outside the normal range, the donor can be accepted if their results comply with local

policies and procedures. Blood transfusion services should have a written policy for management of donors who are found to have a platelet count of less than $150 \times 10^9/L$ or more than $450 \times 10^9/L$ during donation testing.

See if Relevant Haematological Disease
Immune Thrombocytopenia
Platelet Disorder
Thrombosis and Thrombophilia

Additional Information Taking a platelet donation from a donor with a platelet count lower than $150 \times 10^9/L$ is unlikely to provide a therapeutic dose.

Platelet counts outside of the normal range (i.e. less than $150 \times 10^9/L$ or more than $450 \times 10^9/L$) may be due to an underlying disease process. High platelet counts can also be associated with iron deficiency. Transfusion services should ensure that, where abnormal platelet counts are identified as part of routine donation testing, these are reviewed and managed appropriately. Further investigation may be required for donors with persistently abnormal results.

Information Part of this entry is a requirement of the Blood Safety and Quality Regulations 2005.

Reason for change Additions to the Obligatory and See if Relevant sections.

Update Information This entry was last updated in:
WB-DSG Edition 203 Release 73

Platelet Disorders

Includes Individuals with, or carriers of, an inherited platelet disorder. These include:

- Bernard-Soulier disease
- Glanzmann's thrombasthenia
- Hermansky-Pudlak syndrome
- Jacobsen syndrome
- Lowe syndrome
- Paris-Trousseau syndrome
- Platelet release and storage pool defects
- Thrombocytopenia with absent radius syndrome

Acquired platelet disorders due to an underlying condition.

Excludes Donors taking medications that reduce platelet function e.g. Aspirin, Clopidogrel – see:

- Nonsteroidal Anti-Inflammatory Drugs
- Clopidogrel
- Cardiovascular Disease
- Other entries relevant to reason for treatment

Donors identified as having an abnormal platelet count on testing by the blood transfusion service – see Platelet Count.

Thrombotic thrombocytopenic purpura (TTP) – see Thrombosis and Thrombophilia.

Obligatory **Must not donate if:**

- a) The donor has an inherited platelet disorder.
- b) The donor is an affected carrier of an inherited platelet disorder.
- c) The donor has an acquired platelet disorder due to an underlying condition.
- d) The donor had an acquired platelet disorder and the underlying cause precludes donation e.g. malignancy.
- e) There is a history of excessive bleeding or bruising.

- f) The donor is requiring monitoring and/or follow up.
- g) There is any associated immune system or organ involvement e.g. heart, lung, kidney, that precludes donation.
- h) The donor has received a transfusion or plasma exchange since 1st January 1980.
- i) The donor has a platelet disorder and has ever:
- received blood derived and recombinant products e.g. Factor VIIa, ADAMTS13 and/or
 - received or is currently on treatment to reduce or prevent excessive bleeding e.g. desmopressin (DDAVP[®]), tranexamic acid, oral contraceptive pill and similar hormone therapies, intrauterine device (IUD) and/or
 - needed iron supplementation.
- j) The donor has required or been advised they will require prophylactic treatment for surgery, dental treatment, or for any other procedure.
- k) The donor has required or been advised they should receive immunisations subcutaneously rather than intramuscularly.

- Discretionary*
1. If the donor is a non-affected carrier of an inherited platelet disorder and fulfils all other criteria, refer to DCSO.
 2. If the donor had an acquired platelet disorder (except immune thrombocytopenia/ITP – see Immune Thrombocytopenia) that has now fully resolved, the underlying condition does not preclude donation and they fulfil all other criteria, refer to DCSO.

See if Relevant

Cardiovascular Disease
Clopidogrel
Haematological Disease
Immune Thrombocytopenia
Nonsteroidal Anti-Inflammatory Drugs
Platelet Count
Transfusion
Thrombosis and Thrombophilia

Additional Information

Platelet disorders can be inherited or can be acquired, e.g. due to an autoimmune reaction or malignancy such as leukaemia. Symptoms are similar whether there are too many platelets in circulation, too few platelets in circulation or the correct number of platelets are in circulation but they do not work properly.

These disorders can cause significant bleeding, either spontaneously or in response to even minimal trauma or minor procedures. Nose bleeds, bleeding from the gums and petechiae are common. Bleeding can lead to iron deficiency anaemia.

Individuals will have been assessed and advised about their condition and bleeding risk. Most would be at increased risk of bruising and other complications from blood donation so affected individuals must not be accepted.

The diagnosis of the milder forms or carrier status of platelet disorders may arise from family screening, or through testing during investigation for menorrhagia (heavy periods), or bleeding during pregnancy or childbirth. Some people with the carrier state have symptoms which may or may not need treatment and/or are at risk of bleeding and therefore would be at increased risk of bruising and other complications from donation so affected carriers must not be accepted. Affected individuals and affected carriers may have been provided with a Bleeding Disorders Information Card.

Carriers who have been diagnosed through family screening, have not had any symptoms and have had their platelet function investigated and demonstrated to be normal may be able to donate once this information has been confirmed with the donor's specialist or GP by a DCSO.

Treatments include plasma products, plasma exchange or more rarely platelet transfusions. Curative treatment with stem cell or bone marrow transplant is an option for the most severe conditions.

The inherited platelet disorders are often part of a multisystem condition, but carriers are less likely to have significant organ involvement.

The guidance contained in this entry is not intended for use for donors without a platelet disorder, for example for someone who may have taken tranexamic acid for heavy periods

due to an underlying gynaecological cause.

For donors who are identified as having an abnormal platelet count following testing by the blood transfusion service, refer to the [Platelet Count](#) entry.

Reason for change Clarification of the scope of this entry. Expansion of obligatory and discretionary criteria. Addition of relevant links. Additional Information section rewritten to support revised entry.

Update Information This entry was last updated in:
WB-DSG Edition 203 Release 73

Pneumothorax

1. Spontaneous

Obligatory **Must not donate if:**
a) Not recovered.

b) Associated with cystic fibrosis.

c) Associated with emphysema.

See if Relevant [Asthma](#)
[Infection - General](#)
[Respiratory Disease](#)
[Surgery](#)
[Tuberculosis](#)

Additional Information Spontaneous pneumothorax most often affects tall thin men around the ages of 20 to 40 years. It also affects people with lung disease such as cystic fibrosis, emphysema and tuberculosis. It reduces lung function and so can decrease the amount of oxygen entering the blood. Removing blood from an affected person may worsen or cause breathing problems.

Update Information This entry was last updated in:
DSG-WB Edition 203, Release 01.

Reason for change Cystic fibrosis has been added as a reason not to donate.

Relevant links have been added together with 'Additional Information'.

2. Traumatic

See [Accident](#)

Update Information This entry was last updated in:
DSG-WB Edition 202, Release 02.

Update Information This entry was last updated in:
DSG-WB Edition 203, Release 01.

Update Information This entry was last updated in:
DSG-WB Edition 202, Release 02.

Polycythaemia and Raised Haemoglobin

Obligatory **Must not donate.**

Discretionary If specialist investigation has excluded Polycythaemia Rubra Vera, or another myeloproliferative neoplasm, and no treatment or further investigation is planned, the donor can be accepted for whole blood donation or for double red cell donation. Donors with a haemoglobin above the normal range should not usually be accepted for plasma or platelet donation.

See if Relevant

Cardiovascular Disease
Haematological Disease
Haemoglobin Disorders
Haemoglobin Estimation
Respiratory Disease

Additional Information In men, haemoglobin concentrations in excess of 180 g/l or red cell counts in excess of 6.5×10^{12} and in women, haemoglobin concentrations in excess of 165 g/l or red cell counts in excess of 5.6×10^{12} should be repeated. If found to be persistently raised the donor should not be accepted and referred for investigation.

Polycythaemia is commonly linked to malignant or pre-malignant conditions or to the body's response to a shortage of oxygen. Apparent polycythaemia is caused by a decreased plasma volume. All of these are reasons not to accept a donation, either because of the association with malignancy, or because of the potential to harm the donor.

Individuals with 'high affinity' haemoglobins can develop polycythaemia because of the reduced oxygen carrying capacity of their blood. This would be detrimental to a recipient of their blood and donation may be harmful to the donor. For these reasons they should not be accepted

Reason for change Clarification of the suitable donation types for donors with a haemoglobin above the normal range has been added.

Update Information This entry was last updated in:
DSG-WB Edition 203, Release 58

Porphyria

Obligatory **Must not donate if:**
Suffers from porphyria

Discretionary If the potential donor suffers from Acute Porphyria, Acute Intermittent Porphyria (AIP), Varigate Porphyria (VP) or Hereditary Coproporphyrinuria (HCP), it is 12 months or more since their last acute attack and they have no current skin lesions, accept.

See if Relevant Hepatitis

Additional Information Acute porphyrias (AIP, VP and HCP) may be associated with skin lesions and raised blood porphyrins independently of acute attacks. Theoretically the recipient of the blood could develop skin lesions, and we therefore exclude anyone with active skin lesions.

Porphyria Cutanea Tarda (PCT) is almost always an acquired condition associated with underlying liver disease, usually hepatitis of viral or unknown origin. These patients are often treated by venesection, however because of the risk of transmission of the agent that caused the condition the blood is not suitable for transfusion.

With Erythropoietic Protoporphyrinuria (EPP) and Congenital Erythropoietic Porphyria (CEP) the patient is often anaemic because of the condition. Also in these conditions there are porphyrins in the red cells and red cell life span is reduced so the blood is not suitable for donation.

Reason for change This is a new entry.

Update Information This entry was last added in:
DSG-WB Edition 203, Release 05.

Pre- and Post-Exposure Prophylaxis for HIV prevention

Obligatory **Must not donate if:**

Donor has taken Pre-Exposure Prophylaxis (PrEP) or Post-Exposure Prophylaxis (PEP) in the previous three months.

Assess any donor using PrEP or PEP for blood safety risks relating to sexual activity.

Discretionary

If it is over three months since the donor has taken PrEP or PEP and there is no other blood safety risk, accept

See if Relevant Blood Safety Entry
HIV
Non-consented Exposure to Human Body Fluids

Additional Information The use of Pre-Exposure Prophylaxis (PrEP), e.g. Truvada[®], to prevent HIV is increasing. Individuals taking PrEP are unlikely to be eligible to donate due to criteria within the blood safety entry. However, PrEP is also available via private prescription and/or online pharmacies and may be used by individuals who would not otherwise be deferred.

Use of PrEP may interfere with testing for HIV by delaying seroconversion or giving unclear results in a positive donor. For this reason, it is important that donors who have taken PrEP in the previous three months are not accepted to donate, even if they do not have another blood safety risk.

Post-Exposure Prophylaxis (PEP) has a similar mechanism of action to PrEP and may also interfere with testing results. In the UK PEP is prescribed to people who have been exposed to someone who may have HIV. This includes sexual activity or a needle stick injury. Donors who have received PEP will usually be ineligible to donate for the same reason they were given PEP.

If the underlying reason for taking PrEP or PEP warrants a longer deferral period, this should be applied.

This guidance will be updated as more information about the use and effects of PrEP/PEP becomes available.

Reason for change This entry was revised to support the implementation of recommendations from the FAIR study.

Update Information This entry was last updated in:
DSG-WB Edition 203, Release 57

Pregnancy

Obligatory **Must not donate if:**

- a) Pregnant.
- b) Less than 6 months have passed since delivery or termination.
- c) Resulted in a malignant (invasive) hydatidiform mole.
- d) Resulted in a non-malignant (non-invasive) hydatidiform mole and treatment and follow up is ongoing.

Discretionary If the pregnancy ended before the 12th week of pregnancy without significant blood loss, if follow up is complete and it is more than 7 days from last dose of methotrexate (if taken), and it is agreed by a Physician member of the designated clinical support, accept.

See if Relevant Anaemia - Discretionary 1. Iron deficiency
Malignancy
Surgery
Transfusion
Trying to Conceive

Additional Information During pregnancy, particularly in the later part, a woman loses a considerable amount of iron to the baby. It is important to allow time for this lost iron to be replaced through the mother's diet. Donating during pregnancy will make it very likely that the pregnant woman will become short of iron and this may lead to anaemia and even threaten the pregnancy. Iron usage in pregnancy occurs mostly between 12 and 35 weeks either to increase the number of red cell of the mother, or for the growth of the baby (after 30 weeks). Pregnancies of less than 12 weeks have little impact on the mother's iron stores. However if there was significant bleeding due to a miscarriage or ectopic pregnancy a full 6 months from the date of this event is advisable before the lady donates.

Methotrexate is now increasingly used to medically treat ectopic pregnancy, to avoid surgery and protect the fallopian tube. This method of treatment, if successful, is not associated with significant bleeding but a week is needed for any residual methotrexate to clear the system.

A mother can donate if she is still breast-feeding, provided that a longer period than 6 months from delivery has passed.

If a woman is trying to become pregnant they can donate if they have not missed a period and are not under investigation or on infertility treatment. If they are on treatment or under investigation for infertility see the link for 'Trying to Conceive'.

Hydatidiform moles may be malignant. If they are, the woman will not be able to donate. In other cases it is important for treatment and follow up to be completed so that the possibility of malignancy is excluded.

Repeat anti-HLA, anti-HNA and/or anti-HPA antibody testing should be undertaken when donors return after pregnancy, regardless of duration, when:

- it is intended to collect components for which the blood service has implemented TRALI risk reduction measures based on antibody testing,

or,

- it is intended to collect HPA-matched components

Reason for change Advice to consider repeat HLA, HNA or HPA antibody testing after pregnancy has been added.

Update Information This entry was last updated in:
DSG-WB Edition 203, Release 43.

Prion Associated Diseases

Includes Familial, sporadic and variant Creutzfeldt-Jacob disease (CJD), Gerstmann-Sträussler-Scheinker disease and fatal familial insomnia.

Obligatory **Must not donate if:**

1. Diagnosed with any form of CJD, or other human prion disease.
2. Identified at increased risk of developing a prion associated disorder.
This includes:
 - a) Individuals at familial risk of prion-associated diseases (have had two or more blood relatives develop a prion-associated disease or have been informed following genetic counselling they are at risk).
 - b) Individuals who have been told that they have been put at increased risk from surgery, transfusion or transplant of tissues or organs.
 - c) Individuals who have been told that they may be at increased risk because a recipient of their blood or tissues has developed a prion related disorder.
 - d) Recipients of dura mater grafts.
 - e) Recipients of corneal, scleral or other ocular tissue grafts.
 - f) Recipients of human pituitary derived extracts.
 - g) **Since January 1st 1980:**
Recipients of a transfusion or allogeneic human tissue (except recipients of donated human eggs, sperm or embryos).

Discretionary If the donor has had two or more blood relatives develop a prion-associated disease and, following genetic counselling, they have been informed that they are not at risk, accept. This requires confirmation by a **'Designated Clinical Support Officer'**.

See if Relevant

Tissue and Organ RecipientsTransfusionTrying to Conceive

Additional Information A 'Position Statement on Creutzfeldt-Jakob Disease' is available in the 'Document Library' of 'www.transfusionguidelines.org'.

The use of human gonadotrophin and growth hormone of pituitary origin had stopped in the UK by 1986. Dura mater use stopped in the UK by 1993. The situation in other countries varied so specific dates cannot be given.

Recipients of donated human eggs, sperm or embryos can be accepted to donate, provided they meet the other criteria outlined in the 'Trying to Conceive' entry.

Information Part of this entry is a requirement of the Blood Safety and Quality Regulations 2005.

It also Includes Department of Health decisions about individuals who have been identified at an increased risk of developing a prion related disease.

Reason for change The deferral for recipients of donated eggs or embryos has been removed in line with the most recent update of the SaBTO Microbiological Safety Guidelines.

Update Information This entry was last updated in:
DSG-WB Edition 203, Release 61.

Proctitis

Obligatory **Must not donate if:**
a) Due to ulcerative colitis.
b) Due to Crohn's disease.
c) Requiring treatment.

Discretionary If other causes do not exclude and not on treatment, accept.

See if Relevant Inflammatory Bowel Disease
Malignancy
Sexually Transmitted Disease

Additional Information Proctitis has been linked to stress and food intolerance. It is also associated with infection and this may be sexually transmitted, particularly through anal sex. It may be caused by inflammatory bowel disease (ulcerative colitis and Crohn's disease) and post radiation therapy. The latter is likely to have been given for malignancy so, as with inflammatory bowel disease, will lead to exclusion from donation.

Information Part of this entry is a requirement of the Blood Safety and Quality Regulations 2005.

Reason for change The 'see if relevant' section was updated as part of the implementation of recommendations from the FAIR study.

Update Information This entry was last updated in:
DSG-WB Edition 203, Release 57

Prostate Problems

Obligatory **Must not donate if:**
a) Due to malignancy.
b) The donor has an indwelling urinary catheter, or it is less than seven days since catheterisation.
c) On treatment with anti-androgens.

	d) Accompanied by infection.
	e) Has been referred to a specialist for investigation of a high PSA (Prostate-Specific Antigen) level.
<i>Discretionary</i>	a) For benign prostatic problems, if not requiring treatment, or only taking alpha blockers to relieve symptoms, accept.
	b) If PSA (Prostate-Specific Antigen) levels are being monitored but no referral, biopsy or other treatment is planned, accept.
<i>See if Relevant</i>	<u>Anti-Androgens</u> <u>Infection-General</u> <u>Malignancy</u> <u>Urinary Catheterisation</u> <u>Surgery</u>
<i>Additional Information</i>	Prostate problems become increasingly common as men age. They may cause difficulty in passing water, having to pass water more frequently, or pain and discomfort. Men with benign prostatic hypertrophy (BPH) who do not require treatment, or whose only treatment is with alpha blockers, may donate. If they are being treated with <u>Anti-Androgens</u> (dutasteride (Avodart [®]) or finasteride (Proscar [®])) special precautions are needed while taking these drugs and for some time afterwards. Malignancy must lead to permanent deferral. Infection, or the possibility of infection, associated with catheterisation will also lead to deferral. The interpretation of PSA (Prostate-Specific Antigen) levels depends on a number of factors. If the levels were thought to have been significantly abnormal, the individual would have been referred for biopsy or other investigations or treatment.
<i>Information</i>	Part of this entry is a requirement of the Blood Safety and Quality Regulations 2005.
<i>Reason for change</i>	The deferral for donors who use a urinary catheter has been reworded and the See if Relevant section has been revised.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 61.

Psoriasis

<i>Includes</i>	Psoriatic Arthritis
<i>Obligatory</i>	Must not donate if: a) Has ever taken etretinate (Tigason [®]). b) Less than 36 months from the last dose of acitretin (Neotigason [®]). c) Less than 6 months from the last dose of any treatment that may affect the immune system. d) Generalised or severe. e) There is secondary infection.
<i>Discretionary</i>	a) If mild, the venepuncture site is unaffected and only using topical treatment, accept. b) If the donor: <ul style="list-style-type: none"> • has been established on oral treatment for their disease with only one of the following drugs: Methotrexate, Sulfasalazine, Hydroxychloroquine or Azathioprine, and • their disease is controlled by medication, and • the dose of the drug has not increased in the previous 6 months, and • the venepuncture site is not affected, and

- the donor is well,

accept.

c) If the donor:

- is receiving PUVA or UVA therapy for their skin disease, and
- their disease is controlled, and
- the venepuncture site is not affected, and
- the donor is well,

accept.

See if Relevant Autoimmune Disease
Monoclonal antibody therapy and other Biological Modalities
Steroid Therapy

Additional Information Psoriasis is primarily a skin condition caused by an autoimmune process. Sometimes the disease is treated with powerful drugs and/or ultraviolet radiation to suppress the underlying autoimmune process. This may be with treatment with PUVA, methotrexate, ciclosporin, hydroxycarbamide etc. and this may alter the body's defence mechanisms to infection.

Etretinate (Tigason[®]) and acitretin (Neotigason[®]) can cause birth defects in babies exposed to them while inside the womb. It is important to allow time for the drug to be cleared from the blood of a donor. It takes longer to clear some drugs than others.

Reason for change The text has been further updated to ensure consistency with other DSG references to immunosuppression.

Update Information This entry was last updated in:
WB-DSG Edition 203 Release 74

Pyrexia

Also Known As Fever.

Definitions **Pyrexia:**
A temperature of 38.0°C or above.

1. Not Related to Travel

Obligatory **Must not donate if:**
Less than two weeks from an episode of pyrexia.

Discretionary If related to a common cold or other upper respiratory tract infection, but not influenza, from which the donor is now recovered or recovering, accept.

See if Relevant Infection - Acute
Infection - General

Additional Information A raised temperature may be a sign of an infection, which could be passed on through a donation. Waiting two weeks from when the temperature returns to normal reduces the risk of infection being transmitted by the donation.

There is no evidence that common colds and upper respiratory tract infections can be passed on by donation but it is still necessary to wait until any such infection is obviously getting better before allowing donation.

2. Related to Travel

See Malaria
Tropical Viruses
West Nile Virus

Viral Haemorrhagic Fever
The Geographical Disease Risk Index

<i>Reason for change</i>	The definitions and scope have been revised. The See if Relevant list in part 2 has been expanded.
<i>Update Information</i>	This entry was last updated in: WB-DSG Edition 203 Release 72

Rabies

1. Affected Individual

<i>Obligatory</i>	Must not donate if: Diagnosed with Rabies even if now recovered.
<i>Discretionary</i>	If exposure to a potentially rabid animal has been managed with passive immunisation and/or immunisation, accept if it is at least 12 months post exposure and fully cleared by the treating physician.
<i>Additional Information</i>	Once symptomatic, rabies is almost always fatal. There is not enough information on individuals who have recovered to know if they may still present an infection risk and, if so, for how long.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 01.
<i>Reason for change</i>	This is a new entry.

2. Immunisation - Post Exposure

<i>Obligatory</i>	Must not donate until: At least 12 months post exposure and fully cleared by the treating physician.
<i>See if Relevant</i>	<u>Immunoglobulin Therapy</u> <u>Animal Bite (Non-Human)</u>
<i>Additional Information</i>	It is essential that any rabies virus has been eliminated from the system before a donation is accepted. Waiting at least 12 months post exposure and until the individual is fully cleared by the treating physician should make sure that the virus has been cleared.
<i>Information</i>	This entry is compliant with the Blood Safety and Quality Regulations 2005.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 55.
<i>Reason for change</i>	A link has been added to 'Animal Bite (Non-Human)'.

3. Immunisation - Non-Exposed

<i>Discretionary</i>	If non-exposed, accept.
<i>Information</i>	This entry is compliant with the Blood Safety and Quality Regulations 2005.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 202, Release 02.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 01.

Update Information This entry was last updated in:
DSG-WB Edition 203, Release 55.

Update Information This entry was last updated in:
DSG-WB Edition 202, Release 02.

Radiation Therapy

<i>Obligatory</i>	<p>1. Must not donate if: For malignancy other than basal cell carcinoma.</p> <p>2. For other treatments: Refer to a 'Designated Clinical Support Officer'.</p>
<i>Discretionary</i>	<p>a) If fully recovered and is acceptable according to immunosuppression advice, accept.</p> <p>b) If for basal cell carcinoma or ductal carcinoma in situ of the breast, all treatment has been completed, the donor has been discharged from follow up and is eligible under the Malignancy Guideline, accept.</p>
<i>See if Relevant</i>	<p><u>Autoimmune Disease</u> <u>Immunodeficiency</u> <u>Malignancy</u></p>
<i>Additional Information</i>	<p>Radiation therapy is sometimes used for non-malignant conditions, particularly for some skin conditions. It is often used as a substitute for other treatments that work by suppressing the immune system such as high dose steroids and cytotoxic drugs. More information is likely to be required before a decision can be made as to if an individual can donate. This why a referral to a 'Designated Clinical Support Officer' is required.</p>
<i>Reason for change</i>	<p>Link updated from 'Immunosuppression' to 'Immunodeficiency' in the 'See if Relevant' section.</p>
<i>Update Information</i>	<p>This entry was last updated in: WB-DSG Edition 203 Release 73</p>

Radionuclides

<i>Definitions</i>	<p>Radionuclides: These are unstable materials that emit radioactivity when they decay. They are used in some special investigations carried out in radiology (X-ray) and medical physics departments. They may be breathed in, taken by mouth or given by injection.</p>
<i>Obligatory</i>	<p>1. Radioactive iodine therapy: Must not donate if: a) For malignancy.</p> <p>b) Administered in the preceding six months for a non-malignant condition.</p> <p>2. Other treatment or investigation: Refer to a 'Designated Clinical Support Officer'.</p>
<i>See if Relevant</i>	<p><u>Investigations</u> <u>Malignancy</u> <u>Thyroid Disease</u></p>
<i>Additional Information</i>	<p>In general, those used for diagnostic purposes are cleared within 24 hours. Some, e.g. radioactive iodine, have long half-lives and affected donors must not be accepted unless at least six months have passed. This is because we do not wish to transfuse radio-active material to recipients, particularly where it may affect a child or an unborn baby.</p>
<i>Reason for change</i>	<p>A 'Definition' of 'Radionuclides' has been added.</p> <p>The 'Additional Information' has been extended.</p>

Update Information This entry was last updated in:
DSG-WB Edition 203, Release 01.

Raynaud's Syndrome

Obligatory **If part of an autoimmune process:**
See:
Autoimmune Disease

Discretionary If not part of an autoimmune process, even if the donor is taking vasodilators, accept.

Reason for change Both the 'Obligatory' and the 'Discretionary' entries have been modified to be consistent with updates to the entries for 'Blood Pressure - High' and 'Autoimmune Disease'.

Update Information This entry was last updated in:
DSG-WB Edition 203, Release 01.

Respiratory Disease

Obligatory **Must not donate if:**
a) Out of breath on minimal exertion.
b) Has acute or chronic infection including bronchiectasis.
c) Has cystic fibrosis.

See if Relevant Asthma
Autoimmune Disease
Infection - General
Sarcoidosis
SARS
Steroid Therapy

Additional Information If a potential donor is out of breath on minimal exertion (for instance, climbing a single flight of stairs), taking a unit of blood may reduce the amount of oxygen that can be carried in the blood to a level that makes them unwell. Bronchiectasis is associated with recurrent acute infections and chronic infection. Although these do not usually cause the sufferer harm the transmission of these may result in significant problems for severely ill and often immunocompromised recipients.

Reason for change To clarify the position for potential donors with Bronchiectasis

Update Information This entry was last updated in:
DSG-WB Edition 203, Release 27.

Rheumatic Fever

Obligatory **Must not donate if:**
a) It is less than 24 months from any symptomatic disease.
b) Has caused permanent heart valve damage.

Additional Information The Blood Safety and Quality Regulations 2005 state donation is not allowed until 24 months following the date of cessation of symptoms

Rheumatic fever can cause damage to the heart and this could make it unsafe to give blood.

Information Part of this entry is a requirement of the Blood Safety and Quality Regulations 2005.

Reason for change The 'Obligatory' entry has been changed and the 'Additional Information' has been amended.

Update Information This entry was last updated in:
DSG-WB Edition 203, Release 01.

Sarcoidosis

1. Acute

<i>Obligatory</i>	Must not donate if: a) Not recovered. b) Less than five years from both finishing all treatment and full recovery.
<i>Discretionary</i>	If more than five years since finishing all treatment and full recovery, accept
<i>Additional Information</i>	Acute sarcoidosis is normally a self limiting disease and does not require treatment in about 90% of cases. The cause is not known but there appears to be an immune defect that can run in families. Because of the uncertainty with this condition, only potential donors who have fully recovered and been off all treatment for at least five years may donate.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 01.
<i>Reason for change</i>	New guidance to accept after full recovery and off all treatment for at least five years has been added. 'Additional Information' has been added.

2. Chronic

<i>Obligatory</i>	Must not donate.
<i>Additional Information</i>	Chronic Sarcoidosis can cause a range of problems, particularly with the lungs but also with the heart, that may pose risks for a potential donor. The treatments used may also cause immunosuppression. For these reasons people with this condition should not donate.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 01.
<i>Reason for change</i>	'Additional Information' has been added.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 01.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 01.

Sex Worker

<i>Obligatory</i>	Must not donate if: Has ever received money or drugs for sex.
<i>Discretionary</i>	If three or more months have elapsed since the donor last received money or drugs for sex, accept
<i>See if Relevant</i>	<u>Blood Safety Entry</u>
<i>Additional Information</i>	In this context sex is defined as vaginal, oral or anal sex with or without a condom/protective
<i>Information</i>	This is a requirement of the Blood Safety and Quality Regulations 2005.
<i>Reason for change</i>	This entry was updated to remove the reference to a separate entry for Northern Ireland. This is to reflect changes in donor selection criteria for donors in Northern Ireland (1st June 2020) which are in line with the other UK Blood Services and the SaBTO Donor Selection Criteria Review Report (2017).
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 50.

Sexually Transmitted Disease

1. Affected Individual

Obligatory **Refer to WBDSG entry for the condition.**

If there is no specific entry, must not donate if:
Less than three months from completing treatment.

See if Relevant Blood Safety Entry

For chlamydia (excluding Lymphogranuloma Venereum) see:
Infection - Acute

For genital warts see:
Warts

For genital herpes see:
Herpes Simplex

For syphilis see:
Syphilis

2. Current or Former Sexual Partner of Affected Individual

Obligatory **Refer to WBDSG entry for the condition.**

If there is no specific entry or the entry has no guidance on assessing sexual partners, must not donate if:

- a) The potential donor is undergoing, or waiting for, investigations.
- b) The potential donor required treatment and it is less than three months since completing that treatment.
- c) The potential donor did not require treatment and it is less than three months from the last sexual contact with the infected partner.

Discretionary a) If the donor's sexual partner has been diagnosed with chlamydia, genital warts or genital herpes and the donor is not undergoing treatment or investigation, accept.

b) If there is no WBDSG entry for the condition, or the entry has no guidance on assessing sexual partners, and it is more than three months since the donor's sexual partner completed treatment, accept.

See if Relevant Blood Safety Entry

For chlamydia (excluding Lymphogranuloma Venereum) see:
Infection - Acute

For genital warts see:
Warts

For genital herpes see:
Herpes Simplex

For syphilis see:
Syphilis

Additional Information Certain sexually transmitted infections, such as gonorrhoea, are more likely to be associated with other sexually transmitted infections and/or blood borne viruses that can be passed on

through blood and component donation. A three-month deferral is required so that there is less risk of other infections being missed by the Blood Services and then being passed on to a recipient of donated material.

Reason for change This entry was updated to support the implementation of recommendations from the FAIR study; the deferral period after a sexually transmitted disease, or treatment after sexual contact with an infected person, has been reduced to three months.

Update Information This entry was last updated in TDSG-BM Edition 203, Release 57

Shunts, Stents and Devices

Includes **Shunts**

- For hydrocephalus e.g. ventriculo-peritoneal, -atrial, -pleural and lumboperitoneal shunts

Stents

- Vascular stents including coronary artery stents
- Urinary tract stents including ureteric stents
- Ophthalmic stents including nasolacrimal and Schlemm canal stents

Pacemakers

- Cardiac pacemakers
- Gastric pacemakers

Implanted neuromodulator and nerve stimulator devices, used for:

- Neuropathic pain – includes spinal cord and peripheral nerve stimulators
- Bladder dysfunction - includes sacral nerve stimulators
- Gastroparesis – sometimes referred to as gastric pacemakers

Glucose monitoring devices

- Real time continuous glucose monitors (rtCGM)
- Intermittently scanned glucose monitors (isCGM) – flash glucose monitors

Obligatory **Must not donate.**

Discretionary a). If the indication for an implanted neuromodulator device does not preclude donation and the site of implantation is fully healed, accept.

b). If an ophthalmic stent has been successfully inserted with good effect, the area has fully healed and there is no infection, accept.

c) If the reason for using a glucose monitoring device does not preclude donation and there is no inflammation or infection at, or around, the site of the skin sensor, accept.

See if Relevant Arrhythmia
Cardiovascular Disease
Diabetes Mellitus
Eye disease
Gastrointestinal Disease
Kidney and Bladder Disease
Neurosurgery
Spina Bifida

Additional Information Some shunts and stents can be a source of bacterial infection due to their location in the body and infection can be present without symptoms.

Bacteria can be a serious threat to anybody receiving blood or blood components. This is because bacteria can multiply to dangerous levels after collection.

Implanted neuromodulator devices and some stents are not a covert infection risk once the implantation site is fully healed and there are no signs of infection or inflammation.

Care should be taken to ensure that the underlying condition that requires the use of a neuromodulator device does not preclude donation, and that the donor is well at the time of donation.

Care should be taken to ensure that the reason that a glucose monitoring device is being used does not preclude donation or make it inadvisable at that time. These devices are not a covert infection risk as any local infection or inflammation should be apparent. The Diabetes Mellitus entry should always be referred to for diabetic donors using these devices.

Reason for change Addition of advice for donors with glucose monitoring devices.
Addition of Diabetes Mellitus to See if Relevant section.

Update Information This entry was last updated in:
WB-DSG Edition 203 Release 72

Sickle-Cell Trait

Obligatory **Whole Blood donor:**
Not suitable for intra-uterine or neonatal use.

Discretionary For adult use only, accept.

Additional Information The red blood cells from people with sickle cell trait can be safely transfused into most adults. They are however not thought to be suitable for intra-uterine or neonatal use as there is a higher risk of the cells sickling and causing harm to the baby.

For some individuals with sickle cell trait it will not be possible to process their blood. For this reason they may be asked not to donate.

Reason for change To allow component donors with sickle-cell trait to donate.

Update Information This entry was last updated in:
DSG-WB Edition 203, Release 16

Skin Disease

Obligatory **Must not donate if:**

- a) The donor has a condition that is infected or infectious e.g. Scabies.
- b) History of malignancy.
- c) The venepuncture site is affected.
- d) Required application of steroid, tacrolimus (Protopic[®]) or pimecrolimus (Elidel[®]) creams over large areas for periods of more than three weeks in the last six months.
- e) Ever been treated with Etreinate (Tigason[®]).
- f) Less than 36 months from the last dose of acitretin (Neotigason[®]).
- g) Less than four weeks from the last does of isotretinoin (Roaccutane[®]) or Alitretinoin (Toctino[®]).
- h) Has any current open skin wounds or infection.

Discretionary

- a) If occasional use of steroid, tacrolimus (Protopic[®]) or pimecrolimus (Elidel[®]) or other creams over small areas of skin and none of the above apply, accept.
- b) If chronic superficial fungal infection (e.g. ringworm, athlete's foot, chronic fungal nail infection or tinea) on local therapy only or has been in contact with an infected individual, accept.
- c) If in contact with scabies but not obviously infected, accept.
- d) If malignancy was a basal cell carcinoma (rodent ulcer) and treatment is completed and all wounds healed, accept.

For donors with Lichen Sclerosus requiring treatment other than topical steroid therapy only, excluding Etreinate (Tigason[®]):

- e) If more than 24 months from completing treatment, have no areas of open wound or infection, have no history of associated malignancy and symptoms are controlled with or without intermittent use of topical steroid therapy only, accept.

See Is there is a specific A-Z index entry for the treatment and/or condition you are concerned about?

See if Relevant

- Acne
- Anaemia
- Autoimmune disease
- Dermatitis
- Herpes Simplex
- Immunodeficiency
- Infection - General
- Malignancy
- Psoriasis
- Steroid Therapy
- Surgery
- Thrush
- Thyroid disease
- Wounds, Mouth and Skin Ulcers

Additional Information A donor who has been in contact with scabies but has no symptoms (e.g. itching) does not pose a risk to other donors or staff.

Damaged skin can increase the risk of infection contaminating a donation. For this reason a venepuncture should not be performed through an area of affected skin.

Many malignancies spread through the blood stream. It is therefore considered safer not to accept donations of blood from people who have been diagnosed with malignancy. Treated basal cell carcinoma is an exception to this as it is not spread through the blood stream.

Initial treatment of Lichen Sclerosus is through specialist care with potent steroid therapies.

Treatment can also be with methotrexate and retinoids such as Etreinate (Tigason[®]) or acitretin (Neotigason[®]). If taken systemically these can cause birth defects for babies exposed to them before birth. It is important to allow time for the drug to be cleared from the blood of a donor. Some drugs take longer to be cleared than others.

Under normal circumstances the use of topical treatment with steroid, tacrolimus and pimecrolimus will not result in blood levels which cause suppression of the adrenal system or immune response. Side effects are more likely if there is a skin barrier defect or high doses are used over large areas for extended periods. A large area of skin is defined as >9% (Wallace Rule of Nines). 1% is equal to the area of the closed digits and palm of the donor's hand.

Reason for change Link updated from 'Immunosuppression' to 'Immunodeficiency' in the 'See if Relevant' section.

Update Information This entry was last updated in:
WB-DSG Edition 203 Release 73

Sleep Apnoea

<i>Obligatory</i>	Must not donate.
<i>Discretionary</i>	<p>If:</p> <ul style="list-style-type: none"> • the donor's symptoms are well controlled with lifestyle change and/or treatment, and • the donor reports no daytime sleepiness (and there is no restriction on activities such as driving), and • no underlying cause for sleep apnoea has been identified which would preclude donation, <p>accept.</p>
<i>See if Relevant</i>	<p><u>Central Nervous System Disease</u> <u>Cerebrovascular Disease and Intracranial Haemorrhage</u> <u>Donor Weight</u> <u>Surgery</u></p>
<i>Additional Information</i>	<p>Sleep Apnoea can lead to daytime sleepiness, difficulty in concentration and an increased risk of accidents. Not everyone with sleep apnoea will require treatment. Corrective measures that can improve symptoms include:</p> <ul style="list-style-type: none"> • weight loss • the use of gum shield • smoking and alcohol reduction/cessation • sleep hygiene <p>Surgery may be required for some individuals e.g. tonsillectomy, uvuloplasty, mandibular surgery, or nasal septum surgery.</p> <p>Some individuals may use CPAP (Continuous Positive Airway Pressure) and BiPAP (Bi-level Positive Airway Pressure) Ventilation devices while sleeping. These use a close-fitting mask to deliver pressurised air to the lungs through the respiratory cycle as treatment for Sleep Apnoea.</p>
<i>Reason for change</i>	Link updated from 'Central Nervous System Disease' to 'Cerebrovascular Disease and Intracranial Haemorrhage' in the 'See if Relevant' section.
<i>Update Information</i>	This entry was last updated in: WB-DSG Edition 203 Release 73

Smallpox Immunisation

1. Immunised Individual

<i>Obligatory</i>	<p>Must not donate if:</p> <p>a) The inoculation site has not fully healed.</p> <p>b) Any secondarily infected site has not fully healed.</p> <p>c) Less than eight weeks from inoculation or from the appearance of any secondarily infected site.</p>
<i>Additional Information</i>	Smallpox immunisation is with live virus. By eight weeks, the infection caused by the inoculation should have been controlled. If the wound has not healed it is possible that there may still be infection present. We do not want to pass the virus, or other infection, on to other donors, staff or to people receiving donated material.
<i>Information</i>	This entry is compliant with the Blood Safety and Quality Regulations 2005.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 202, Release 02.

2. Contact

<i>Obligatory</i>	Must not donate if: a) Any secondarily infected site has not yet healed. b) Less than eight weeks after secondarily infected site appeared.
<i>Discretionary</i>	If no new skin lesions, accept.
<i>Additional Information</i>	Close contacts of vaccinees (household or direct bodily contact) may become secondarily infected from direct skin contact with an infected inoculation site or from virus on clothing, bedding, dressings etc. If infection occurs, a new skin rash, blister or sore appears at the site of contact, which could be anywhere on the body. The rash represents a secondary vaccination site and presents exactly the same potential risk to patients, other donors and staff as that from a person who has been intentionally immunised.
<i>Information</i>	This entry is compliant with the Blood Safety and Quality Regulations 2005.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 202, Release 02.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 202, Release 02.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 202, Release 02.

Smoking

<i>Discretionary</i>	Accept.
<i>See if Relevant</i>	<u>Smoking Cessation</u>
<i>Additional Information</i>	Smokers may donate. However all donation sessions have a no smoking policy to comply with the law.
<i>Reason for change</i>	This is a new entry.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 01.

Smoking Cessation

<i>Includes</i>	Bupropion (Amfebutamone [®] , Zyban [®]), nicotine replacement therapy (Nicorette [®] , Nicotinell [®] , NiQuitin [®]) and varenicline (Champix [®]).
<i>Obligatory</i>	Must not donate if: Experiencing symptoms related to treatment.
<i>Discretionary</i>	Donors using nicotine replacement therapy (patches, sprays etc), bupropion (Amfebutamone [®] , Zyban [®]) or varenicline (Champix [®]), if well, accept.
<i>See if Relevant</i>	<u>Complementary Therapy</u> (includes acupuncture)
<i>Additional Information</i>	Anti-smoking treatments can cause dizziness and nausea. Taking a donation from people who are affected, may make these symptoms worse.
<i>Reason for change</i>	Varenicline (Champix [®]) has been added to the list of anti-smoking treatments.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 01.

South American Trypanosomiasis

Also Known As Chagas disease.

Excludes Donors who will only donate plasma for fractionation. See South American Trypanosomiasis - plasmapheresis donors

1. Affected Individual

Obligatory **Must not donate.**

Additional Information South American trypanosomiasis is caused by infection with a protozoal parasite, *trypanosoma cruzi*. It is a persistent infection that is known to be transmitted by transfusion. At present there is no certain cure for the infection, so anyone who has ever been infected cannot donate.

2. Risk

Obligatory **Must not donate if:**

- a) Born in South America or Central America (including Mexico).
- b) Mother was born in South America or Central America (including Mexico).
- c) Has had a transfusion in South America or Central America (including Mexico).
- d) Has lived and/or worked in rural subsistence farming communities in these countries for a continuous period of four weeks or more.

Discretionary If at least four months following the date of last exposure (or if transfused prior to 1980) and a validated test for *T. cruzi* antibody is negative, accept.

See if Relevant Geographical Disease Risk Index for countries with *T. cruzi* risk
Transfusion

Additional Information Infection with *T. cruzi* is very common in many parts of South or Central America and is often symptomless. It can be passed from an infected mother to her unborn baby and by transfusion. The insect that passes the infection on is only common in rural areas and the greater time that an individual has spent living in housing conditions with thatched roofs or mud lined walls which harbour the insect vector, the greater their risk of becoming infected. Testing is available and should be performed if there is a possibility of infection. Waiting four months from the last time of exposure allows time for the antibodies that are tested for to develop.

Camping or trekking in the jungle in South or Central America (including Mexico) is not considered of high enough risk to merit exclusion.

Information This is a requirement of the Blood Safety and Quality Regulations 2005.

Reason for change Entry updated to exclude donors who will only donate plasma for fractionation.

Update Information This entry was last updated in:
WB-DSG Edition 203 Release 71

South American Trypanosomiasis - plasmapheresis donors

For donors who will donate whole blood, platelets and other cellular components see South American Trypanosomiasis

1. Affected Individuals

Obligatory **Must not donate**

Additional Information South American trypanosomiasis is caused by infection with a protozoal parasite, *Trypanosoma cruzi*. It is a persistent infection that is known to be transmitted by transfusion. At present there is no certain cure for the infection, so anyone who has ever been infected cannot donate.

2. Risk

<i>Excludes</i>	This entry only applies for donors who will only donate plasma for fractionation. It should not be used for donors who will donate whole blood, platelets and other cellular components.
<i>Discretionary</i>	Accept
<i>See if Relevant</i>	<u>The 'Geographical Disease Risk Index'</u>
<i>Additional Information</i>	The processes used to fractionate plasma include several measures that inactivate or remove <i>T. cruzi</i> parasites. This means that the Trypanosoma Cruzi risks described in the GDRI do not need to be applied for donors who will only donate plasma for fractionation. <i>T. cruzi</i> antibody testing is not required for these donors.
<i>Reason for change</i>	This is a new entry.
<i>Update Information</i>	This entry was last updated in: WB-DSG Edition 203 Release 71

Spina Bifida

<i>Obligatory</i>	<p>Must not donate if:</p> <p>a) Has an indwelling shunt.</p> <p>b) The donor has an indwelling urinary catheter, or it is less than seven days since catheterisation.</p> <p>c) Has a pressure sore.</p>
<i>See if Relevant</i>	<p><u>Anti-Androgens</u></p> <p><u>Indwelling Shunts and Stents and Implanted Devices</u></p> <p><u>Infection-General</u></p> <p><u>Malignancy</u></p> <p><u>Surgery</u></p> <p><u>Urinary Catherisation</u></p>
<i>Additional Information</i>	All of the conditions under 'Obligatory' put the potential donor at increased risk of bacteria being present in the blood stream. Bacteria can be a serious threat to anybody receiving blood or blood components. This is because they can multiply to dangerous levels after collection.
<i>Reason for change</i>	The deferral for donors who use a urinary catheter has been reworded and the See if Relevant section has been revised.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 61.

Splenectomy

<i>Obligatory</i>	<p>Must not donate if:</p> <p>a) For malignancy.</p>
-------------------	---

b) For a myeloproliferative disorder.

c) For haemolytic anaemia.

Discretionary a) If for trauma, when recovered, even if taking prophylactic antibiotics, accept.

b) If for immune thrombocytopenia, if at least five years from recovery, even if taking prophylactic antibiotics, accept.

See if Relevant Haematological Disease
Immune Thrombocytopenia
Malignancy
Surgery
Transfusion

Additional Information If haemolysis is severe enough to require splenectomy, it is likely to significantly reduce affect red cell survival in storage. This may be dangerous for any recipient.

Information This is a requirement of the Blood Safety and Quality Regulations 2005.

Reason for change The permanent deferral of individuals who have had a splenectomy for immune thrombocytopenia has been removed.

Relevant links and 'Additional Information' have been added.

Update Information This entry was last updated in:
DSG-WB Edition 203, Release 01.

Steroid Therapy

Obligatory **Must not donate if:**

a) Taking steroid tablets, injections, or enemas, or applying creams over large areas for periods of more than three weeks in the last six months.

b) Less than seven days after completing a course of oral or injected steroids for asthma, other disorders associated with allergy or a musculoskeletal condition.

c) A donor has needed long term (six months or more) treatment within the last 12 months.

Discretionary a) If occasional use of creams over small areas of skin for minor skin complaints, accept.

b) If using steroid inhalers for prophylaxis, accept.

c) If using steroid eye drops, nasal spray or ear drops for control of allergic symptoms, accept.

d) If more than seven days from completing a course of intramuscular, periarticular or intra-articular injected steroids for a musculoskeletal condition, accept unless the musculoskeletal condition itself would lead to deferral.

See if Relevant Adrenal Failure
Allergy
Asthma
Autoimmune Disease
Skin Disease

Additional Information A large area of skin is defined as >9% (Wallace Rule of Nines). 1% is equal to the area of the closed digits and palm of the donor's hand.

Steroid therapy in high doses causes immunosuppression. This may mask infective and inflammatory conditions that would otherwise prevent donation.

Some individuals have to take replacement steroid hormones because they do not produce enough themselves. The dose of these must be increased during times of stress. It is considered that taking blood from people who need replacement therapy may put them at unnecessary risk.

Long term steroid therapy may cause temporary adrenal dysfunction. Waiting 12 months from the last dose allows time for the adrenal glands to recover.

Reason for change The text has been updated to ensure consistency with other DSG references to immunosuppression. The see if relevant section has been revised.

Update Information This entry was last updated in:
WB-DSG Edition 203 Release 71

Stoma

Obligatory **Must not donate if:**
a) For malignancy.

b) Inflammatory bowel disease.

Discretionary If the reason for the stoma is not of itself a reason to exclude and the stoma is healthy, accept.

See if Relevant Disabled Donor
Endoscopy
Inflammatory Bowel Disease
Malignancy
Surgery

Additional Information A stoma is usually performed either for malignancy or inflammatory bowel disease. It may be temporary or permanent.

If it is clear that a stoma has been performed for a different reason, that itself would not lead to deferral (e.g. following an accident or non-malignant obstruction), it is possible that the donor may be accepted. If there is any doubt:
Refer to a '**Designated Clinical Support Officer**'.

Information This is a requirement of the Blood Safety and Quality Regulations 2005.

Reason for change Additional links and 'Additional Information' have been added.

Update Information This entry was last updated in:
DSG-WB Edition 203, Release 01.

Superficial Thrombophlebitis

Also Known As Superficial vein thrombosis; Thrombophlebitis; Phlebitis

Definitions Inflammation of a superficial vein due to a blood clot. For the purposes of donor selection, superficial thrombophlebitis is not considered to be a significant clotting episode unless the clot has extended to a deep vein.

Obligatory **Must not donate if:**
a) It is less than 7 days since recovery and cessation of treatment; or
b) It is less than 14 days since recovery from an associated infection; or
c) The donor is under investigation for recurrent superficial thrombophlebitis; or
d) The donor has a history of recurrent superficial thrombophlebitis associated with thrombophilia; or
e) An underlying cause has been identified which would preclude donation; or
f) It is associated with poor skin integrity of the lower limbs, arising as a complication of varicose veins and/or chronic venous insufficiency.

- Discretionary* a) If associated with Deep Vein Thrombosis (DVT), refer to the Thrombosis and Thrombophilia entry.
- b) Otherwise, if:
- It is at least 7 days since recovery and treatment (including NSAIDs or anticoagulants), and
 - It is at least 14 days since recovery from an associated infection, and
 - The donor is not under investigation, and
 - Any underlying cause does not preclude donation, and
 - The donor does not have chronic skin damage which could pose an infection risk,

Accept.

See if Relevant Thrombosis and Thrombophilia
Varicose Veins and Chronic Venous Insufficiency

Additional Information Superficial thrombophlebitis is a common condition usually, but not exclusively, affecting the lower limbs. It is caused by clot formation in a superficial vein which in turn gives rise to inflammation with associated pain, tenderness, redness and hardness of the vein. Usually the condition is mild and self-limiting, settling over a few weeks. Treatment is typically pain relief and NSAIDs. Antibiotics should only be prescribed if there is associated infection.

Superficial thrombophlebitis can sometimes occur in association with DVT. Some patients with thrombophlebitis may be treated with anticoagulants to reduce any risk of the clot extending to a deep vein.

Risk factors for superficial thrombophlebitis include varicose veins; a previous history of thrombophlebitis; IV cannulation; female sex; the oral contraceptive pill or hormone replacement therapy; thrombophilia, increasing age; some autoimmune diseases; and cancer. Provided a serious underlying cause is not suspected or has been excluded, a history of thrombophlebitis on its own is not a reason for deferral. However donors with a history of thrombophilia associated with repeated episodes of thrombophlebitis should be deferred.

Individuals with complications of varicose veins affecting the lower limb are at risk of recurrent superficial thrombophlebitis. It is important that donors with recurrent episodes are asked about any skin damage, such as inflamed venous eczema or skin ulceration, before being accepted. This is to reduce the risk of bacterial contamination of donated blood arising from a breach of the normal skin defences.

Reason for change This is a new entry. It replaces the previous 'Phlebitis' entry.

Update Information This entry was last updated in:
DSG-WB Edition 203, Release 68

Surgery

Definitions **Recovery from surgery:** Donors can be considered to be recovered if they:

- are well
- are back to activities of daily living (e.g. housework, employment, driving)
- have regained mobility

Major Surgery for the purposes of donor selection:
Any surgical procedure where recovery is not achieved within two months.

Obligatory **Must not donate if:**

a) For malignancy or other condition that would preclude donation.

b) All wounds are not healed.

c) There are signs or symptoms of any infection.

- d) Not recovered.
- e) Less than four months from major surgery.
- f) Less than seven days from other surgery.
- g) Less than four months from any flexible endoscopic procedure.
- h) Requiring post-operative treatment or follow-up, except routine physiotherapy.
- i) Received a transfusion since 1st January 1980.
- j) If waiting for surgery that is:
 - expected to occur within three months, or
 - required due to possible malignancy
- k) Less than three months from a surgical procedure performed outside of the UK and Republic of Ireland (ROI).
- l) Less than seven days after completing postoperative prophylactic anticoagulant treatment.

Discretionary

a) If all other criteria are met and the donor has received a metal-on-metal hip replacement even if being monitored for blood chromium or cobalt levels, accept.

b) If the donor is waiting for surgery that is not required for possible malignancy, and:

- the procedure is not expected to take place within three months, or
- the procedure is minimally invasive and it is not expected to take place within one month,

accept.

c) If the donor has recovered from surgery within the UK and ROI, and:

- it is more than four months since major surgery, or
- it is more than seven days since any other form of surgery, and
- it is more than four months since a flexible endoscopic procedure, and
- there was no malignancy and the reason for surgery does not otherwise preclude donation, and
- the donor did not receive a transfusion since 1st January 1980, and
- all wounds are healed, and
- there are no signs or symptoms of infection, and
- the donor has been discharged from postoperative follow-up, and
- the donor does not require ongoing postoperative treatment except routine physiotherapy, and
- it is more than seven days from finishing any anticoagulant treatment given to prevent postoperative thrombosis e.g. DVT,

accept.

d) If it is more than three months since a surgical procedure performed outside of the UK and ROI, and all other criteria for surgery performed within the UK and ROI are met (see point c above), accept.

See if Relevant

Anaesthetic
Anticoagulant Therapy
Cervical Dysplasia
Disabled Donor
Dental Treatment
Endoscopy
Eye Disease
Malignancy
Neurosurgery
Tissue and Organ Recipients

Transfusion
Wounds, Mouth and Skin Ulcers
Xenotransplantation

Additional Information Surgery may cause significant blood loss. It is important that donors waiting for an operation should not be put at risk of anaemia or poor iron stores by donating prior to planned surgery. Unless the type of surgery planned is unlikely to result in significant blood loss the donor should be deferred until after their planned surgery. This will minimize their own chance of needing a transfusion, which would of course prevent them from continuing as a donor. It is also important not to hinder the recovery of the donor. This requires waiting until they are fully recovered before they donate again.

This guidance presumes that a validated NAT test for hepatitis C is negative. If this test is stopped the guidance will change.

Surgery may place the donor at risk of infection, either from unhealed wounds, or due to infection risks from infected staff or equipment. Although these risks are very small it is important to wait long enough for the risks to have gone or until the tests performed by the Blood Services can pick up any infection that they test for that may have been transmitted to the donor through their surgery. As there may be uncertainty about these risks for surgery performed outside of the UK and ROI, a deferral period of three months is required.

Minimally invasive surgery includes superficial skin procedures and procedures performed under infiltration with local anaesthetic agents and/or sedation. This does not include procedures performed under regional anaesthesia (e.g. spinal, epidural) which may be used where joints and major body cavities may be accessed. The use of general anaesthesia may not indicate the invasiveness of a procedure and should not be used as a substitute to assessment of the donor regarding the procedure and their recovery. Donors can be accepted for donation once it is more than seven days since a surgical procedure as long as they also fulfil all other criteria. Donors who have had minimally invasive surgical procedures are unlikely to have systemic effects from the surgery requiring recovery time. However, care should be taken to ensure that all wounds are dry and healing. An open wound is a risk for bacteria entering the blood. Bacteria can be a serious threat to anybody receiving blood or blood components. This is because bacteria can multiply to dangerous levels after collection.

Donors being monitored for chromium or cobalt levels following a metal-on-metal hip replacement can be accepted for donation.

Completion of postoperative monitoring, treatment and follow-up should be confirmed for every donor returning to donate. Thromboprophylaxis may be continued, usually for a few weeks only, after discharge from hospital. Donors who are recovered and are attending only physiotherapy appointments for ongoing rehabilitation can be accepted.

Information Part of this entry is a requirement of the Blood Safety and Quality Regulations 2005.

Reason for change The entry has been revised to include a definition of recovery and amendment of the definition of major surgery. The deferral after major surgery has been shortened. Information regarding donor eligibility after non-major surgery has been added. Specific guidance for surgery overseas, donors awaiting surgery and postoperative thromboprophylaxis has been added.

Update Information This entry was last updated in:
DSG-WB Edition 203, Release 65

Syphilis

1. Affected Individual

Obligatory **Must not donate.**

See if Relevant Blood Safety Entry

Additional

<i>Information</i>	Many donors with treated syphilis will persistently test positive to the screening tests used by the Blood Services, even if treated many years ago. This will mean they will not be able to donate.
<i>Information</i>	Part of this entry is a requirement of the Blood Safety and Quality Regulations 2005.

2. Current or Former Sexual Partner of Affected Individual

<i>Obligatory</i>	Must not donate if: a) The potential donor was diagnosed with syphilis (See 1. Affected Individuals). b) It is less than three months since last sexual contact with an infected partner.
<i>Discretionary</i>	a) If it is more than three months from the last sexual contact with an infected partner, accept. b) If it is more than three months since the infected partner has completed treatment, accept.
<i>See if Relevant</i>	<u>Blood Safety Entry</u>
<i>Reason for change</i>	This entry was updated to support the implementation of the recommendations from the FAIR study; the deferral period after sexual contact with an infected person, has been reduced to three months.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 57

Tendonitis

<i>Discretionary</i>	If the donor is only taking nonsteroidal anti-inflammatory drugs, accept.
<i>See if Relevant</i>	<u>Disabled Donor</u> <u>Drug Index - preparations which may affect platelet function</u> <u>Infection - General</u> <u>Nonsteroidal Anti-Inflammatory Drugs</u> <u>Steroid Therapy</u> <u>Surgery</u>
<i>Additional Information</i>	This entry includes inflammatory conditions affecting tendons, their sheaths and bursas. Treatment may be with rest, nonsteroidal anti-inflammatory drugs (these affect platelet function), steroid injections or tablets and surgery.
<i>Reason for change</i>	A 'Discretionary' entry has been added. This entry has been expanded to include additional relevant links and additional information.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 01.

Tetanus

1. Affected Individual

<i>Obligatory</i>	1. If treated with immunoglobulin: See: <u>Immunoglobulin Therapy</u> 2. Must not donate if: Not fully recovered
<i>Discretionary</i>	If fully recovered and is acceptable according to immunoglobulin therapy advice, accept.

<i>See if Relevant</i>	<u>Wounds, Mouth and Skin Ulcers</u>
<i>Additional Information</i>	Tetanus is a severe illness and usually requires treatment with high dose immunoglobulin. This may exclude the individual from donation.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 03 Issue 01.
<i>Reason for change</i>	Additional link added.

2. Immunisation

<i>Obligatory</i>	Must not donate if: Less than four weeks from exposure to a tetanus risk injury or receipt of passive immunisation with tetanus immunoglobulin.
<i>Discretionary</i>	a) If not exposed i.e. prophylactic tetanus toxoid Immunisation only or a tetanus toxoid booster, accept. b) If treated with single dose anti-tetanus immunoglobulin (intra muscular) and more than four weeks from exposure, accept.
<i>See if Relevant</i>	<u>Immunoglobulin Therapy</u> <u>Wounds, Mouth and Skin Ulcers</u>
<i>Additional Information</i>	Active or passive immunisation may mask infection. It is important to wait four weeks to ensure that the potential donor is not infected.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 03 Issue 01.
<i>Reason for change</i>	Additional link has been added.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 03 Issue 01.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 03 Issue 01.

Threadworms

<i>Discretionary</i>	Even if on treatment, accept.
<i>Additional Information</i>	Threadworms are a common problem in children but can also infect adults. The infection is usually harmless and should not affect fitness to donate.
<i>Reason for change</i>	'Additional Information' has been added.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 01.

Thrombosis and Thrombophilia

<i>Definitions</i>	Thrombophilia: This is a condition in which there is an increased tendency for blood clots to form. It is often inherited and may be discovered through family studies. Not all individuals with a thrombophilic condition will suffer from blood clots.
<i>Obligatory</i>	For Acquired Thrombophilia, see: Is there a specific A-Z entry for the underlying cause? Must not donate if: a) Due to atherosclerosis (e.g. coronary thrombosis). b) Two or more episodes of thrombosis requiring treatment.

- c) Less than seven days after completing anticoagulant therapy.
- d) Has thrombophilia and has had one or more episodes of thrombosis.
- e) Has thrombophilia associated with a history of recurrent superficial thrombophlebitis.
- f) History of Vaccine Induced Thrombotic Thrombocytopenia (VITT), Thrombotic Thrombocytopenic Purpura (TTP) or Heparin Induced Thrombocytopenia (HIT).

Discretionary

- a) If a first episode of thrombosis, such as deep vein thrombosis (DVT), retinal vein thrombosis or pulmonary embolism (PE):
- If no underlying cause that excludes the donor has been identified, and
 - The donor is not known to have thrombophilia, and
 - The donor is well and anticoagulant therapy (if used) has been stopped for at least seven days,

accept.

- b) If the potential donor has thrombophilia and,
- The donor is not on antithrombotic therapy, and
 - The donor has never had an episode of thrombosis, and
 - The donor has not been treated with antithrombotic therapy for recurrent pregnancy loss, and
 - The donor has never been treated with plasma-derived clotting factor concentrates, and
 - If relevant, the underlying cause of an acquired thrombophilia (see additional information) does not exclude the donor,

accept.

- c) If the potential donor has a history of Axillary Vein Thrombosis, **refer to a DCSO**.

- d) If the donor has a history of superficial thrombophlebitis (superficial vein thrombosis) see Superficial Thrombophlebitis

See if Relevant

Anticoagulant Therapy
Autoimmune Disease
Cardiovascular Disease
Drug Index - preparations which may affect platelet function
Malignancy
Nonsteroidal Anti-Inflammatory Drugs
Superficial Thrombophlebitis

Additional Information

Thrombophilia is a broad medical term which describes a multifactorial condition where the blood has an increased tendency to clot. Individuals with thrombophilia can present with arterial or venous thrombosis. The causes of thrombophilia include inherited and acquired disorders, and a combination of causes may be present.

Inherited causes of thrombophilia may be discovered through family testing. These include:

- Antithrombin, Protein C and Protein S deficiency
- Factor V Leiden and prothrombin gene mutations

Acquired causes of thrombophilia may present later in life and can be associated with:

- Malignancy including myeloproliferative neoplasms
- Antiphospholipid syndrome and other autoimmune connective tissue disorders. These may be associated with a lupus anticoagulant and/or anti-cardiolipin antibodies on laboratory testing.

Retinal Vein Thrombosis (also known as Retinal Vein Occlusion) is a form of retinal vascular disease and can affect central or branch retinal veins. The condition is uncommon under the age of 60 but becomes more frequent in later life. The condition may be associated with risk factors including hypertension, hyperlipidaemia, diabetes mellitus, atherosclerosis, and smoking.

VITT, TTP and HIT are rare disorders characterised by arterial or venous thrombosis in combination with a low platelet count (due to platelet consumption). Donors who recover from these disorders are unlikely to be eligible to donate due to the therapy they received (e.g. the primary treatment for TTP is plasma exchange with FFP) or an underlying condition (e.g. the indication for Heparin therapy that triggered HIT). VITT was recognised as a complication of some SARS-CoV-2 (COVID-19) vaccinations.

Axillary Vein Thrombosis can be precipitated by excessive use of the arm (e.g. sports or working above head level) but other precipitants include venous compression in thoracic outlet syndrome, diabetes, smoking, malignancy and venous cannulation. The donor may be eligible to donate if the underlying cause has been identified and corrected, but this should be balanced with the remote risk of local complications from a subsequent donation.

Superficial thrombophlebitis, also known as superficial vein thrombosis, is a common condition usually, but not exclusively, affecting the lower limbs. It is characterised by inflammation in a superficial vein associated with clot formation. This is different to, and less serious than, a deep vein thrombosis (DVT). If the superficial clot extends to where the superficial and deep veins join, a DVT can develop. Superficial thrombophlebitis normally settles within two to six weeks. Some individuals may be treated with anticoagulants to reduce the risk of extension. Recurrent superficial thrombophlebitis is sometimes associated with a diagnosis of thrombophilia.

<i>Information</i>	Part of this entry is a requirement of the Blood Safety and Quality Regulations 2005.
<i>Reason for change</i>	Revision of guidance and information for superficial thrombophlebitis. Addition of link to the Superficial Thrombophlebitis entry.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 68

Thrush

Also Known As Candida, candidiasis, moniliasis or yeast.

1. Oral

Obligatory **Must not donate if:**

- Unexplained.
- Related to immunosuppression.
- Less than seven days after completion of any treatment.

See if Relevant Antibiotic Therapy
Asthma
Immunodeficiency

Additional Information Oral thrush is uncommon, unless there is an underlying illness, or there has been recent treatment with antibiotics. It may also be a problem in people using steroid inhalers or antiseptic mouthwashes.

If the infection returns quickly after stopping treatment, this is very suggestive of underlying immunosuppression. The individual should not be accepted unless they have been properly investigated and an underlying immunodeficiency has been excluded. This is why we require any treatment to have been stopped for at least seven days.

2. Genitoanal

Obligatory **Must not donate if:**

- Related to immunosuppression.
- Less than seven days after receiving systemic (oral) therapy.

<i>Discretionary</i>	If not related to immunodeficiency, even if using local therapy, accept.
<i>See if Relevant</i>	<u>Antibiotic Therapy</u> <u>Immunodeficiency</u>
<i>Additional Information</i>	Vaginal thrush is common and is not usually a sign of a more serious problem. Penile thrush is less common and is usually a problem in uncircumcised men. Both types of thrush can affect the whole of the perineal area including the anus. The yeast that causes thrush is usually present on everybody's skin so it should not normally be considered as a sexually transmitted infection. If the infection requires systemic (oral) treatment, and returns quickly after stopping this treatment, it is suggestive of underlying immunosuppression. The individual should not be accepted unless they have been properly investigated and an underlying immunodeficiency has been excluded. This is why we require any systemic treatment to have been stopped for at least seven days.
<i>Reason for change</i>	Link updated from 'Immunosuppression' to 'Immunodeficiency' in the 'See if Relevant' section.
<i>Update Information</i>	This entry was last updated in: WB-DSG Edition 203 Release 73

Thyroid Disease

<i>Obligatory</i>	Must not donate if: a) Under investigation. b) Malignant. c) Less than six months from treatment with radioactive iodine therapy. d) Less than 24 months from stopping treatment with anti-thyroid tablets. e) Less than 8 weeks since commencing thyroid replacement therapy (thyroxine).
<i>Discretionary</i>	If on stable maintenance thyroid replacement therapy (thyroxine) and there have been no dose changes in the last 4 weeks, accept.
<i>See if Relevant</i>	<u>Autoimmune Disease</u> <u>Malignancy</u> <u>Surgery</u>
<i>Additional Information</i>	An over or an under active thyroid increases the risk of heart disease. Treatments used to treat an overactive thyroid are potentially harmful to the unborn child of a transfused mother.
<i>Reason for change</i>	The acceptance criteria for donors on long term thyroxine has been reviewed.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 39.

Tissue and Organ Recipients

<i>Excludes</i>	Recipients of donated human eggs, sperm and embryos.
<i>Obligatory</i>	Must not donate if: 1. At any time has: a) Needed immunosuppression. b) Had dura mater transplanted.

- c) Had a stored autologous tissue, matrix implant or organ transplanted.
- d) Had ocular tissue transplanted.
- e) Had a Xenotransplant performed.

2. Since January 1st 1980:

Has had an allogeneic human tissue or organ transplant.

- Discretionary*
- a) If before January 1st 1980 an allogeneic tissue or organ transplant, other than those listed above, was performed and there is no other reason to exclude the donor, accept.
 - b) If at any time a non-stored autologous tissue or organ has been transplanted, accept.
 - c) If has received an acellular non-human matrix graft, accept.

See Surgery

See if Relevant Dental Treatment
Eye Disease
Immunodeficiency
Prion Associated Diseases
Transfusion
Trying to Conceive
Xenotransplantation

Additional Information The transfer of tissues or organs between individuals and species has led to the spread of infection. The above guidelines are intended to minimize these risks.

There is now a concern that this could also happen with vCJD. This is because in the autumn of 2003 a UK recipient of blood, taken from a healthy donor who later developed vCJD, died from vCJD. Since then, there have been several cases of infection with the vCJD prion in recipients of blood from donors who have later developed vCJD.

In view of this, people who have received a tissue or organ transplant since 1980, will be excluded from donation in the same way as recipients of transfusion are. This date is before BSE, which is believed to have caused vCJD, was prevalent.

Following an update to the SaBTO Microbiological Safety Guidelines, recipients of donated human eggs, sperm or embryos can be accepted to donate. Care should be taken to ensure they also meet the other criteria included in the 'Trying to Conceive' entry.

Stored autologous tissue has been replaced in the wrong individual. Because of the associated infection risk these donors are not allowed to donate. It is important to check that any tissue transplanted has not be stored (e.g. chondrocytes).

Information This entry reflects guidance from the former Committee on the Microbiological Safety of Blood Tissues and Organs of the Department of Health.

Reason for change Link updated from 'Immunosuppression' to 'Immunodeficiency' in the 'See if Relevant' section.

Update Information This entry was last updated in:
 WB-DSG Edition 203 Release 73

Topical Medication

Obligatory **Must not donate if:**
 a) The site of venepuncture is affected.

b) There is broken or infected skin

Discretionary If the condition being treated does not exclude, accept.

See if Relevant Acne
Alopecia
Dermatitis

Infection - GeneralPsoriasisSteroid Therapy

<i>Additional Information</i>	Any area of broken skin can be a means for bacterial entering the blood. This risk is higher if the venesection site is affected. Bacteria can be a serious threat to anybody receiving blood or blood components. This is because bacteria can multiply to dangerous levels after collection.
<i>Reason for change</i>	Appropriate links and 'Additional Information' have been added.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 01.

Toxoplasmosis

<i>Obligatory</i>	Must not donate if: Less than six months from recovery.
<i>Additional Information</i>	This is a common parasitic infection, often spread by cat faeces or eating undercooked meat. It can be spread through transfusion. It may have serious consequences or even prove fatal for the recipient. Usually it does not cause symptoms, as the body's immune system easily overcomes the parasite. If the infection has caused symptoms that has lead to it being diagnosed, waiting six months from recovery will make it unlikely that it will be passed on by donation.
<i>Information</i>	This is a requirement of the Blood Safety and Quality Regulations 2005.
<i>Reason for change</i>	Entry has been simplified following a risk assessment.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 12.

Transfusion

<i>Obligatory</i>	<p>1. Must not donate if: At any time the donor has:</p> <p>a) Received, or thinks they may have received, a transfusion of blood or blood components in a country endemic for malaria or South American trypanosomiasis.</p> <p>b) Received treatment with blood derived coagulation factor concentrates. This includes prothrombin complex to reverse over-anticoagulation.</p> <p>2. Must not donate if: Since January 1st 1980:</p> <p>a) Anywhere in the world the donor has received, or thinks they may have received, a transfusion with red cells, platelets, fresh frozen plasma (FFP), cryoprecipitate, cryodepleted plasma, convalescent plasma, granulocytes, buffy coat preparations, intravenous or subcutaneous human normal immunoglobulin. This includes mothers whose babies have required intra-uterine transfusion.</p> <p>b) Has had a plasma exchange performed.</p>
<i>Discretionary</i>	<p>1.</p> <p>a) If on medical inquiry it is unlikely that the donor has been transfused, accept.</p> <p>b) If treatment with human immunoglobulin has been limited to specific immunoglobulin given as prophylaxis (e.g. anti-D, anti-tetanus or hepatitis immunoglobulin etc.), accept.</p> <p>2. Autologous Transfusion in:</p> <ul style="list-style-type: none"> • the United Kingdom • North America • Australasia • Western Europe (at any time) • EU member states (from February 2005) <p>If only the donor's own blood has been used, accept.</p>

3. Donor transfused before 1st January 1980:

a) If before 1st January 1980 the donor received, or thinks they may have received, a transfusion in a country endemic for malaria or South American trypanosomiasis, check the Geographical Disease Risk Index. If transfused in an at-risk country and a validated malarial antibody test and/or (as appropriate) a validated test for *T. cruzi* antibody is negative, accept.

b) If the transfusion was not within a risk area for either malaria or South American trypanosomiasis, accept.

See if Relevant Bleeding Disorder
Geographical Disease Risk Index
Immunodeficiency
Immunoglobulin Therapy
Malaria
Prion Associated Diseases
South American Trypanosomiasis

Additional Information Transfused donors have previously contributed to the spread of some diseases. This happened with hepatitis C.

Transfusions in some countries may have put the donor at risk of malaria or South American trypanosomiasis. It is necessary to exclude these infections before accepting the donor.

Coagulation concentrates:

People who have received blood derived coagulation concentrates (these are made from the blood of many donors) may have been put at risk of infections that can be passed through blood.

Donors transfused since 1980:

In the autumn of 2003 a UK recipient of blood, taken from a healthy donor who later developed variant Creutzfeldt-Jakob Disease (vCJD), died from vCJD. Since then there have been several cases of infection with the vCJD prion in recipients of blood from donors who have later developed vCJD. In view of this, people transfused, or possibly transfused, since 1980 are now excluded from donation. This date is before Bovine Spongiform Encephalopathy, which is believed to have caused vCJD, was prevalent.

Plasma exchange results in a patient being exposed to multiple donors. In view of the increased vCJD risk, donations may not be taken from individuals who have had a plasma exchange performed since 1980.

Donors transfused before 1996 (Infected Blood Inquiry):

The Infected Blood Inquiry (2024) recommended that anyone who received a transfusion before 1996 should be offered hepatitis C testing, unless they have already been tested.

Advise the donor to discuss/seek testing for hepatitis C with their GP or another clinical service if:

- they were transfused before 1996, and
- they are currently deferred from donating blood, and
- they haven't been tested for hepatitis C by a health care provider since, and
- they haven't donated blood since the start of 1996.

Donors who are accepted to donate do not need to see their GP as their blood will be tested as part of routine blood donation screening.

Information This entry reflects guidance from SaBTO (Advisory Committee on the Safety of Blood, Tissues and Organs) and its predecessor, the Committee on the Microbiological Safety of Blood Tissues and Organs of the Department of Health.

Reason for change Addition of guidance for the management of previously-deferred donors, following the recommendation of the Infected Blood Inquiry (2024) regarding hepatitis C testing.

Update Information This entry was last updated in:
WB-DSG Edition 203 Release 75

Transgender and Non-Binary Individuals

<i>Definitions</i>	<p>Transgender and non-binary individuals</p> <p>Trans is an umbrella term to describe people whose gender is not the same as, or does not sit comfortably with, the sex they were assigned at birth. Trans people may describe themselves using one or more of a wide variety of terms including (but not limited to) transgender, non-binary or gender queer. Gender affirming hormone therapy may be used as part of transition by transgender and non-binary individuals.</p>
<i>Discretionary</i>	<p>a) If the donor is taking masculinising hormone therapy (e.g. testosterone) to support their transition, the donor is well and the donor has been on treatment for more than 12 months, accept.</p> <p>b) If the donor is taking feminising hormone therapy, and the donor is well, accept.</p>
<i>See if Relevant</i>	<p><u>Anti-Androgens</u> <u>Haemoglobin Estimation</u> <u>Hormone Replacement and Sex Hormone Therapy</u> <u>Surgery</u></p>
<i>Additional Information</i>	<p>The higher haemoglobin concentration of men, compared to women, is related to testosterone levels. Testosterone therapy will result in the haemoglobin concentration rising. The opposite will be true if a person is taking feminising therapy.</p> <p>Waiting 12 months after starting masculinising hormone therapy, ensures that donation does not interfere with the assessment and laboratory monitoring of their treatment. A high haemoglobin (polycythaemia) can be a complication of this therapy and blood donation may mean this complication is not recognised. Once treatment is stabilised, it may be appropriate to offer the donor an individualised haemoglobin screening range consistent with their therapy (e.g. Haemoglobin 135 to 180 g/L for donors taking testosterone). Donors must be counselled to inform the Blood Service if their treatment changes or discontinues.</p> <p>As well as hormones, donors may take other medication to modify the effect of sex hormones as part of gender-affirming treatment. This may include hormone blockers, such as anti-androgens, which could affect the donor's eligibility.</p> <p>For blood services that use leucocyte antibody screening as a TRALI risk reduction measure, donors who were assigned female at birth should be included.</p>
<i>Reason for change</i>	Entry title changed. Instruction to base haemoglobin screening on gender removed. Addition of guidance re gender-affirming hormone therapy.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 70

Travel

<i>See if Relevant</i>	<p><u>Air Crew and Air Traffic Controllers</u> <u>Geographical Disease Risk Index</u> <u>Hazardous Activity</u> <u>Infection - General</u> <u>Malaria</u> <u>South American Trypanosomiasis</u></p>
<i>Additional Information</i>	<p>Donating before or after travel should not be a problem provided the donor is well hydrated. Travelling, particularly by plane, can be dehydrating and this may increase the risk of developing a thrombosis. If the donor is dehydrated, they should be advised to delay donating until they are well hydrated to avoid an increased risk of fainting.</p> <p>If the donor is likely to be exercising in conditions where the amount of available oxygen is low (e.g. at high altitude) it may be sensible to delay donation within two weeks of travel, so as to avoid the possibility of increasing the risk of adverse events.</p>
<i>Reason for change</i>	The See if Relevant section has been revised.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 66

Tropical Viruses

<i>Includes</i>	Chikungunya Virus, also known as CHIKV Dengue Virus, also known as Dengue Fever Yellow Fever, also known as YF Zika Virus, also known as ZIKV, and Zika Virus Fever
<i>Excludes</i>	Donors who will only donate plasma for fractionation. See <u>Tropical Viruses - plasmapheresis donors</u>
<i>Definitions</i>	Tropical Virus Endemic Areas: are shown in the 'Geographical Disease Risk Index' (GDRI) as a Tropical Virus Risk.
<i>Obligatory</i>	<p>Must not donate if:</p> <p>a) It is less than six months from a donor's return from a Tropical Virus Risk endemic area and the donor has been diagnosed with Chikungunya, Dengue, Yellow Fever or Zika virus infection whilst there or following their return to the UK.</p> <p>b) It is less than six months from a donor's return from a Tropical Virus Risk endemic area and the donor has either had a history of symptoms suggestive of Chikungunya, Dengue, Yellow Fever or Zika virus infection whilst there or following their return to the UK.</p> <p>c) In other cases it is less than four weeks from a donor's return from a Tropical Virus Risk endemic area.</p>
<i>Discretionary</i>	All donors may be accepted six months after their return from an affected area or resolution of symptoms. This may be reduced to four weeks, if they have had neither symptoms nor evidence of infection.
<i>See if Relevant</i>	<u>Infection - General</u> <u>Malaria</u> <u>South American Trypanosomiasis</u> <u>The 'Geographical Disease Risk Index'</u>
<i>Additional Information</i>	<p>Chikungunya, Dengue, Yellow Fever and Zika virus are spread by the day-flying mosquito species <i>Aedes aegypti</i> and <i>Aedes albopictus</i>. As these mosquitoes are typically found in tropical and subtropical regions, the main geographical areas affected by tropical virus infection are the Caribbean, South and Central America, Mexico, Africa, the Pacific Islands, Southeast Asia, Indian sub-continent, Hawaii and northern parts of Australia. The range of <i>Aedes albopictus</i> is also increasing into more temperate zones leading to outbreaks of tropical virus disease in new areas. There have been outbreaks of Dengue and Chikungunya in parts of Europe.</p> <p>Chikungunya is an alpha virus that can cause a wide spectrum of disease. This may range from no or minimal symptoms to death. Most commonly it causes arthritis (typically in the knee, ankle and small joints of the extremities), high fever and a maculopapular rash.</p> <p>Chikungunya virus is found in countries in Asia, Africa, Central and South America, and in the islands of the Caribbean. There is no evidence of person-to-person transmission except through blood transfer. Transfusion-transmission from an asymptomatic individual has not been documented. Nevertheless, restrictions after travel to a Chikungunya virus risk area were introduced to reduce any risk of transmission through blood or tissue donation.</p> <p>Dengue Virus is a flavivirus that typically gives rise to abrupt high fever with a range of accompanying symptoms.</p> <p>Dengue fever (DF) is the most common insect-borne disease worldwide. Dengue is currently considered endemic in approximately 140 countries. Transfusion-transmission has been reported.</p> <p>Overall, up to 75% of cases are asymptomatic or mild. If symptoms occur, they can range from non-specific acute febrile illness to severe disease including dengue haemorrhagic fever and dengue shock syndrome. Mild cases may be misdiagnosed as other febrile illnesses.</p> <p>Yellow Fever Virus is a flavivirus which is found in Africa, South America, Central America and parts of the Caribbean. Symptoms of Yellow Fever include high temperature, headache, nausea and vomiting, muscle pains and backache. One in four individuals may suffer from jaundice and bleeding from the gastrointestinal tract and other sites.</p>

Zika Virus is a flavivirus which was known to occur in Africa and parts of Southeast Asia. More recently, Zika Virus has been associated with epidemic outbreaks in the Pacific region and in the Americas. As well as mosquito-borne infection,

Zika Virus can be spread through sexual transmission. Infection is usually asymptomatic or presents as a mild self-limiting febrile illness. More severe disease and hospitalisation are rare but infection during pregnancy carries a high risk of congenital abnormalities in the baby. Zika Virus infection may be mistaken for Chikungunya or Dengue infections as these viruses often co-circulate.

Position statements are available in the JPAC Document Library.

<i>Information</i>	This entry is compliant with the Blood Safety and Quality Regulations 2005.
<i>Reason for change</i>	Entry updated to exclude donors who will only donate plasma for fractionation.
<i>Update Information</i>	This entry was last updated in: WB-DSG Edition 203 Release 71

Tropical Viruses - plasmapheresis donors

For donors who will donate whole blood, platelets and other cellular components see Tropical Viruses

<i>Includes</i>	Chikungunya Virus, also known as CHIKV Dengue Virus, also known as Dengue Fever Yellow Fever, also known as YF Zika Virus, also known as ZIKV, and Zika Virus Fever
<i>Excludes</i>	This entry only applies for donors who will only donate plasma for fractionation. It should not be used for donors who will donate whole blood, platelets and other cellular components.
<i>Definitions</i>	Tropical Virus Endemic Areas are shown in the ' <u>Geographical Disease Risk Index</u> ' (GDRI) which includes details of the specific viral risks present.
<i>Obligatory</i>	Must not donate if: a) It is less than six months from a donor's return from a Yellow Fever risk area and the donor has been diagnosed with Yellow Fever whilst there or following their return to the UK. b) It is less than six months from a donor's return from a Tropical Virus Risk endemic area and the donor has either had a history of symptoms suggestive of Yellow Fever whilst there or following their return to the UK. c) The donor was diagnosed with Chikungunya Virus, Dengue Virus or Zika Virus infection and the donor has not fully recovered from their illness.
<i>Discretionary</i>	If the donor has returned from a Chikungunya Virus, Dengue Virus or Zika Virus risk area and the donor is well, accept. If it more than 28 days since the donor has returned from a Yellow Fever risk area, and the donor has been well while there and after their return to the UK, accept.
<i>See if Relevant</i>	<u>Infection – General</u> <u>Malaria</u> <u>South American Trypanosomiasis</u> <u>The 'Geographical Disease Risk Index'</u>
<i>Additional Information</i>	Chikungunya, Dengue, Yellow Fever and Zika virus are spread by the day-flying mosquito species <i>Aedes aegypti</i> and <i>Aedes albopictus</i> . As these mosquitoes are typically found in tropical and subtropical regions, the main geographical areas affected by tropical virus infection are the Caribbean, South and Central America, Mexico, Africa, the Pacific Islands, Southeast Asia, Indian sub-continent, Hawaii and northern parts of Australia. The range of

Aedes albopictus is also increasing into more temperate zones leading to outbreaks of tropical virus disease in new areas. There have been outbreaks of Dengue and Chikungunya in parts of Europe.

Chikungunya is an alpha virus that can cause a wide spectrum of disease. This may range from no or minimal symptoms to death. Most commonly it causes arthritis (typically in the knee, ankle and small joints of the extremities), high fever and a maculopapular rash.

Chikungunya virus is found in countries in Asia, Africa, Central and South America, and in the islands of the Caribbean. There is no evidence of person-to-person transmission except through blood transfer. Transfusion-transmission from an asymptomatic individual has not been documented. Nevertheless, restrictions after travel to a Chikungunya virus risk area were introduced to reduce any risk of transmission through blood or tissue donation.

Dengue Virus is a flavivirus that typically gives rise to abrupt high fever with a range of accompanying symptoms. Dengue fever (DF) is the most common insect-borne disease worldwide. Dengue is currently considered endemic in approximately 140 countries. Transfusion-transmission has been reported.

Overall, up to 75% of cases are asymptomatic or mild. If symptoms occur, they can range from non-specific acute febrile illness to severe disease including dengue haemorrhagic fever and dengue shock syndrome. Mild cases may be misdiagnosed as other febrile illnesses.

Yellow Fever Virus is a flavivirus which is found in Africa, South America, Central America and parts of the Caribbean. Symptoms of Yellow Fever include high temperature, headache, nausea and vomiting, muscle pains and backache. One in four individuals may suffer from jaundice and bleeding from the gastrointestinal tract and other sites.

Zika Virus is a flavivirus which was known to occur in Africa and parts of Southeast Asia. More recently, Zika Virus has been associated with epidemic outbreaks in the Pacific region and in the Americas. As well as mosquito-borne infection, Zika Virus can be spread through sexual transmission. Infection is usually asymptomatic or presents as a mild self-limiting febrile illness. More severe disease and hospitalisation are rare but infection during pregnancy carries a high risk of congenital abnormalities in the baby. Zika Virus infection may be mistaken for Chikungunya or Dengue infections as these viruses often co-circulate.

The processes used to fractionate plasma include several measures that inactivate or remove viruses. This means that some travel risks described in the GDRI do not need to be applied for donors who will only donate plasma for fractionation.

<i>Reason for change</i>	This is a new entry.
<i>Update Information</i>	This entry was last updated in: WB-DSG Edition 203 Release 71

Tuberculin PPD Test

<i>Obligatory</i>	Must not donate unless: No further investigations or treatment is planned
<i>See if Relevant</i>	<u>Sarcoidosis</u> <u>Tuberculosis</u>
<i>Additional Information</i>	The tuberculin PPD Test, sometimes known as a Mantoux test, is used to test for exposure to Tuberculosis, or to see if past immunisation with BCG remains effective. It may also be used as part of the investigation of sarcoidosis.
<i>Reason for change</i>	This is a new entry, replacing the previous entries for the Heaf test (now discontinued) and the Mantoux test.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 01.

Tuberculosis

Also Known As TB.

1. Affected Individual

Obligatory **Must not donate if**
 a) Infected.
 b) Less than 24 months from confirmation of cure.
 c) Under follow-up.

Discretionary Donors with a diagnosis of Latent TB can donate, as long as they are not currently undergoing investigation or treatment.

Donors on antibiotic treatment for Latent TB only can donate 7 days after their last dose.

See if Relevant For BCG immunization:
Immunisation - Live
Tuberculin PPD Test

Reason for change Advice and background information on Latent TB has been added.

2. Contact

Obligatory **Must not donate until:**
 Screened and cleared.

Discretionary If the donor has been informed that they do not need to be screened, accept.

See if Relevant For BCG immunization:
Immunisation - Live
Tuberculin PPD Test

Additional Information Close contacts may have undiagnosed disease.

Reason for change The links and 'Additional Information' have been updated.

Update Information This entry was last updated in:
 DSG-WB Edition 203, Release 55

Turner's Syndrome

Discretionary If in good health, accept.

See if Relevant Cardiovascular Disease
Kidney and Bladder Disease

Additional Information Turner's syndrome is a chromosomal abnormality that occurs in about one in 2,000 female births. There may be associated problems affecting the cardiovascular and renal systems that should be enquired for, as they may affect donor safety.

Reason for change Relevant links and 'Additional Information' has been added.

Update Information This entry was last updated in:
 DSG-WB Edition 203, Release 22.

Urinary Catheterisation

<i>Includes</i>	Self-catheterisation, indwelling urinary catheter
<i>Obligatory</i>	<p>Must not donate if:</p> <p>a) Has an indwelling urinary catheter.</p> <p>b) It is less than seven days since catheterisation.</p>
<i>Discretionary</i>	If it is seven days or more since catheterisation, there are no symptoms suggestive of urinary tract infection and the underlying condition does not prevent donation, accept.
<i>See if Relevant</i>	<u>Kidney and Bladder Disease</u>
<i>Additional Information</i>	<p>Self-catheterisation is usually needed regularly every day for bladder emptying.</p> <p>Catheterisation including self-catheterisation is also used to administer drugs directly into the bladder. It is important to ensure that the underlying condition requiring this treatment does not prevent donation.</p> <p>Catheterisation is likely to cause bacteraemia following the procedure. Bacteria can be a serious threat to anybody receiving blood or blood components. This is because they can multiply to dangerous levels after collection. The waiting time after catheterisation is to allow any bacteria that have entered the blood stream to be cleared.</p> <p>Indwelling urinary catheters are associated with ongoing, sometimes asymptomatic, urinary infection with the associated risk of bacteraemia. The underlying condition may prevent donation.</p>
<i>Reason for change</i>	<p>Title changed to allow inclusion of guidance for individuals with indwelling catheters as well as those who undertake self-catheterisation.</p> <p>Discretionary guidance to be able to accept some donors who require catheterisation periodically has been added, and additional information regarding this has been included.</p> <p>A link to Kidney and Bladder Disease has been added.</p>
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 61.

Valproate and Topiramate

<i>Obligatory</i>	Must not donate.
<i>Discretionary</i>	If it is more than seven days since the last dose of valproate or topiramate, and the reason for therapy does not preclude donation, accept.
<i>Additional Information</i>	<p>Sodium valproate (Epilim[®], Episenta[®]) and the related drugs valproic acid (Depakote[®], Convulex[®]) are anticonvulsant medications used in the treatment of epilepsy. They are also used for treatment of bipolar disorder and as prophylaxis for migraine.</p> <p>Topiramate (Topamax[®]) is an anticonvulsant medication used in the treatment of epilepsy and as prophylaxis for migraine.</p> <p>Exposure during pregnancy to either valproate or topiramate is a known cause of birth defects. As it is not possible to know whether an individual donation will be transfused to a pregnant woman, donors taking any form of sodium valproate, valproic acid or topiramate must be deferred until at least one week after stopping treatment.</p>
<i>Reason for change</i>	A deferral has been added for donors taking topiramate. The name of the entry has been updated to reflect this.
<i>Update Information</i>	This entry was last updated in: WB-DSG Edition 203 Release 74

Varicose Veins and Chronic Venous Insufficiency

<i>Obligatory</i>	<p>Must not donate if:</p> <p>a) The donor has inflamed, broken or ulcerated skin.</p> <p>b) The donor has superficial thrombophlebitis.</p> <p>c) The donor has chronic venous insufficiency associated with persistent skin changes affecting skin integrity.</p> <p>d) It is within one week of treatment by injection (sclerotherapy).</p> <p>e) The donor has had laser therapy which has not yet healed.</p>
<i>Discretionary</i>	<p>If fully recovered from any non-surgical treatment and at least seven days after sclerotherapy, accept.</p> <p>For surgical treatment, refer to the <u>Surgery</u> guideline.</p>
<i>See if Relevant</i>	<p><u>Laser Treatment</u></p> <p><u>Phlebitis</u></p> <p><u>Superficial Thrombophlebitis</u></p> <p><u>Surgery</u></p> <p><u>Thrombosis and Thrombophilia</u></p> <p><u>Wounds, Mouth and Skin Ulcers</u></p>
<i>Additional Information</i>	<p>Varicose veins are not a reason for deferral. However if there is active inflammation, thrombosis or ulceration, the donor should be deferred. This is to minimise any risk of thrombosis in the donor or transfer of infection to the recipient.</p> <p>There are many treatments for varicose veins, including sclerotherapy (injection of material to block the vein), endothermal or laser ablation, and surgery to remove affected veins. Newer treatments may also be available. It is important that the donor is fully recovered from any treatment.</p> <p>Chronic Venous Insufficiency is a condition where the normal flow of blood from the lower limbs back to the heart is impaired, leading to pooling of blood (stasis) in the legs. It can give rise to persistent inflammation and ulceration of the skin. If this occurs, it is a blood safety risk, due to breach of the normal skin barriers to bacterial infection.</p>
<i>Reason for change</i>	<p>Addition of chronic venous insufficiency to the title.</p> <p>New guidance for donors with persistent skin damage relating to venous insufficiency.</p> <p>Clarification of deferral requirements after treatment.</p> <p>Revision of See if Relevant section to reflect other DSG changes.</p>
<i>Update Information</i>	<p>This entry was last updated in: DSG-WB Edition 203, Release 68</p>

Vertigo

<i>Obligatory</i>	<p>Must not donate if:</p> <p>Experiencing dizzy spells.</p>
<i>Discretionary</i>	<p>If the donor has Meniere's disease, if well on the day, even if on treatment to prevent attacks, accept.</p>
<i>See if Relevant</i>	<p><u>Infection - General</u></p>
<i>Additional Information</i>	<p>Vertigo is a feeling of everything spinning around. It can be accompanied by nausea and sickness and lead to the affected person falling. There are many different causes and, if known, the cause should be looked up in the index.</p>

Because faintness after donation can cause similar symptoms it is recommended that people affected by vertigo should only donate if they are not experiencing any symptoms.

<i>Reason for change</i>	This is a new entry.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 203, Release 01.

Viral Haemorrhagic Fever

Definitions Includes Crimean-Congo Fever, Ebola Virus Disease, Lassa Fever and Marburg Fever.

Viral Haemorrhagic Fever Endemic Areas are shown in the 'Geographical Disease Risk Index' (GDRI) as a Viral Haemorrhagic Fever Risk. Outbreak information is also listed but is not required for Whole Blood and Components Donor Selection Guideline users.

1. Affected Individual

Obligatory **Must not donate if:**
Ever diagnosed with a Viral Haemorrhagic Fever.

2. Contact or traveller to endemic country

Obligatory **Must not donate if:**
Less than 6 months from last contact with an affected individual or travel to an endemic country.

Discretionary If more than 6 months from last contact, completion of investigations or return to the UK from endemic country, accept.

3. Sexual Partners of Affected Individuals

Obligatory **Must not donate if:**
The donor has had sex with an individual who had been diagnosed with a Viral Haemorrhagic Fever at any time before their last sexual contact.

See if Relevant The Geographical Disease Risk Index for countries with a current endemic Viral Haemorrhagic Fever risk

Additional Information There is evidence of persistent virus in individuals who recover from several forms of Viral Haemorrhagic Fever. For this reason, it is necessary to defer the sexual partners of these individuals.

Reason for change A permanent deferral has been introduced for donors who have had sex with an individual who has been diagnosed with a Viral Haemorrhagic Fever.

Update Information This entry was last updated in:
DSG-WB Edition 203, Release 55.

Vitamins and Other Nutritional Supplements

Obligatory **Must not donate if:**
On prescribed medication to treat a deficiency.

Discretionary a) Medication to prevent recurrence, as opposed to treat a deficiency (e.g. B12 for treated pernicious anaemia or folic acid for treated folate deficiency), accept.
b) If on oral self-medication, accept.

c) If on Vitamin D supplement to treat risk of vitamin D deficiency, accept.

See if Relevant Anaemia
Osteopenia

<i>Additional Information</i>	<p>People who are on treatment to cure a vitamin or other nutritional deficiency other than Vitamin D supplementation to prevent or treat osteopenia should not donate, even if they pass the haemoglobin-screening test.</p> <p>Once treatment is completed, even if they then require maintenance treatment, they should be accepted or excluded on the basis of the underlying condition that required treatment. As an example, a person with pernicious anaemia (vitamin B12 deficiency) should not be accepted until their anaemia is fully corrected. Once fully recovered, they may be accepted, even though receiving maintenance treatment to prevent recurrence.</p> <p>Vitamins and other nutritional supplements are often prescribed to prevent deficiency. For example, this might be for coeliac disease or for people wanting to conceive. Providing any underlying condition is not a reason to exclude the donor, they should be accepted.</p>
<i>Reason for change</i>	<p>Advice about Vitamin D supplementation and a link to osteopenia has been added, see the letter from UK Chief Medical Officers of UK Feb 2012: www.gov.uk/government/publications/vitamin-d-advice-on-supplements-for-at-risk-groups</p>
<i>Update Information</i>	<p>This entry was last updated in: DSG-WB Edition 203, Release 22.</p>

Warts

<i>Includes</i>	Molluscum contagiosum and verrucas.
<i>Obligatory</i>	Must not donate if: Treatment has left unhealed areas.
<i>Discretionary</i>	If there are no open wounds, even if on treatment, accept.
<i>See if Relevant</i>	<u>Sexually Transmitted Disease</u> <u>Surgery</u> <u>Wounds, Mouth and Skin Ulcers</u>
<i>Additional Information</i>	<p>Warts (including verruca) are caused by infection with the human papilloma virus (HPV) of which there are over 100 different types. They may occur on the skin and mucous membranes. The virus is spread by skin to skin contact and it can be very infectious. Genital warts are possibly the commonest sexually transmitted disease but they do not necessarily indicate high risk sexually activity, so no specific deferral is required.</p> <p>Molluscum contagiosum is also caused by a virus and can be managed in the same way as warts.</p> <p>Treatment may lead to unhealed wounds or sores and these pose a risk for bacteria entering the blood. Bacteria can be a serious threat to anybody receiving blood or blood components. This is because bacteria can multiply to dangerous levels after collection.</p>
<i>Reason for change</i>	This entry was revised to support the implementation of recommendations from the FAIR study; the instruction to discuss the possibility of high risk sexual activity has been removed.
<i>Update Information</i>	<p>This entry was last updated in: DSG-WB Edition 203, Release 57</p>

West Nile Virus

<i>Excludes</i>	Donors who will only donate plasma for fractionation. See West Nile Virus - plasmapheresis donors
<i>Definitions</i>	West Nile Virus (WNV) Endemic Areas: These are shown in the ' Geographical Disease Risk Index ' (GDRI).
<i>Obligatory</i>	Must not donate if: a) It is less than six months from a donor's return from a WNV endemic area and the donor has been diagnosed with WNV whilst there or following their return. b) It is less than six months from a donor's return from a WNV endemic area and the donor has either had a history of symptoms suggestive of WNV whilst there or within 28 days of their return. c) In other cases it is less than four weeks from a donor's return from a WNV endemic area.
<i>Discretionary</i>	1) All donors may be accepted six months after their return from an affected area. This may be reduced to four weeks if they have had neither symptoms nor evidence of infection. For donors who have been back in the UK for less than four weeks, who have not been diagnosed with WNV infection and who have not had symptoms suggestive of WNV infection, if a validated NAT for WNV is to be undertaken on the donated component(s), accept. 2) Donors who have been back in the UK for less than six months, who have had symptoms suggestive of WNV infection while abroad or within 28 days of return, (but no firm diagnosis of WNV infection) if a validated NAT for WNV is to be undertaken on the donated component (s), accept.
<i>See if Relevant</i>	The 'Geographical Disease Risk Index'
<i>Additional Information</i>	West Nile Virus is a flavivirus, similar to Dengue, which causes a wide spectrum of infection. This may range from no or minimal symptoms to death. It is geographically widespread, including areas in Europe and other parts of the world not affected by Malaria, and it has reached epidemic proportions in North America in recent years. There it has caused illness and death post transfusion and post transplantation of tissues and organs. It is spread by mosquitoes and so is more prevalent at times of the year when mosquitoes are active. As the problem can vary both in relation to geography and time of the year it is not possible to state areas from which donors need to be deferred and dates of disease activity. These are provided in the ' Geographical Disease Risk Index '. A ' Position Statement on West Nile Virus (WNV) ' is available in the 'Document Library' of ' www.transfusionguidelines.org '.
<i>Reason for change</i>	Entry updated to exclude donors who will only donate plasma for fractionation.
<i>Update Information</i>	This entry was last updated in: WB-DSG Edition 203 Release 71

West Nile Virus - plasmapheresis donors

For donors who will donate whole blood, platelets and other cellular components see [West Nile Virus](#)

<i>Excludes</i>	This entry only applies for donors who will only donate plasma for fractionation. It should not be used for donors who will donate whole blood, platelets and other cellular components.
<i>Definitions</i>	West Nile Virus (WNV) Endemic Areas: are shown in the 'Geographical Disease Risk Index' (GDRI).
<i>Obligatory</i>	Must not donate if:

The donor was diagnosed with West Nile Virus and the donor has not fully recovered from their illness.

Discretionary If the donor has returned from a WNV risk area and the donor is well, accept.

See if Relevant The 'Geographical Disease Risk Index'

Additional Information West Nile Virus is a flavivirus, similar to Dengue Virus, which causes a wide spectrum of infection. This may range from no or minimal symptoms to death. It is geographically widespread, including areas in Europe and other parts of the world not affected by Malaria, and it has reached epidemic proportions in North America in recent years. Mild cases may be misdiagnosed as other febrile illnesses.

The processes used to fractionate plasma include several measures that inactivate or remove viruses. This means that some travel risks described in the GDRI do not need to be applied for donors who will only donate plasma for fractionation.

Reason for change This is a new entry.

Update Information This entry was last updated in:
WB-DSG Edition 203 Release 71

Wounds, Mouth and Skin Ulcers

Obligatory **Must not donate if:**
a) Has infected wounds, or skin ulcers, sores or mouth ulcers.

b) Has persistently inflamed or broken skin, associated with cardiovascular disease, chronic venous insufficiency, lymphoedema, diabetes mellitus or other medical condition.

Discretionary If an individual has an uninfected wound or small non-infected aphthous ulcers only, accept.

See if Relevant Autoimmune Disease
Cardiovascular Disease
Diabetes Mellitus
Infection - General
Malignancy
Surgery
Tetanus - 2. Immunisation
Varicose Veins and Chronic Venous Insufficiency

Additional Information An infected wound, a sore or an ulcer is a risk for bacteria entering the blood. Bacteria can be a serious threat to anybody receiving blood or blood components. This is because bacteria can multiply to dangerous levels after collection.

A small individual aphthous ulcer in an otherwise healthy person does not pose such a risk. Donors with recurrent severe aphthous ulceration may have a serious underlying condition, such as an autoimmune disease.

Persistently inflamed or broken skin, usually of the lower limbs, is a complication of a range of medical conditions which affect the cardiovascular and/or lymphatic systems. There is an increased risk of bacterial contamination in a blood donation if the donor's normal skin integrity is impaired.

Reason for change Addition of guidance for donors with persistent damage to the skin of their lower limbs.

Update Information This entry was last updated in:
DSG-WB Edition 203, Release 68

Xenotransplantation

Includes Heterografts, non-human organ perfusion, xenografts and xenotransplant recipients.

Definitions **Xenotransplantation:**
Any procedure that involves the transplantation, implantation, or infusion into a human recipient of either (a) live cells, tissues, or organs from a non-human animal source, or (b) human body fluids, cells, tissues, or organs that have had ex vivo contact with live, non-human animal cells, tissues, or organs. Xenotransplantation products include live cells, tissues and organs.

Biological products, drugs, or medical devices sourced from non-living cells, tissues or organs from non-human animals including, but not limited to, porcine insulin, porcine heart valves and acellular porcine collagen matrix are not considered xenotransplantation products.

Inoculation injuries from non human sources are not considered to be Xenotransplants.

1. Recipient

Obligatory **Must not donate if:**
Material from a **living** non-human animal source has been directly or indirectly in contact with the donor's blood supply. This does not include animal bites.

See if Relevant Animal Bite (Non-Human)
Non-Consented Exposure to Human Body Fluids

Additional Information Exposure to non-human animal material, particularly when the person exposed is immunosuppressed, may result in unusual infections, that would not normally affect humans, being passed on to recipients of donated material. Inoculation injury, involving non-human animals, does not fall into the category of xenotransplantation

2. Current or Former Sexual Partner of Xenotransplant Recipient

Obligatory **Must not donate.**

Additional Information Sexual partners of individuals who have received a xenotransplant may potentially be at risk of acquiring an unusual infection that may be passed on by donated material. Because the duration of any risk is not known, deferral must be permanent.

Information This is a requirement of the Blood Safety and Quality Regulations 2005.

Reason for change Reference to specific products has been removed from the Definitions section

Update Information This entry was last updated in:
DSG-WB Edition 203, Release 10 Issue 67

XMRV

Discretionary Donors who have been tested positive for XMRV, accept.

Additional Information As there is no evidence that XMRV is implicated in human disease, a positive test is not a bar to blood donation.

Reason for change This is a new entry.

Update Information This entry was last updated in:
DSG-WB Edition 203, Release 10 Issue 01

Zanamivir

<i>Also Known As</i>	Relenza [®] .
<i>Obligatory</i>	Must not donate if: a) Taking zanamivir (Relenza [®]) as treatment for influenza. b) At any time in the seven days prior to, or while taking zanamivir, the donor has had symptoms of influenza, (a temperature of more than 38 degrees centigrade, or a history of fever and two or more of the following symptoms: cough, headache, runny nose, diarrhoea or vomiting).
<i>Discretionary</i>	If the potential donor is taking zanamivir as prophylaxis, they have not been advised to be confined to home, and have not had any symptoms of influenza, accept.
<i>See if Relevant</i>	<u>Infection - Acute</u>
<i>Additional Information</i>	Zanamivir is a viral neuraminidase inhibitor (neuraminidase is an enzyme that helps the virus spread from cell to cell). It is used to treat influenza and for post-exposure prophylaxis of influenza. It appears to be a very safe drug with little evidence for teratogenic (potential to cause birth defects) or mutagenic (potential to cause malignancy) effect.
<i>Information</i>	Part of this entry is a requirement of the Blood Safety and Quality Regulations 2005.
<i>Update Information</i>	This entry was last updated in: DSG-WB Edition 202, Release 13.

Updates

Specification of Current Version		
Publication	WB-DSG	
Edition	203	
Release	75	
Issue	01	17 February 2025

All changes to WB-DSG Edition 203 after Release 01

Release	Date	Change Notifications	
		Title	CN No.
75	17 February 2025	Hepatitis C testing of donors transfused before 1996 (IBI update)	<u>03 - 2025</u>
74	26 November 2024	Migraine and Topiramate	<u>51 - 2024</u>
		Monoclonal Antibody Therapies	<u>49 - 2024</u>
		Haemolytic Anaemia	<u>48 - 2024</u>
		Topical Anti-Androgens	<u>47 - 2024</u>
		COVID-19 Vaccinations	<u>46 - 2024</u>
73	07 October 2024	Links to Immunodeficiency entry in WB-DSG	<u>43 - 2024</u>
		Cerebrovascular Disease and Intracranial Haemorrhage	<u>30 - 2024</u>
		Platelet Disorders	<u>27 - 2024</u>
72	31 July 2024	Faints	<u>26 - 2024</u>
		Dental Treatment	<u>25 - 2024</u>
		Upper Age Limit for Returning Donors	<u>22 - 2024</u>
		Cardiovascular Disease	<u>21 - 2024</u>
		Bleeding Disorder	<u>20 - 2024</u>
		Pyrexia and Infection	<u>19 - 2024</u>
		Animal Bite (Non-Human)	<u>18 - 2024</u>
		Osteoarthritis	<u>17 - 2024</u>
		Drugs and Platelet Donation	<u>16 - 2024</u>
		Shunts, Stents and Devices	<u>15 - 2024</u>
Fertility	<u>14 - 2024</u>		
71	02 May 2024	Travel Criteria for Plasmapheresis Donors	<u>08 - 2024</u>
		Immunosuppression	<u>07 - 2024</u>

70	18 April 2024	Transgender and Non-Binary Donors	<u>09 - 2024</u>
		Hepatitis A and Parvovirus B19	<u>06 - 2024</u>
69	04 July 2023	Chronic Fatigue Syndrome	<u>29 - 2023</u>
		Blood Safety Entry	<u>28 - 2023</u>
		Air Crew and Air Traffic Controllers	<u>27 - 2023</u>
68	09 May 2023	Hepatitis B and Hepatitis of Unknown Cause	<u>13 - 2023</u>
		Tropical Viruses	<u>12 - 2023</u>
		Recurrent Thrombophlebitis	<u>11 - 2023</u>
		Frequency of Donation	<u>10 - 2023</u>
67	31 January 2023	Mpox (Monkeypox)	<u>04 - 2023</u>
		Donor Weight	<u>03 - 2023</u>
		Dental Treatment	<u>02 - 2023</u>
66	18 October 2022	Platelet Count	<u>59 - 2022</u>
		Liver Disease	<u>58 - 2022</u>
		Thrombosis and Thrombophilia	<u>57 - 2022</u>
		Air Crew and Air Traffic Controllers	<u>13 - 2023</u>
65	30 August 2022	Table of Immunisations	<u>49 - 2022</u>
		Hepatitis of Unknown Cause	<u>48 - 2022</u>
		Coronavirus Infection	<u>46 - 2022</u>
		Surgery	<u>45 - 2022</u>
		Sleep Apnoea	<u>44 - 2022</u>
		Addiction and Drug Abuse	<u>43 - 2022</u>
64	26 April 2022	Monkeypox	<u>40 - 2022</u>
		Hepatitis B	<u>39 - 2022</u>
64	31 May 2022	Body Piercing	<u>28 - 2022</u>
		Conn's Syndrome	<u>27 - 2022</u>
		Ehlers Danlos Syndrome	<u>26 - 2022</u>
		Narcolepsy	<u>25 - 2022</u>
		Appendix 4 - Management of post donation illness	<u>22 - 2022</u>
		Tropical Viruses	<u>21 -</u>

			<u>2022</u>
		Inflammatory Bowel Disease	<u>20 - 2022</u>
		Indwelling Shunts and Stents	<u>19 - 2022</u>
62	07 April 2022	Coronavirus Infection	<u>29 - 2022</u>
61	22 February 2022	Trying to Conceive	<u>08 - 2022</u>
		Non-Contagious Disease	<u>07 - 2022</u>
		Malignancy	<u>06 - 2022</u>
		Kidney & Bladder Disease	<u>05 - 2022</u>
		Complementary Therapy	<u>04 - 2022</u>
		Alopecia	<u>03 - 2022</u>
		Allergy	<u>02 - 2022</u>
60	20 January 2022	Coronavirus Infection	<u>01 - 2022</u>
59	21 December 2021	Blood Safety Entry	<u>45 - 2021</u>
58	22 June 2021	Plasma Changes	<u>15 - 2021</u>
57	26 May 2021	The FAIR Study	<u>16 - 2021</u>
56	05 May 2021	Hypercholesterolaemia	<u>08 - 2021</u>
		Diabetes Mellitus	<u>07 - 2021</u>
		Coronavirus Infection	<u>06 - 2021</u>
55	21 January 2021	Coronavirus Infection	<u>04 - 2021</u>
		Acitretin in Acne, Psoriasis and Skin Disease	<u>72 - 2020</u>
		Body Piercing	<u>71 - 2020</u>
		Drug Index	<u>70 - 2020</u>
		Animal Bites and Rabies	<u>69 - 2020</u>
		South American Trypanosomiasis	<u>58 - 2020</u>
		Viral Haemorrhagic Fever	<u>57 - 2020</u>
		Sodium Valproate	<u>56 - 2020</u>
		Tropical Viruses	<u>55 - 2020</u>
		Latent Tuberculosis	<u>54 - 2020</u>
		Haemochromatosis	<u>53 - 2020</u>
		Cervical Carcinoma in Situ	<u>52 - 2020</u>

		Clopidogrel	<u>51 - 2020</u>
54	16 December 2020	COVID-19 Vaccine	<u>73 - 2020</u>
53	13 November 2020	Clinical Trials	<u>65 - 2020</u>
		Coronavirus Infection	<u>64 - 2020</u>
52	21 September 2020	Transfusion	<u>48 - 2020</u>
51	08 June 2020	Coronavirus Infection	<u>29 - 2020</u>
50	01 June 2020	Northern Ireland - Donor Selection Changes 2020	<u>28 - 2020</u>
49	07 May 2020	Haemaglobin Estimation	<u>26 - 2020</u>
48	07 May 2020	Coronavirus Infection	<u>13 - 2020</u>
47	24 February 2020	Coronavirus Infection	<u>09 - 2020</u>
46	17 February 2020	Coronavirus Infection	<u>08 - 2020</u>
45	24 January 2020	Coronavirus Infection	<u>03 - 2020</u>
44	03 September 2019	Palpitations & Arrhythmias	<u>27 - 2019</u>
		Familial Psuedohyperkalaemia	<u>26 - 2019</u>
43	28 May 2019	HCV (Northern Ireland)	<u>11 - 2019</u>
		HTLV	<u>10 - 2019</u>
		Cervical Carcinoma in Situ	<u>09 - 2019</u>
		Blood Safety Entry (England, Scotland, Wales)	<u>08 - 2019</u>
		Blood Safety Entry (Northern Ireland)	<u>07 - 2019</u>
		Pre- and Post-Exposure Prophylaxis for HIV Prevention	<u>04 - 2019</u>
		Pregnancy	<u>03 - 2019</u>
		Malignancy	<u>02 - 2019</u>
		Hepatitis C	<u>01 - 2019</u>
42	17 September 2018	Narcolepsy	<u>21 - 2018</u>
		Hepatitis A	<u>20 - 2018</u>
41	10 May 2018	Hepatitis E	<u>17 - 2018</u>
40	27 November 2017	Surgery	<u>52 - 2017</u>
		Endoscopy	<u>51 - 2017</u>
		Bleeding Disorder	<u>49 - 2017</u>
		Syphilis	<u>47 -</u>

			<u>2017</u>
		Sex Worker	<u>45 - 2017</u>
		Non-Consented Exposure to Human Body Fluids	<u>44 - 2017</u>
		HTLV	<u>41 - 2017</u>
		Homosexual and Bisexual Individuals	<u>39 - 2017</u>
		HIV	<u>37 - 2017</u>
		Hepatitis C	<u>35 - 2017</u>
		Hepatitis B	<u>33 - 2017</u>
		Complementary Therapy	<u>31 - 2017</u>
		Body Piercing	<u>29 - 2017</u>
		Blood Safety Entry	<u>26 - 2017</u>
39	26 September 2017	Drug Index	<u>19 - 2017</u>
		Thyroid	<u>18 - 2017</u>
38	01 August 2017	Malaria	<u>15 - 2017</u>
		Tissue and Organ Recipients	<u>14 - 2017</u>
		Radiation Therapy	<u>13 - 2017</u>
		Eye Disease	<u>12 - 2017</u>
		Autoimmune Disease and Osteopenia	<u>11 - 2017</u>
37	21 February 2017	Hepatitis A	<u>04 - 2017</u>
		Central Nervous System Disease	<u>03 - 2017</u>
36	01 November 2016	Cardiac Surgery clarification	<u>37 - 2016</u>
35	01 September 2016	Blood Safety Entry & Homosexual and Bisexual Individuals	<u>36 - 2016</u>
34	23 August 2016	Tropical Viruses	<u>35 - 2016</u>
33	05 July 2016	Immunoglobulin	<u>21 - 2016</u>
		High Haemoglobin	<u>20 - 2016</u>
		Disabled Donor	<u>19 - 2016</u>
		Acne	<u>18 - 2016</u>
32	02 February 2016	Viral Haemorrhagic Fever	<u>15 - 2016</u>
		Tropical Viruses	<u>14 - 2016</u>
31	18 January 2016	Viral Haemorrhagic Fever	<u>11 - 2016</u>

		West Nile Virus	<u>09 - 2016</u>
		Tropical Viruses	<u>08 - 2016</u>
		Kidney and Bladder Disease	<u>06 - 2016</u>
		Memorial Tattoos and Body Piercing	<u>05 - 2016</u>
		Appendix 2 - Table of Immunisations	<u>04 - 2016</u>
		Hepatitis E	<u>03 - 2016</u>
		Hepatitis A	<u>02 - 2016</u>
		Glycogen Storage Disease	<u>01 - 2016</u>
30	06 October 2015	Chronic Fatigue Syndrome	<u>26 - 2015</u>
		Alopecia and Autoimmune Disease	<u>25 - 2016</u>
29	23 June 2015	Injectable Tanning Agents	<u>15 - 2015</u>
		<i>The changes made to the Blood Safety Entry have also been applied to the Blood and Tissue Safety Entry (Northern Ireland). Injectable Tanning Agents has been added to index and linked to Blood Safety Entry (England, Scotland and Wales).</i>	
		Complementary Therapy	<u>12 - 2015</u>
		Communication Difficulties	<u>11 - 2015</u>
		Central Nervous System	<u>10 - 2015</u>
28	17 March 2015	Nonsteroidal Anti-Inflammatory Drugs and Drug Index	<u>05 - 2015</u>
		Malignancy	<u>04 - 2015</u>
		Kidney and Bladder Disease	<u>03 - 2015</u>
		Asthma	<u>02 - 2015</u>
27	17 December 2014	Thrombosis	<u>48 - 2014</u>
		Respiratory Disease	<u>47 - 2014</u>
		Mental Health Problems	<u>46 - 2014</u>
		Tuberculosis	<u>45 - 2014</u>
		Hepatitis of Unknown Cause	<u>44 - 2014</u>
26	20 October 2014	Viral Haemorrhagic Fever Risk	<u>43 - 2014</u>
25	14 October 2014	West Nile Virus	<u>42 - 2014</u>
24	19 August 2014	Chikungunya Virus Risk	<u>41 - 2014</u>
23	08 July 2014	Chikungunya Virus Risk	<u>27 - 2014</u>
22	17 June 2014	Chikungunya Virus	<u>26 - 2014</u>

		Vitamins and Other Nutritional Supplements	<u>24 - 2014</u>
		Osteopenia	<u>23 - 2014</u>
		Kidney and Bladder Disease	<u>22 - 2014</u>
		Complementary Therapy	<u>21 - 2014</u>
		Central Nervous System Disease	<u>20 - 2014</u>
		Autoimmune Disease	<u>19 - 2014</u>
		Accept	<u>18 - 2014</u>
21	13 February 2014	Chikungunya Virus	<u>01 - 2014</u>
20	11 February 2014	Surgery	<u>22 - 2014</u>
		South American Trypanosomiasis	<u>21 - 2014</u>
		Skin Disease and Dermatitis (Alitretinoin)	<u>20 - 2014</u>
		Malignancy	<u>18 - 2014</u>
19	03 September 2013	West Nile Virus	<u>14 - 2013</u>
18	09 July 2013	Wounds, Mouth and Skin Ulcers	<u>07 - 2013</u>
		Cardiovascular Disease	<u>06 - 2013</u>
		Chest Pain	<u>05 - 2013</u>
		Chronic Infection	<u>03 - 2013</u>
		Hepatitis B	<u>02 - 2013</u>
17	04 June 2013	West Nile Virus	<u>01 - 2013</u>
16	05 February 2013	Compression Illness	<u>29 - 2012</u>
		Kidney Disease	<u>28 - 2012</u>
		Sickle-Cell Trait	<u>27 - 2013</u>
15	31 October 2012	West Nile Virus	<u>26 - 2012</u>
		West Nile Virus	<u>25 - 2012</u>
14	31 October 2012	Mobilised Granulocytes	<u>24 - 2012</u>
		Cardiovascular Disease	<u>23 - 2012</u>
		Cupping/Wet Cupping	<u>22 - 2012</u>
		West Nile Virus	<u>21 - 2012</u>
		Malaria	<u>20 - 2012</u>
13	28 August 2012	West Nile Virus	<u>19 -</u>

			<u>2012</u>
12	29 June 2012	Cardiac Surgery	<u>13 - 2012</u>
		Cardiovascular Disease	<u>12 - 2012</u>
		Skin Disease	<u>11 - 2012</u>
		Dermatitis	<u>10 - 2012</u>
		Pregnancy	<u>09 - 2012</u>
		Psoriasis	<u>08 - 2012</u>
		Toxoplasmosis	<u>07 - 2012</u>
		Acne	<u>06 - 2012</u>
11	28 March 2012	West Nile Virus	<u>03 - 2012</u>
10	24 January 2012	Hepatitis C	<u>26 - 2012</u>
		XMRV	<u>25 - 2012</u>
		Surgery	<u>24 - 2012</u>
		Cardiovascular Disease	<u>23 - 2012</u>
		Donor Weight	<u>22 - 2012</u>
09	07 November 2011	West Nile Virus	<u>19 - 2011</u>
		Homosexual and Bisexual Individuals	<u>17 - 2011</u>
		Blood Safety Entry	<u>16 - 2011</u>
08	04 October 2011	West Nile Virus	<u>18 - 2011</u>
07	31 August 2011	West Nile Virus	<u>15 - 2011</u>
06	08 August 2011	West Nile Virus	<u>11 - 2011</u>
05	07 June 2011	Porphyria	<u>05 - 2011</u>
		Syphilis	<u>04 - 2011</u>
		Immunisation - Live	<u>03 - 2011</u>
		Trying to Conceive	<u>02 - 2011</u>
		Pregnancy	<u>01 - 2011</u>
04	01 February 2011	Donor Weight and Donation Volumes	<u>13 - 2010</u>
03	01 November 2010	Chronic Fatigue Syndrome	<u>08 - 2010</u>
02	01 September 2010	West Nile Virus	<u>09 - 2010</u>
01	01 June 2010		

Appendix 1 - Estimated Blood Volume for Female donors (after Nadler) by height and weight

Estimated Blood Volume for Female donors (after Nadler) by height and weight																
Height cm	Weight Kg															Height 4' 10" or less
	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	
149 or less	Less than 3500 ml															
150	3039	3072	3105	3138	3171	3205	3238	3271	3304	3337	3370	3403	3436	3469	3502	4' 11"
151	3063	3096	3129	3163	3196	3229	3262	3295	3328	3361	3394	3427	3460	3493	3526	
152	3088	3121	3154	3187	3220	3253	3286	3319	3352	3386	3419	3452	3485	3518	3551	4' 11½"
153	3113	3146	3179	3212	3245	3278	3311	3344	3377	3410	3443	3477	3510	3543	3576	
154	3138	3171	3204	3237	3270	3303	3336	3369	3403	3436	3469	3502	3535	3568	3601	5' ½"
155	3163	3196	3230	3263	3296	3329	3362	3395	3428	3461	3494	3527	3560	3593	3626	
156	3189	3222	3255	3288	3322	3355	3388	3421	3454	3487	3520	3553	3586	3619	3652	5' 1 ½"
157	3215	3248	3282	3315	3348	3381	3414	3447	3480	3513	3546	3579	3612	3645	3678	
158	3242	3275	3308	3341	3374	3407	3440	3473	3507	3540	3573	3606	3639	3672	3705	5' 2"
159	3269	3302	3335	3368	3401	3434	3467	3500	3533	3566	3600	3633	3666	3699	3732	
160	3296	3329	3362	3395	3428	3461	3494	3527	3561	3594	3627	3660	3693	3726	3759	5' 3"
161	3323	3356	3390	3423	3456	3489	3522	3555	3588	3621	3654	3687	3720	3753	3787	
162	3351	3384	3417	3451	3484	3517	3550	3583	3616	3649	3682	3715	3748	3781	3814	5' 4"
163	3379	3413	3446	3479	3512	3545	3578	3611	3644	3677	3710	3743	3776	3810	3843	
164	3408	3441	3474	3507	3540	3573	3607	3640	3673	3706	3739	3772	3805	3838	3871	5' 4 ½"
165	3437	3470	3503	3536	3569	3602	3635	3668	3702	3735	3768	3801	3834	3867	3900	
166	3466	3499	3532	3565	3599	3632	3665	3698	3731	3764	3797	3830	3863	3896	3929	5' 5"
167	3496	3529	3562	3595	3628	3661	3694	3727	3760	3794	3827	3860	3893	3926	3959	
168 or more	More than 3500 ml															5' 6" or more
	7st 12	8st	8st 3	8st 5	8st 7	8st 9	8st 11	9st	9st 2	9st 4	9st 6	9st 8	9st 11	9st 13	10st 1	
	110lb	112lb	115lb	117lb	119lb	121lb	123lb	126lb	128lb	130lb	132lb	134lb	137lb	139lb	141lb	

Appendix 2 - Table of Immunisations

Diseases Protected against	Comments and example trade names of adult preparations	
Anthrax	Rarely given available only through Public Health England	<u>Non-Live</u>
Cholera	Oral vaccine. Dukoral [®]	<u>Non-Live</u>
COVID-19 (SARS-CoV-2)	All COVID-19 vaccines licenced in the UK are Non-Live	<u>Non-Live</u>
HiB	Haemophilus influenza, Menitorix [®]	<u>Non-Live</u>
Hepatitis A	May be combined with typhoid or hepatitis B <ul style="list-style-type: none"> • Hepatitis A only: Vaqta[®], Avaxim[®], Havrix[®] • Combined with typhoid: Viatim[®] • Combined with hepatitis B: Ambirix[®], Twinrix[®] 	<u>Non-Live</u>
Hepatitis B	May be combined with hepatitis A If unexposed and more than 7 days from last immunisation, accept. See: Hepatitis B – Immunisation Engerix [®] , Fendrix [®] , HBvaxPRO [®] , Ambirix [®] , Twinrix [®]	<u>Non-Live</u>
HPV	Cervarix [®] , Gardasil [®]	<u>Non-Live</u>
Influenza-intra-nasal	Live vaccine given by intra-nasal spray, age 2-18 Fluenz Tetra [®]	<u>Live</u>
Influenza, injection	Annual 'flu jab', given by injection Several preparations, updated annually	<u>Non-Live</u>
Japanese Encephalitis	Travel. Ixiaro [®]	<u>Non-Live</u>
Measles, Mumps, Rubella	MMR vaccines. M-M-RvaxPro[®], Priorix[®]	<u>Live</u>
Meningitis	Meningococcal group C: NeisVac-C [®] , Menjugate Kit [®] Meningococcal group B: Bexsero [®] , Trumenba [®] MenACWY Quadrivalent vaccine: Menveo [®] , Nimenrix [®] , MenQuadfi [®] Combined with <i>H. influenzae</i> type b (Hib): Menitorix [®]	<u>Non-Live</u>
Mpox (formerly known as Monkeypox)	During the 2022 Mpox outbreak, contacts of Mpox cases may receive Imvanex, a live attenuated non-replicating Smallpox vaccine. For donor selection purposes this should be assessed as a non-live vaccine. See DSG entry for Mpox	<u>Non-Live</u>
Pertussis	Usually pregnant women, Boostrix-IPV [®] Repavax [®]	<u>Non-Live</u>
Pneumococcal disease	Given to people with specific risks: for example, people who have had a splenectomy or people over 65. Pneumovax [®] 23	<u>Non-Live</u>
Polio, injected	Would usually be given Diphtheria, Tetanus and Polio (injection). Boostrix-IPV [®] , Revaxis [®] , Repevax [®]	<u>Non-Live</u>
Polio, oral	Not in routine use in UK. May be used abroad	<u>Live</u>
Rabies	Given to non-exposed individuals if occupation or activity has an exposure risk, or for some travellers to endemic areas: Rabipur [®] , Verorab [®]	<u>Non-Live</u>
Shingles	There are two vaccines available to prevent	

	shingles: Zostavax[®] and Shingrix[®] ; see rows below. Ensure the correct guidance is applied depending on the vaccine given. If vaccine name not certain, treat as a Live vaccine	
	<ul style="list-style-type: none"> • Zostavax[®] for shingles prevention 	<u>Live</u>
	<ul style="list-style-type: none"> • Shingrix[®] for shingles prevention 	<u>Non-Live</u>
Smallpox	Note this live vaccine requires an 8-week deferral If given, see DSG entry for Smallpox Immunisation. See also Mpox	<u>Live</u>
Tetanus	Would usually be given as Diphtheria, Tetanus and Polio. Boostrix-IPV [®] , Revaxis [®] , Repevax [®]	<u>Non-Live</u>
Tick-borne encephalitis	TicoVac [®]	<u>Non-Live</u>
Tuberculosis	BCG vaccine	<u>Live</u>
Typhoid - injected	Typhim Vi [®] Combined with hepatitis A: Viatim [®]	<u>Non-Live</u>
Typhoid - Oral	Given in capsule form Vivotif[®]	<u>Live</u>
Varicella (chickenpox)	Usually given to healthcare workers. Varilrix[®], Varivax[®]	<u>Live</u>
Yellow Fever	Stamaril	<u>Live</u>

Appendix 3 - Maximum permitted Extra Corporeal Volume for component donors

Female Donors

Height (cm)	Weight (kg)											Height
	50	55	60	65	70	75	80	85	90	95	100	
150	486	513	539	566	592	619	645	672	698	724	751	4'11"
153	498	524	551	577	604	630	657	683	710	736	763	5'
155	506	533	559	586	612	638	665	691	718	744	771	5'1"
158	519	545	572	598	625	651	677	704	730	757	783	5'2"
160	527	554	580	607	633	660	686	713	739	766	792	5'3"
163	541	567	594	620	647	673	700	726	752	779	805	5'4"
165	550	576	603	629	656	682	709	735	762	788	815	5'5"
168	564	591	617	644	670	696	723	749	776	802	829	5'6"
170	574	600	627	653	680	706	733	759	786	812	839	5'7"
173	589	615	642	668	695	721	748	774	801	827	854	5'8"
175	599	626	652	679	705	732	758	785	811	837	864	5'9"
178	615	642	668	695	721	748	774	801	827	853	880	5'10"
180	626	653	679	706	732	759	785	812	838	864	891	5'11"
183	643	670	696	723	749	775	802	828	855	881	908	6'
	7.9st	8.7st	9.5st	10.2st	11st	11.8st	12.6st	13.4st	14.2st	15st	15.7st	
	Weight											

Male Donors

Height (cm)	Weight (kg)											Height
	50	55	60	65	70	75	80	85	90	95	100	
150	552	578	604	630	655	681	707	733	758	784	810	4'11"
153	564	590	616	642	667	693	719	745	770	796	822	5'
155	573	599	624	650	676	702	727	753	779	805	830	5'1"
158	586	611	637	663	689	714	740	766	792	817	843	5'2"
160	595	620	646	672	698	723	749	775	801	826	852	5'3"
163	608	634	660	686	711	737	763	789	814	840	866	5'4"
165	618	644	669	695	721	747	772	798	824	850	875	5'5"
168	633	658	684	710	736	761	787	813	839	864	890	5'6"
170	643	668	694	720	746	771	797	823	849	874	900	5'7"
173	658	684	710	735	761	787	813	838	864	890	916	5'8"
175	669	695	720	746	772	798	823	849	875	901	926	5'9"
178	685	711	737	763	788	814	840	866	891	917	943	5'10"
180	697	722	748	774	800	825	851	877	903	928	954	5'11"
183	714	740	765	791	817	843	868	894	920	946	971	6'
	7.9st	8.7st	9.5st	10.2st	11st	11.8st	12.6st	13.4st	14.2st	15st	15.7st	
	Weight											

Appendix 4 - Management of post donation illness

This appendix gives guidance on the management of donations taken from donors who report post donation illness with a (probable) infectious cause.

The actions are based upon the nature and potential severity of the illness, relevant incubation period and the risk of the illness causing harm to a transfusion recipient.

Recipient notification and lookback/traceback investigations are outside the scope of this guidance. Please follow local policies and procedures.

Infections

Infection	Incubation	Action for donation
Bordetella Pertussis (Whooping Cough)	IP 7-10 days	Discard if within 10 days
Borrelia Burgdorferi (Lyme Disease)	IP 3-30 days	Discard up to 30 days if donor diagnosed with acute Lyme disease Chronic Lyme disease no action required
Costochondritis/ Coxsackie virus (Bornholm Disease)	IP 1-7 days	Discard up to 1 week
Chickenpox /Varicella Zoster	IP 10-21 days	Discard if within 3 weeks
COVID-19 (SARS-CoV-2)	IP 2-14 days	Discard if: <ul style="list-style-type: none"> • A SARS-CoV-2 test has been taken and COVID-19 confirmed; and • Symptoms and/or the positive test result occurred in the 48-hour period after donation. If a SARS-CoV-2 test is negative or has not been taken, refer to the relevant advice on this page for the donor's symptoms.
Coxsackie A (Hand, foot & mouth disease)	IP 3-7 days Usually Coxsackie A, but can be other enteroviruses	Discard up to 7 days
Epstein-Barr Virus (Glandular Fever)	IP 30-50 days	Discard up to 50 days
Hepatitis (acute, viral)	IP HAV 2-6 weeks	Discard up to 7 weeks
	IP HBV 6 weeks to 6 months	Discard all in date components*
	IP HCV up to 6 months	Discard all in date components*
	IP HEV 2-8 weeks	Discard up to 9 weeks
Herpes Simplex (Oral and genital)	IP 2-12 days for primary infection. Primary viraemia during IP, secondary viraemia at time of symptom	Discard up to 14 days for primary infection
	Recurrent infection	No action if recurrent lesion/s and lesions were absent or healing when donated
HIV		Discard all in-date components at any interval after donation*
HTLV		Discard all in-date components at any interval after donation
Influenza	IP 1-5 days (Influenza A)	No action unless severe systemic symptoms.
	IP 4-5 days (adenovirus)	
	Defined as fever	If present, discard up to 5 days

	/myalgia +/- cough /cold symptoms	
Legionella (Legionnaire's Disease/Pontiac Fever)	IP up to 3 weeks	Discard up to 3 weeks
Measles	IP 10-21 days	Discard up to 3 weeks
Monkeypox	21 days	Discard up to 21 days. Follow local processes for public health notification if the component has been transfused. If the donor has reported contact with Monkeypox in the 21 days before donation, place the donation on hold and seek public health advice to determine the risk.
Mumps	IP 16-18 days Primary & secondary viraemia	Discard up to 3 weeks
Mycoplasma	IP 1-4 weeks M. pneumoniae	Discard up to 4 weeks
	Mostly headache, malaise, fever, 5-10% progress to pneumonia	
Parvovirus B19 (Fifth disease, Slapped Cheek)	IP 13-20 days	Discard up to 3 weeks
Rubella(German Measles)	IP 14-21 days	Discard up to 3 weeks
TB		Discard all in-date components at any interval post donation. Look-back to relevant transfused recipients
West Nile Virus	IP 3-15 days	Discard up to 15 days

IP –Incubation Period

* HBV, HCV and HIV Seek microbiological advice regarding recall of previous donations if the donor's history and/or testing results suggest this is an acute (recent) infection

Conditions

Condition	Comments	Action for donation
Appendicitis		No action if confirmed appendicitis and asymptomatic at the time of donation.
Bornholm Disease	See Costochondritis/ Coxsackie virus	
Chest infection		No action unless systemic symptoms; if present discard up to 5 days
Common Cold		No action unless symptoms
Conjunctivitis		No action providing well on the day
Diarrhea & vomiting	Causes may include	Discard up to 14
	Salmonella (IP 12-72 hrs)	If this is an episode of food poisoning which occurred after the donation, no action required
	Shigella (IP 1-7 days)	
	Campylobacter (IP 1-11 days)	
	Rotavirus (IP 24-72 hrs)	
	Norovirus (IP 1-2 days)	
	Cryptosporidium (IP 2-5 days)	
	Yersinia (IP 4 days)	
	With all the above likely to be significant bacteremia or viraemia	

	Staphylococcal, Clostridium and B. cereus food poisoning is all toxin induced	
Fifth Disease, Slapped Cheeks	See Parvovirus B19	
German Measles	See Rubella above	
Glandular Fever /Kissing Disease	See Epstein-Barr Virus	
Hand, Foot and Mouth Disease	See Coxsackie A	
Jaundice	Assess whether infective cause possible	Discard all in-date components after any notification if infection is a possible cause: <ul style="list-style-type: none"> • See specific entry if infective cause identified • Discard not required if a non-infective cause has been identified (e.g. drug reaction)
Legionnaire's Disease /Pontiac Fever	See Legionella	
Lyme Disease	Refer Borrelia Burgdorferi	
Malaria	Any disclosure of illness or risk after donation	Follow local policies and procedures
Shingles (Herpes Zoster)	Possible viraemia for 48 hours from symptoms and/or rash	Discard if rash or any symptoms develop within 48 hours. Symptoms include tingling of skin, pain or eruption of vesicles
Skin disease: Cellulitis /erysipelas	Streptococcus Pyogenes	Discard up to 1 week
Skin disease: Impetigo	Group A Streptococcus Staphylococcus Aureus IP 3-5 days	No action if no systemic symptoms; if present, discard up to 1 week
Sore throat	May include: Rhinovirus Group A strep (IP 2-4 days) EBV (IP 0-50 days)	If a sore throat is accompanied by simple cold symptoms and no systemic symptoms, no action is required Systemic symptoms include malaise, myalgia, fever, headache. If systemic symptoms, discard up to 1 week If glandular fever, discard up to 50 days
Transmissible Spongiform Encephalopathy (Prion Disease)	If informed of a possible or confirmed case of prion associated disease, recall (do not discard) any in date components.	Follow local policies and procedures.
UTI	Symptomatic at donation Asymptomatic at donation	Discard No action unless systemic symptoms when discard up to 5 days Systemic symptoms include malaise, myalgia, fever and headache.
Whooping cough	See Bordetella Pertussis	

IP – Incubation Period

Drugs and Platelet Donation

Principles

Some drugs can alter platelet function for several hours or days after they are taken.

If a donor is otherwise eligible to donate but has taken a drug affecting platelets in the 48 hours prior to donation, their donation should not be used to manufacture blood components that are intended to treat thrombocytopenia or platelet dysfunction; this includes apheresis platelets, pooled platelets and some whole blood components.

This rule applies to drugs taken systemically, i.e. medication that is taken orally or rectally, or medication that is taken by injection.

This rule does not apply to creams or gels applied topically to the skin. Donors who have used topical medications which are listed in this index, can be accepted for platelets immediately.

Drugs which affect platelet function are usually taken as painkillers and anti-inflammatory medications. Some can be purchased directly from shops or pharmacies without a prescription. They may be sold under their generic drug name (e.g. aspirin, ibuprofen) or they may be sold under a brand name.

This entry lists Non-Steroidal Anti-Inflammatory Drugs (NSAIDs) and other painkillers which affect platelet function. It does not include every medication which could affect platelets. Some drugs, such as clopidogrel, have not been included because a donor taking them would not be eligible to donate anyway.

How to use this index

If a donor reports taking anti-inflammatory or pain-killing medication, but has been assessed as eligible to donate, their medication must be checked against this entry.

The entry has been redrafted to list only generic drug names and no longer includes brand names. This is because some brand names are used for several formulations, not all of which contain the same active drugs. It is important that staff confirm the exact drug taken by a donor.

Up-to-date product information for branded medications available in the UK can be found online at www.gov.uk/government/organisations/medicines-and-healthcare-products-regulatory-agency, bnf.nice.org.uk and www.medicines.org.uk/emc. If there is doubt about the active drug in any individual branded product, staff should apply the 48 hour rule for platelet production.

Post donation information

Donors may contact the blood service to report they had taken NSAIDs before their donation. If any blood components containing therapeutic platelets have been manufactured from the donor's blood, these should be discarded. If such components have been transfused, a DCSO should assess the need to notify the treating clinician.

Drug Index

Drug Index

A

Aceclofenac

Acemetacin

Aspirin

Azapropazone

C

Celecoxib

D

Dexketoprofen

Diclofenac potassium

Diclofenac sodium

E

Etodolac

Etoricoxib

F

Felbinac

Fenbufen

Fenoprofen

Flurbiprofen

I

Ibuprofen

Indometacin

K

Ketoprofen

Ketorolac trometamol

M

Mefenamic Acid

Meloxicam

N

Nabumetone

Naproxen

P

Parecoxib

Piroxicam

S

Sulindac

T

Tenoxicam

Tiaprofenic Acid

Tolfenamic Acid